

Current understandings on dust formation in supernovae

Takaya Nozawa

(NAOJ, Division of theoretical astronomy)

Main collaborators:

Masaomi Tanaka (NAOJ, theory)

Keiichi Maeda (Kyoto University)

Takashi Kozasa, Asao Habe (Hokkaido University)

Hideyuki Umeda (University of Tokyo)

Ken'ichi Nomoto (Kavli IPMU)


0-1. My research interests

- **Formation of dust in the ejecta of supernovae**
- **Dynamics, destruction, and heating of dust grains in high-velocity shock waves**
- **Evolution of grain size distribution in galaxies and interstellar extinction curves**
- **Origin of dust grains in the early universe and the roles of dust in star formation**
- **Formation of dust in stellar winds and mass-loss history at late phases of stellar evolution**

0-2. Introduction

SNe are important sources of interstellar dust?

- abundant metal (metal : $N > 5$)
- low temperature ($T < \sim 2000$ K)
- high density ($n > \sim 10^6$ cm⁻³)



mass-loss winds
of AGB stars
expanding ejecta
of supernovae

- huge amounts of dust grains ($> 10^8 M_{\text{sun}}$) are detected in host galaxies of quasars at redshift $z > 5$
 - **0.1 M_{sun} of dust per SN** is needed to explain such massive dust at high- z (e.g. Dwek et al. 2007)

contribution of dust mass from AGB stars and SNe


$$n(\text{AGB stars}) / n(\text{SNe}) \sim 10\text{-}20$$

$M_{\text{dust}} = 0.01\text{-}0.05 M_{\text{sun}}$ per AGB (Zhukovska & Gail 2008)

$M_{\text{dust}} = 0.1\text{-}1.0 M_{\text{sun}}$ per SN (e.g., Nozawa et al. 2003, 2007)


Dust Formation in the ejecta of SNe

at ~1 days




1. Observations of Dust Formation in SNe (and SNRs)

1-1. Summary of observed dust mass in CCSNe






Far-IR to sub-mm observations are essential for revealing the mass of dust grains produced in the ejecta of SNe

1-1. Summary of observed dust mass in CCSNe





Far-IR to sub-mm observations are essential for revealing the mass of dust grains produced in the ejecta of SNe

1-2. Herschel detects cool dust in SN 1987A



~180K
~10⁻⁵ Msun



on 4 Oct 2003
Gemini T-ReCS
($\lambda = 10.36 \mu\text{m}$)
2 pixels : 0.18"
(Bouchet+04)


Herschel detects cool (~20K) dust of ~0.4-0.7 Msun toward SN 1987A!

1-3. Resolving cool dust in SN 87A with ALMA


ALMA Cycle 0 Proposal
 'Detecting cool dust in SN1987A'
 (TN, Tanaka, et al.)






Band 7 (850 μm)



Band 9 (450 μm)



CASA simulation
 with extended
 config. (4 hrs)



0.1 Msun of silicate
→ 5 σ detection at Band 9 !!

1-4. Successful ALMA proposals for SN 1987A

~~2011.0.00241.S~~

PI	Exec	Country	Institute
Mozzari, Takaya	EA	Japan	The University of Tokyo
COI			
Tanaka, Masahiro	EA	Japan	The University of Tokyo
Moriya, Takashi	EA	Japan	University of Tokyo
Minamidani, Tetsuhiko	EA	Japan	Hokkaido University
Kozasa, Takashi	EA	Japan	Hokkaido University

This proposal was ranked in the highest priority !!

Our proposal was not executed

Band 9 extended configuration


2011.0.00273.S

PI	Exec	Country	Institute
Indebetouw, Remy	NA	United States	Virginia, University of
COI			
McGray, Richard	NA	United States	Colorado at Boulder, Univ of
Matsuura, Mikako	EU	United Kingdom	London, University of
Andjelic, Milica	OTHER	Serbia	Belgrade, University of
Arbutins, Bojan	OTHER	Serbia	Belgrade, University of
Baes, Maarten	EU	Belgium	Ghent University
Bolatto, Alberto	NA	United States	Maryland, University of
Burrows, David	NA	United States	Pennsylvania State University
Chevalier, Roger	NA	United States	Virginia, University of
Gaensler, Bryan	OTHER	Australia	Sydney, University of
Long, Knox	NA	United States	Space Telescope Science Institute
Lundqvist, Peter	EU	Sweden	Stockholm University
Mebner, Margaret	NA	United States	Space Telescope Science Institute
Marcalde, Jon	EU	Spain	Valencia, University of
Marti-Vidal, Ivan	EU	Germany	Max-Planck-Institute for Radio Astronomy
OTSUKA, Masaoaki	EA/NA	Taiwan	Academia Sinica
Sandstrom, Karin	EU	Germany	Max-Planck-Institute for Astronomy
Sonneborn, George	NA	United States	National Aeronautics and Space Administration
Steveley-Smith, Lister	OTHER	Australia	International Centre for Radio Astronomy Research
van Leeu, Jacco	EU	United Kingdom	Keele University
Urosavic, Dejan	OTHER	Serbia	Belgrade, University of
Vlahakis, Catherine	CL	Chile	Chile, University of

Band 3, 6, 7, 9 compact configuration


Wesson, Roger	EU	United Kingdom	London, University of
Dwek, Eli	NA	United States	National Aeronautics and Space Administration
Bouchet, Patrice	EU	France	CEA Saclay
Lakicevic, Masa	EU	Germany	European Southern Observatory
Potter, Toby	OTHER	Australia	International Centre for Radio Astronomy Research

1-5. ALMA reveals dust formed in SN 1987A




blue: H α
 green: 1.6 μ m
 red: CO(2-1)
 → >0.1 M \odot of CO

SED of 25-years old SN 1987A



Indebetouw+2014




ALMA spatially resolves cool (~20K) dust of ~0.5 M \odot formed in the ejecta of SN 1987A
 → SNe could be production factories of dust grains

1-6. Possible target: SNR 1E0102-72.3 in SMC


young (compact) supernova remnants in LMC/SMC !

▪ SNR 1E0102-72.3

- O-rich (Type Ib?)
- age : ~1000 yr
- $M_{\text{warm}} \sim 10^{-3} M_{\text{sun}}$
- $M_{\text{cool}} \sim ???$



Stanimirovic+05




Gaetz+00


▪ Cassiopeia A

- O-rich (Type IIb)
- age : 330 yr
- $M_{\text{warm}} < 10^{-2} M_{\text{sun}}$
- $M_{\text{cool}} \sim 0.07 M_{\text{sun}}$

Cas A model (Nozawa+10)




1-7. SNR 1E0102-72.3 is too extended!





1-8. Flux estimates necessary for detection

**1987A
Band 9**




**1E0102
Band 9**



**full operation
1 sigma, 1 hr**

**effective flux
(flux per beam)**




1-9. Summary of observations of SN dust

- **ALMA confirmed the presence of huge amounts of newly formed dust in the ejecta of SN 1987A**
 - heavy elements (Si, Mg, Fe, C) are locked up in dust grains
 - SNe are main production factories of dust grains
 - **It seems too hard to detect thermal emission from cool dust in any other SNe/SNRs with ALMA**
 - young SNRs in LMC/SMC are too extended
 - **A part of dust grains formed in the SN ejecta will be destroyed by the reverse shocks**
- what fraction of dust grains can survive to be injected?**
- destruction efficiency depends on dust composition and size
 - infrared to submm observations have few information on the the composition and size distribution of dust

Evolution of dust in SN remnants

$$T = (1-2) \times 10^4 \text{ K}$$
$$n_{\text{H},0} = 0.1-1 \text{ cm}^{-3}$$





2. Theoretical Studies on Dust Formation in SNe

2-1. Dust formation in primordial SNe

Nozawa+2003, ApJ, 598, 785



Population III SNe model (Umeda & Nomoto 2002)

- SNe II : $M_{\text{ZAMS}} = 13, 20, 25, 30 M_{\text{sun}}$ ($E_{51}=1$)
- PISNe : $M_{\text{ZAMS}} = 170 M_{\text{sun}}$ ($E_{51}=20$), $200 M_{\text{sun}}$ ($E_{51}=28$)




- nucleation and grain growth theory (Kozasa & Hasegawa 1987)
- no mixing of elements within the He-core
- complete formation of CO and SiO

2-2. Dust formed in Type II-P SNe





C / O < 1 → all C atoms are locked up in CO
C / O > 1 → all O atoms are locked up in CO



- a variety of grain species can condense according to elemental composition in each layer
- condensation time: **300-600d** after explosion
- average grain radii: **>~0.01 μm**



2-3. Size distribution of newly formed dust



- C, SiO₂, and Fe grains have lognormal-like size distribution, while the other grains have power-law size distribution
- The composition and size distribution of dust formed are almost independent of types of supernova

average grain radius is smaller for PISNe than SNe II-P


2-4. Total mass of dust formed in the ejecta



- Total mass of dust is higher for a higher progenitor mass (MZAMS)
 - SNe II : $m_{\text{dust}} = 0.1\text{-}1.5 M_{\text{sun}}$, $m_{\text{dust}} / m_{\text{metal}} = 0.2\text{-}0.3$
 - PISNe : $m_{\text{dust}} = 10\text{-}30 M_{\text{sun}}$, $m_{\text{dust}} / m_{\text{metal}} = 0.3\text{-}0.4$
- almost all Fe, Mg, and Si are locked up in dust grains, while most of C and O remain in the gas-phase (such as CO)
 - dust-to-metal mass ratio is not high for SNe II

2-5. Temperature and density of gas in SNRs

Nozawa+07, ApJ, 666, 955



Model : $M_{\text{pr}} = 20 M_{\text{sun}}$ ($E_{51} = 1$)
 $n_{\text{H},0} = 1 \text{ cm}^{-3}$

Downward-pointing arrows:
forward shock in upper panel
reverse shock in lower panel

The temperature of the gas swept up by the shocks


→ $10^6 - 10^8 \text{ K}$



Dust grains residing in the shocked hot gas are eroded by sputtering

2-6. Evolution of dust in SNRs

Nozawa+07, ApJ, 666, 955



Model : $M_{pr} = 20 M_{sun}$ ($E_{51}=1$)
 $n_{H,0} = 1 \text{ cm}^{-3}$

Dust grains in the He core collide with reverse shock at $(3-13) \times 10^3 \text{ yr}$

The evolution of dust heavily depends on the initial radius and composition




$a_{ini} = 0.01 \mu\text{m}$ (dotted lines)
→ completely destroyed

$a_{ini} = 0.1 \mu\text{m}$ (solid lines)
→ trapped in the shell

$a_{ini} = 1 \mu\text{m}$ (dashed lines)
→ injected into the ISM

2-7. Dust mass and size ejected from SN II-P

Nozawa+07, ApJ, 666, 955



total mass of dust surviving the destruction in Type II SNRs;
0.07-0.8 Msun ($n_{H,0} = 0.1-1 \text{ cm}^{-3}$)

size distribution of dust after the shock-destruction is dominated by large grains ($> 0.01 \mu\text{m}$)

2-8. Summary of dust production in SNe

- Various grain species can condense in the ejecta
 - almost all Fe, Mg, and Si are locked up in grains
- The fate of newly formed dust within SNRs strongly depends on the initial radii and compositions
- The size distribution of dust surviving the destruction in SNRs is weighted to relatively large size ($> 0.01 \mu\text{m}$).
- The total mass of dust injected into the ISM decreases with increasing the ambient gas density
 - for $n_{\text{H},0} = 0.1\text{-}1 \text{ cm}^{-3}$
 - SNe II-P → $M_{\text{dust}} = 0.1\text{-}0.8 M_{\text{sun}}$
 - making significant contribution to dust budget