

Properties of Dust Accounting for Extinction Laws toward Type Ia Supernovae

(Nozawa, T. 2016, Planetary and Space Science, 133, 36)


Takaya Nozawa

(National Astronomical Observatory of Japan)

Special thanks:

Keiichi Maeda, Takashi Nagao, Ryosuke S. Asano,
Hiroyuki Hirashita, Tsutomu, T. Takeuchi

1-1. Extinction curves in the Milky Way




in the MW
 $R_V = 2.2\text{—}6$
 $R_{V,\text{ave}} = 3.1$

○ CCM relation (Cardelli, Clayton, Mathis 1989)

→ describes the variety of extinction curves in the MW through total-to-selective extinction ratio $R_V = A_V/E(B-V) = A_V/(A_B-A_V)$


1-2. Deriving extinction curves in the MW

Pair method



Light sources: OB stars (or RGs)

- luminous ($\sim 10^5 L_{\text{sun}}$)
→ we can see a large volume
- UV (or IR) bright
→ variation of extinction curves at UV wavelengths



1-3. Applicability of OB stars in pair method


- OB stars can be used only for MW, LMC, SMC, (M31)

→ too faint to be observed in external galaxies

Extinction curves in external galaxies are poorly known



- QSOs and GRB afterglows

→ good light sources to extract the extinction curves at $z = 0-6.5$



However ...

- intrinsic spectral energy distributions are not always established
- local dust may also contribute the observed extinction



Gallerani+2010



2-1. Type Ia SNe as standard light sources

○ Type Ia supernovae (SNe Ia)


- thermonuclear explosion of a WD
- highly luminous ($L_{\text{peak}} \sim 3 \times 10^9 L_{\text{sun}}$)
 - homogeneous peak luminosity
 - used as standard candles
- intrinsic opt/IR spectral established
 - SN 2011fe as an unreddened template
- discovered in all types of galaxies
 - star-forming, elliptical, spiral etc...



good targets to probe the extinction
(dust) properties in external galaxies




2-2. Extinction laws measured for SNe Ia




Individual (nearby) SNe Ia

- highly reddened ($A_v > \sim 2.0$ mag)
- reliable extinction curves
- described by CCM curves with $R_v = 1.5 - 2.0$



Rv values measured for SNe Ia are unusually low ($R_v \sim 1.0-2.5$)

2-3. How peculiar is SNe Ia extinction curves?



○ CCM relation

(Cardelli, Clayton, Mathis 1989)

R_V : total-to-selective
 extinction ratio

$$R_V = A_V / E(B - V)$$

$$= A_V / (A_B - A_V)$$



$$A_\lambda / A_V = a(x) + b(x) / R_V$$

where $x = 1 / \lambda$

- **steeper** extinction curve (**lower** R_V) → **smaller** grains
- **flatter** extinction curve (**higher** R_V) → **larger** grains


3-1. What is the cause for unusually low R_v ?

(1) Unique properties or effects of circumstellar dust around SN Ia

(2) Peculiar properties of interstellar dust in host galaxies of SN Ia

(3) Non-standard properties of extragalactic dust → unlikely

(4) Something is wrong in deriving the extinction laws toward SNe Ia



3-2. Multiple scattering scenario by local dust



○ Multiple scattering scenario

- multiple scattering by circumstellar dust steepens extinction curves

(Wang 2005; Goobar 2008; Amanullah & Goobar 2011)

circumstellar
dust shell of

$$\tau_V \sim 1$$




radiative transfer calculations

3-3. Concern for multiple scattering scenario

Goobar (2008)

- LMC dust (WD01) : $\omega(B)/\omega(V) < 1 \rightarrow$ **steepen**
- MW dust (WD01) : $\omega(B)/\omega(V) < 1 \rightarrow$ **steepen**


WD01 dust model: graphite, silicate, and PAH



Nagao, Maeda, TN (2016)

- LMC dust (MRN77) : $\omega(B)/\omega(V) > 1 \rightarrow$ **flatten**
- MW dust (Pei92) : $\omega(B)/\omega(V) > 1 \rightarrow$ **flatten**

classical dust model: graphite and silicate



Nagao, Maeda, TN (2016)

highly model-dependent !!

Multiple scattering does not always steepen the extinction curves

3-4. Problem on multiple-scattering scenario


No excess of NIR emission

- conservative upper limits of optical depth in B band is $\tau_B < \sim 0.1$
- no massive dusty envelope around SNe Ia


Maeda, TN, Nagao,
Motohara (2015)

circumstellar
dust shell of

$$\tau_V \sim 1$$




If there is a thick dust shell,
we must detect thermal dust
emission as infrared echoes



3-5. What is the cause for unusually low R_V ?

- (1) Unique properties or effects of circumstellar dust → unlikely
- (2) Peculiar properties of interstellar dust in host galaxies of SN Ia
- (3) Non-standard properties of extragalactic dust → unlikely
- (4) Something is wrong in deriving the extinction laws toward SNe Ia



4-1. Fitting to CCM curves with $R_V = 1-2$

What properties of dust cause steep extinction curves?

Data on extinction curves to be fitted

CCM extinction curves with $R_V = 2.0, 1.5, 1.0$ at representative photometric bands


Interstellar dust model

(e.g., Mathis+1977, Draine & Lee 1984)

- graphite & astronomical silicate
- power-law grain size distribution
- **Model 1 (simplest model)**
same size distribution with $q = -3.5$ and $a_{\min} = 0.005 \mu\text{m}$ for two grain species




parameters:

- a_{\max} (upper cutoff radius)
- f_{gs} (graphite-to-silicate mass ratio)



black dots: data of extinction $A_{\lambda i}/A_V$ derived from the CCM formula at photometric bands

4-2. Results of fitting calculations



- **steep extinction curves with $R_V=1-2$ can be described by the power-law grain model**




- $f_{\text{gs}} = 0.45-0.6$

→ $M_{\text{gra}}/M_{\text{total}} = 0.3-0.4$

cf. $f_{\text{gs}} = 0.3-0.7$ in the MW


(Nozawa & Fukugita 2013)

4-3. Implication from dust evolution model



Asano+2013, 2014; Nozawa+2015

Dust evolution model predicts that there may be a phase of a steep extinction curve (but, the calculated R_V is not necessarily low)



**$R_V = 3.8 \rightarrow$
grain size is too small**

4-4. Challenge to dust evolution model

SN 2011iv in NGC1404
© S. Kohle



SN 2014J in M82
© NASA/ESA Foley&McCully




SNe Ia appear in any type of galaxies!

SN 2006X



M100 Spiral Galaxy
© 2006 John Chumack

SN 2006X in M100




SN 2003cg in NGC3169
©ESO, Igor Chkalin



5. Summary of this talk

1) Many studies suggest that R_V values toward SNe Ia are very low ($R_V = 1-2$), compared with $R_V = 3.1$ in the MW



2) Non-detection of IR echoes toward SNe Ia indicates that the low R_V is not caused by circumstellar dust but by interstellar dust in the host galaxies

3) The CCM curves with $R_V = 1-2$ can be nicely fitted by power-law grain size distributions with $a_{max} = 0.05-0.15 \mu m$

