

*This copy is for your personal, non-commercial use only.*

**If you wish to distribute this article to others**, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

**Permission to republish or repurpose articles or portions of articles** can be obtained by following the guidelines [here](#).

**The following resources related to this article are available online at [www.sciencemag.org](http://www.sciencemag.org) (this information is current as of September 14, 2011 ):**

**Updated information and services**, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/333/6047/1258.full.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/333/6047/1258.full.html#related>

This article **cites 50 articles**, 1 of which can be accessed free:

<http://www.sciencemag.org/content/333/6047/1258.full.html#ref-list-1>

This article has been **cited by** 1 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/content/333/6047/1258.full.html#related-urls>

This article appears in the following **subject collections**:

Astronomy

<http://www.sciencemag.org/cgi/collection/astronomy>

# Herschel Detects a Massive Dust Reservoir in Supernova 1987A

M. Matsuura,<sup>1,2\*</sup> E. Dwek,<sup>3</sup> M. Meixner,<sup>4</sup> M. Otsuka,<sup>4</sup> B. Babler,<sup>5</sup> M. J. Barlow,<sup>1</sup> J. Roman-Duval,<sup>4</sup> C. Engelbracht,<sup>6</sup> K. Sandstrom,<sup>7</sup> M. Lakićević,<sup>8,9</sup> J. Th. van Loon,<sup>8</sup> G. Sonneborn,<sup>3</sup> G. C. Clayton,<sup>10</sup> K. S. Long,<sup>4</sup> P. Lundqvist,<sup>11</sup> T. Nozawa,<sup>12</sup> K. D. Gordon,<sup>4</sup> S. Hony,<sup>13</sup> P. Panuzzo,<sup>13</sup> K. Okumura,<sup>13</sup> K. A. Misselt,<sup>6</sup> E. Montiel,<sup>6</sup> M. Sauvage<sup>13</sup>

We report far-infrared and submillimeter observations of supernova 1987A, the star whose explosion was observed on 23 February 1987 in the Large Magellanic Cloud, a galaxy located 160,000 light years away. The observations reveal the presence of a population of cold dust grains radiating with a temperature of about 17 to 23 kelvin at a rate of about 220 times the luminosity of the Sun. The intensity and spectral energy distribution of the emission suggest a dust mass of about 0.4 to 0.7 times the mass of the Sun. The radiation must originate from the supernova ejecta and requires the efficient precipitation of all refractory material into dust. Our observations imply that supernovae can produce the large dust masses detected in young galaxies at very high redshifts.

Supernovae produce most of the heavy elements found in the universe and disperse them into their surrounding galactic environment. They chemically enrich the material from which new generations of stars and planets are formed. The explosion of supernova (SN) 1987A was first observed from Earth on 23 February 1987 in the Large Magellanic Cloud (LMC), only 50 kpc away. Because of its proximity, it has been possible to witness its evolution from explosion to remnant. SN 1987A has thus become one of the most extensively studied extragalactic objects, with ground, airborne, and space observatories covering a wide range of the electromagnetic spectrum. Here, we report far-infrared and submillimeter (submm) observations of SN 1987A.

## Previous observations of Supernova 1987A.

The detection of neutrinos from SN 1987A confirmed that the event marked the explosive death

of a massive star (1, 2). The intensity and evolution of its ultraviolet (UV) and optical light curves showed that the SN's luminosity was powered by the radioactive decays of <sup>56</sup>Ni and <sup>56</sup>Co, yielding 0.06 times the mass of the Sun ( $M_{\odot}$ ) of <sup>56</sup>Fe (3). The  $\gamma$ -rays and x-rays from radioactive nuclei emerged only 4 months after the explosion (4, 5), suggesting that the ejecta were clumpy (6). The large width of near- and mid-infrared spectral lines of heavy elements (7) indicated that the newly synthesized material from the stellar core had undergone substantial mixing into the outer ejecta of the stellar explosion (6). SN 1987A was classified as a type II event on the basis of the detection of hydrogen in its optical spectrum (8).

Examination of plates of the region obtained before the SN explosion allowed the detection of its progenitor, a blue supergiant (Sk-69 202) that was believed to have had an initial mass of 18 to 20  $M_{\odot}$  (8). The progenitor underwent extensive mass loss during its post main-sequence evolution (8). The detection and evolution of UV lines from highly ionized gas revealed previously ejected material that was flash ionized by the x-ray and UV radiation released when the outwardly expanding shock generated by the collapse of the stellar core broke out through the surface of the progenitor star (9). The presence of progenitor material was later confirmed by Hubble Space Telescope observations, which revealed a thin elliptical ring (1.6" by 1.2") (10) containing many knots within it (11). The equatorial ring is part of a system of three rings forming an hourglass-like configuration centered on the explosion. The mass of the ring is estimated to be  $\sim 6 \times 10^{-2} M_{\odot}$  (12).

Mid-infrared spectral and photometric observations of SN 1987A obtained with the Kuiper Airborne Observatory (13) revealed the presence of newly formed dust in the SN, providing direct evidence for the formation of grains in the cooling ejecta of a supernova. The dust condensation process started around day 450 to 600 after the

explosion was observed (14, 15); it is thought that dust had condensed in the clumps within the ejecta (16). The dust mass was estimated to be at least  $\sim 10^{-4} M_{\odot}$  (17).

Several years after the explosion was observed, SN 1987A had evolved from a SN remnant radiating as a result of radioactive decays, mainly <sup>56</sup>Co and <sup>44</sup>Ti (3), into a remnant where the interaction of the SN blast wave with the circumstellar medium is causing the ring to brighten up. This interaction, dominated by the collision between the blast wave and the equatorial ring, manifested through the appearance of UV-optical hot spots strung like beads around the ring (18). Similar ring structures were also observed at x-ray wavelengths and in the mid-infrared (19).

Subsequent spectral observations with the Spitzer Space Telescope on days 6200 to 8000 (20) revealed strong infrared (6 to 30  $\mu\text{m}$ ) emission arising from about  $10^{-6} M_{\odot}$  of silicate dust in the inner ring that had been collisionally heated to a temperature of 180 K by the shocked x-ray emitting gas. The dust in the ring was ejected during the pre-SN evolution of the progenitor star.

**Observations and results.** Using the Herschel Space Observatory's (21) imaging instruments, the Photodetector Array Camera and Spectrometer (PACS) (22) and the Spectral and Photometric Imaging Receiver (SPIRE) (23), we observed SN 1987A as part of the Herschel Inventory of the Agents of Galaxy Evolution in the Magellanic Clouds open time key program (HERITAGE) (24). The observing survey covered five bands, PACS 100- and 160- $\mu\text{m}$  and SPIRE 250-, 350-, and 500- $\mu\text{m}$ , with angular resolutions of 6.69" by 6.89", 10.65" by 12.13", 18.2", 24.9", and 36.3", respectively. SN 1987A was observed on 30 April and 5 August 2010 (days 8467 and 8564 after the explosion was observed).

The PACS and SPIRE images of the combined two epochs (Fig. 1) (25) reveal SN 1987A as a faint but detectable point source coincident with the very-long-baseline interferometry-measured coordinates of SN 1987A: right ascension, 5h 35m 27.990s; declination,  $-69^{\circ} 16' 11.110''$  (J2000) (26). Spatial comparison of these images with previous Spitzer detections with the multiband imaging photometer (MIPS) at 24  $\mu\text{m}$  and the infrared array camera (IRAC) at 8  $\mu\text{m}$  (20) confirms the location of SN 1987A with respect to nearby interstellar features. In the PACS 100- and 160- $\mu\text{m}$  images, SN 1987A is clearly detected as a point source that is well separated from the nearby interstellar medium (ISM) dust. In the SPIRE bands, the 250- $\mu\text{m}$  image shows SN 1987A as a point source, but the SN begins to merge with the nearby ISM dust emission at 350  $\mu\text{m}$  and blends completely at 500  $\mu\text{m}$ , owing to the increasing beam size at longer wavelengths. The Hubble and Spitzer Space Telescopes and Herschel Space Observatory images (Fig. 1, A and C to H) show that SN 1987A is located in a region with relatively low amounts of ISM gas and dust.

<sup>1</sup>Astrophysics Group, Department of Physics and Astronomy, University College London, Gower Street, London WC1E 6BT, UK. <sup>2</sup>Mullard Space Science Laboratory, University College London, Holmbury St. Mary, Dorking, Surrey RH5 6NT, UK. <sup>3</sup>Observational Cosmology Laboratory, Code 665, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA. <sup>4</sup>Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA. <sup>5</sup>Department of Astronomy, 475 North Charter St., University of Wisconsin, Madison, WI 53706, USA. <sup>6</sup>Steward Observatory, University of Arizona, Tucson, AZ 85721, USA. <sup>7</sup>Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany. <sup>8</sup>Astrophysics Group, Lennard-Jones Laboratories, Keele University, ST5 5BG, UK. <sup>9</sup>European Southern Observatory, Karl Schwarzschild Straße 2, D-85748 Garching bei München, Germany. <sup>10</sup>Louisiana State University, Department of Physics and Astronomy, 233-A Nicholson Hall, Tower Drive, Baton Rouge, LA 70803-4001, USA. <sup>11</sup>Department of Astronomy, The Oskar Klein Centre, AlbaNova University Center, Stockholm University, SE-106 91 Stockholm, Sweden. <sup>12</sup>Institute for the Physics and Mathematics of the Universe, University of Tokyo, Kashiwa, Chiba 277-8583, Japan. <sup>13</sup>Commissariat à l'Énergie Atomique et aux Énergies Alternatives, Laboratoire Astrophysique, Instrumentation et Modélisation, Irfu/SAP, Orme des Merisiers, F-91191 Gif-sur-Yvette, France.

\*To whom correspondence should be addressed. E-mail: mikako@star.ucl.ac.uk

We obtained photometry from the processed Herschel images (Table 1) (27), assuming that SN 1987A is unresolved; the fluxes are plotted in Fig. 2A. There are two peaks in the spectral energy distribution (SED), one near 20  $\mu\text{m}$  and the other at 150 to 200  $\mu\text{m}$  (Fig. 2A). The total luminosity from 100 to 500  $\mu\text{m}$  is  $\sim 220$  times the luminosity of the Sun ( $L_{\odot}$ ). The mid-infrared (5 to 40  $\mu\text{m}$ ) luminosity is  $1200 L_{\odot}$  (epoch of 7983 days) (20), and the x-ray luminosity is  $\sim 500 L_{\odot}$  (8012 days) (28). Thus, the far-infrared/submm

emission that we detected is an important component of this SN's SED.

**The origin of the far-infrared/submm emission.** The smooth blackbody-like shape of the far-infrared/submm SED (Fig. 2A) suggests continuous emission from dust. However, synchrotron emission and line emission from the same gas that produces the UV, optical, and near-infrared lines are also possible.

Line emission can arise from the supernova ejecta and the circumstellar rings. The rings were


first flash-ionized by the supernova and have since recombined and cooled (12). To estimate the contribution from ionized lines from the inner ring, we extended the photoionization code of (12) to include far-infrared/submm lines. The expected line intensities of the strongest ionized lines are  $0.5 \times 10^{-15}$ ,  $1.2 \times 10^{-15}$ ,  $0.1 \times 10^{-15}$ , and  $0.2 \times 10^{-15} \text{ erg s}^{-1} \text{ cm}^{-2}$  for the 88- $\mu\text{m}$  ([OIII]), 122- $\mu\text{m}$  ([NII]), 158- $\mu\text{m}$  ([CII]), and 205- $\mu\text{m}$  ([NII]) lines, respectively. These line fluxes are equivalent to a 0.2% contribution to the total Herschel in-band fluxes; hence, photoionized lines make a negligible contribution to the far-infrared/submm fluxes. Another possible contribution to the line emission may come from the radiatively cooling gas shocked by the ejecta/ring interaction (11). However, it is unlikely that this contribution could be orders of magnitude higher than those from photoionized gas because the forbidden lines are collisionally de-excited in the high-density shocked gas.

Continuum synchrotron radiation can also contribute to the emission in the Herschel bands. The synchrotron radiation has been measured at radio frequencies up to day 8014 after the explosion (26, 29) (Fig. 2A); we extrapolated the fluxes to far-infrared wavelengths using a power law. The predicted Herschel flux densities from synchrotron radiation are two orders of magnitude lower than those we measured (Fig. 2A), implying that the emission observed by Herschel is mostly due to the continuum emission from dust.

**Dust mass.** Using the flux densities and their uncertainties, we calculated the dust temperature and mass needed to account for the observations (Table 2 and Fig. 2B) (30).

We explored four possible origins for the dust: progenitor dust, ambient ISM dust swept up by SN shocks, a light echo from ISM dust, and SN ejecta dust.

It is thought that the progenitor of SN 1987A ejected  $8 M_{\odot}$  of gas during its red supergiant (RS) phase before the SN explosion (3). The dust ejected this way is expected to be composed mostly of silicates because RSs produce oxygen-rich dust. Adopting a gas-to-dust mass ratio of 300 (31) implies a silicate dust mass from the RS of  $0.03 M_{\odot}$ . The silicate dust mass required to fit the observed emission is much higher ( $>2 M_{\odot}$ ) (Table 2), equivalent to  $>600 M_{\odot}$  of gas. A



**Fig. 1.** (A to H) The Herschel images of SN 1987A, together with the Spitzer infrared (20) and the Hubble optical (56, 57) images. North is top, and east is left. The two horizontal white lines indicate the position of SN 1987A measured from radio observations. (I) Background-subtracted 350- $\mu\text{m}$  image, where the background is estimated from the 250- $\mu\text{m}$  residual image after the subtraction of the PSF at the position of SN 1987A. The PSFs show the resolution of the Herschel instruments. (B) shows an enlarged HST optical image, indicating the morphology of the SN remnant. [Source: (A) Hubble Heritage Team (Association of Universities for Research in Astronomy (AURA)/Space Telescope Science Institute (STScI)/NASA); (B) NASA, ESA, P. Challis, and R. Kirshner (Harvard-Smithsonian Center for Astrophysics)]

**Table 1.** The measured PACS and SPIRE flux densities of SN 1987A. We added in quadrature the uncertainties from the source extraction and the absolute errors in the flux calibration, which are estimated to be 10% for the PACS bands and 7% for the SPIRE bands (47, 48). \*, 3-SD upper limit.

Band	Flux (mJy)
PACS 100 $\mu\text{m}$	$70.5 \pm 8.5$
PACS 160 $\mu\text{m}$	$125.3 \pm 16.1$
SPIRE 250 $\mu\text{m}$	$131.7 \pm 12.1$
SPIRE 350 $\mu\text{m}$	$49.3 \pm 6.5$
SPIRE 500 $\mu\text{m}$	$<57.3^*$



similar evaluation rules out the possibility that dust could have been formed during another evolutionary phase of the progenitor, for example, a luminous blue variable. It is therefore unlikely that the observed far-infrared/submm emission could have originated from dust formed by the SN progenitor.

As it expands, the SN sweeps up dust from the ambient ISM. The SN shock speed is up to  $6000 \text{ km s}^{-1}$  (32), and the distance already traveled by the shocks is up to 30,000 astronomical units (0.6 arcsec). We assumed a gas density of 1 hydrogen atom per  $\text{cm}^3$  in the LMC ISM and adopted a gas-to-dust mass ratio of 300. Within this volume, only a small mass of ISM dust ( $3 \times 10^{-6} M_{\odot}$ ) will have been swept up. This estimate

is consistent with Herschel measurements of the ISM dust emission of  $<10^{-4} M_{\odot}$ , based on the sky background level in the 160- $\mu\text{m}$  band ( $70.4 \pm 9.6 \text{ MJy sr}^{-1}$ ) measured for a 1.3 by 0.4 arcsec<sup>2</sup> region centered at the position of SN, after removal of a point-spread function (PSF). Our measured dust emission requires a much higher dust mass, hence favoring emission from SN dust rather than from swept-up ISM dust.

The original SN energy pulse could potentially have been absorbed and re-emitted by interstellar dust grains to produce infrared echoes. To account for the Herschel fluxes, an infrared echo would have to originate from a dust cloud located at about 11 light years directly behind the SN, which is improbable. Moreover, this


dust would have been heated by the optical and UV light, which had a luminosity of  $L_{\text{UV}} = 10^{41} \text{ erg sec}^{-1}$  (17), implying a flux incident on the dust  $F = L_{\text{UV}}/(4\pi R^2)$ , where  $R = 11$  light years. Taking this energy input as the dust heating rate, equilibrium with the dust cooling rate yields the dust temperatures ranging from 40 K for silicate and 70 to 110 K for graphite, for a range of grain sizes from 0.01 to 0.1  $\mu\text{m}$  (33). These temperatures are substantially higher than those measured by Herschel (Table 2), ruling out an echo as an explanation for the Herschel SED.

Thus, we conclude that the emission observed by Herschel is associated with cold dust in the SN ejecta.

The dust that is being emitted at Herschel wavelengths could be heated by one or more of the following mechanisms: (i) by the x-ray emission arising from the interaction of the blast wave with the equatorial ring, which has a luminosity of about  $500 L_{\odot}$  (20, 28); (ii) by the energy released from the long-lived radioactive  $^{44}\text{Ti}$  isotope present in the ejecta; our measured far-infrared/submm luminosity of  $220 L_{\odot}$  is about half of the predicted  $^{44}\text{Ti}$  luminosity at this epoch ( $\sim 400 L_{\odot}$ ) (34); or (iii) by the ambient diffuse interstellar radiation field, given that the temperature of the far-infrared/submm emitting dust is very similar to that of ISM dust.

**The origin of the cold dust in the ejecta.** The masses predicted for the major refractory elements (Table 3) in the ejecta of SN 1987A (3, 35) restrict the maximum mass of dust that could have formed in the ejecta if all these elements precipitated from the gas phase (Table 4). The total available dust mass is about  $0.71 M_{\odot}$  in either nucleosynthesis model. The maximum mass of silicate dust of 0.51 or  $0.40 M_{\odot}$ , depending on the models, is smaller than the  $2.4 M_{\odot}$  required to fit the observed far-infrared/submm flux with pure silicate emission. The same holds for carbon and iron dust. It is clear that any attempt to fit the observed spectrum with a single dust species requires a larger mass of that particular dust species than is available in the ejecta.

The far-infrared/submm fluxes could, however, be the sum of the emission from several different dust species in the ejecta. Figure 2, C and



**Fig. 2.** (A) The infrared spectral energy distribution of SN 1987A. Herschel detected SN 1987A from 100 to 350  $\mu\text{m}$ . An upper limit is given at 500  $\mu\text{m}$ . The other observational data were collected from the literature: 5.8- to 24- $\mu\text{m}$  photometric data and 5- to 30- $\mu\text{m}$  spectra (20), which measures the warm dust emission; and the radio continuum (26, 29), which traces the synchrotron emission. (B) Model fits to the far-infrared/submm dust emission using different single dust species. Parameters are given in Table 2. (C and D) The sum of the contributions from each dust species to the far-infrared/submm spectrum of SN 1987A. The flux at each wavelength is that derived from fitting the observed spectrum with each dust species alone (Table 2), scaled by the fraction of its mass available in the ejecta from abundance considerations (Table 4). The model 1 and model 2 abundances in Table 4 correspond to (C) and (D), respectively. The hatched areas represent the range of possible fluxes from each species, with the lower and upper boundaries corresponding, respectively, to the scaling of the minimum and maximum and dust masses (uncertainties in Table 2) required to produce the observed spectrum.

**Table 2.** Dust temperatures ( $T_d$ ) and corresponding dust masses ( $M_d$ ) derived by fitting the whole of the far-infrared/submm emission with a single dust species (30). Quantities are insensitive to grain radius, except metallic iron. The absorption efficiency of metallic iron depends on grain radius ( $a$ ), and the table represents the results for grain radii of 0.1 and 0.5  $\mu\text{m}$ .

Dust species	$M_d (M_{\odot})$	$T_d (K)$
Amorphous carbon	$0.35 \pm 0.06$	$21.2 \pm 0.7$
Silicate	$2.4 \pm 0.4$	$17.7 \pm 0.5$
Iron ( $a = 0.1 \mu\text{m}$ )	$3.4 \pm 0.6$	$19.2 \pm 0.7$
Iron ( $a = 0.5 \mu\text{m}$ )	$0.34 \pm 0.06$	$25.7 \pm 0.9$

**Table 3.** Model prediction of the mass of representative refractory elements synthesized in the SN 1987A ejecta. Isotopes that contribute a large fraction of the total atomic mass were accounted for. Model 1, Theoretical calculations of elemental abundance from (3). Model 2, Calculations from an updated version of (35).

Elements	Mass ( $M_{\odot}$ )	
	Model 1	Model 2
C	0.114	0.26
O	1.48	0.24
Mg	0.183	0.04
Si	0.101	0.14
Fe	0.083	0.075

**Table 4.** The dust mass assumes 100% dust condensation of the available elemental mass ( $m_d$ ). The range of dust masses reflects the difference in compositions in Table 3. All silicates are assumed to be in the form of MgO and SiO<sub>2</sub> dust. The mass of carbon dust assumes that no substantial fraction of carbon is locked up in CO molecules.

Dust species	$m_d (M_\odot)$	
	Model 1	Model 2
Amorphous carbon	0.11	0.26
Silicate	0.52	0.37
Iron	0.08	0.08
Total	0.71	0.71

D, shows the predicted far-infrared SEDs arising from dust in the ejecta of SN 1987A using ejecta abundances from models 1 and 2, respectively (Table 4). Figure 2, C and D, suggests that a substantial fraction of the major refractory elements are depleted onto dust and that the total dust mass is about 0.4 to 0.7  $M_\odot$ .

Support for the presence of large amounts of dust in the ejecta may come from Hubble Space Telescope (HST) observations of the ejecta Ha profile in 2004 and 2010 (36), which show the redward part of the profile to be largely missing, which could be interpreted as implying that emission from material at the far side of the ejecta suffers more extinction by intervening dust than material at the near side.

The derived dust mass is about  $10^3$  times as large as and about 20 times as cold as that measured at mid-infrared wavelengths around 600 days after the explosion was observed, soon after it first condensed out of the ejecta (15, 16). Far-infrared and submm observatories with Herschel's sensitivity did not exist at that time. It is possible that this submm dust mass existed at day 600. However, it is also possible that this dust mass grew by accretion over the past 20 years or that initially optical/infrared-thick dust-bearing clumps may have become optically thin at far-infrared and submm wavelengths. Overall, the observations provide evidence for the efficient formation and growth of massive quantities of dust in SN ejecta.

Our result could affect the understanding of the evolution of dust in galaxies. In the LMC, the rate of type II supernovae is estimated to be one per 300 years (37). If all type II supernovae form typically 0.4  $M_\odot$  of dust, the overall SN dust injection rate into the ISM would be more than  $10^{-3} M_\odot \text{ year}^{-1}$ . This is about 10 times as high as the overall dust inputs measured from asymptotic giant branch stars in the LMC (38, 39). With such a high efficiency and with a dust lifetime of  $2 \times 10^8$  to  $4 \times 10^8$  years (40), it would be possible to explain a substantial fraction of the dust in the LMC (about  $1 \times 10^6 M_\odot$ ) (41, 31) by a stellar origin.

The large amount of dust inferred to be present in the ejecta of SN 1987A is consistent with that required to explain the dust masses in high-redshift galaxies (42, 43), that is, 0.1 to 1.0  $M_\odot$  of dust per SN, if the dust is not substantially destroyed during its injection into the ISM and subsequent encounters with interstellar shocks.

Supernovae may therefore be important contributors of dust detected in such galaxies.

#### References and Notes

- K. Hirata *et al.*, *Phys. Rev. Lett.* **58**, 1490 (1987).
- R. M. Bionta *et al.*, *Phys. Rev. Lett.* **58**, 1494 (1987).
- F.-K. Thielemann, M.-A. Hashimoto, K. Nomoto, *Astrophys. J.* **349**, 222 (1990).
- R. Sunyaev *et al.*, *Nature* **330**, 227 (1987).
- T. Dotani *et al.*, *Nature* **330**, 230 (1987).
- D. Arnett, B. Fryxell, E. Mueller, *Astrophys. J.* **341**, L63 (1989).
- S. H. Moseley *et al.*, *Astrophys. J.* **347**, 1119 (1989).
- W. D. Arnett, J. N. Bahcall, R. P. Kirshner, S. E. Woosley, *Annu. Rev. Astron. Astrophys.* **27**, 629 (1989).
- R. A. Chevalier, C. Fransson, *Nature* **328**, 44 (1987).
- C. J. Burrows *et al.*, *Astrophys. J.* **452**, 680 (1995).
- P. Grönningsson *et al.*, *Astron. Astrophys.* **492**, 481 (2008).
- S. Maitila *et al.*, *Astrophys. J.* **717**, 1140 (2010).
- S. H. Moseley *et al.*, *Nature* **340**, 697 (1989).
- P. A. Whitelock *et al.*, *Mon. Not. R. Astron. Soc.* **240**, 7 (1989).
- D. H. Wooden *et al.*, *Astrophys. J. Suppl. Ser.* **88**, 477 (1993).
- L. B. Lucy, I. J. Danziger, C. Gouffes, P. Bouchet, in *Structure and Dynamics of the Interstellar Medium, Proceedings of IAU Colloq. 120, G. Tenorio-Tagle, M. Moles, J. Melnick, Eds., Also Lecture Notes in Physics, vol 350* (1989), pp. 164–179.
- R. McCray, *Annu. Rev. Astron. Astrophys.* **31**, 175 (1993).
- D. N. Burrows *et al.*, *Astrophys. J.* **543**, L149 (2000).
- P. Bouchet *et al.*, *Astrophys. J.* **650**, 212 (2006).
- E. Dwek *et al.*, *Astrophys. J.* **722**, 425 (2010).
- G. L. Pilbratt *et al.*, *Astron. Astrophys.* **518**, L1 (2010).
- A. Poglitsch *et al.*, *Astron. Astrophys.* **518**, L2 (2010).
- M. J. Griffin *et al.*, *Astron. Astrophys.* **518**, L3 (2010).
- M. Meixner *et al.*, *Astron. Astrophys.* **518**, L71 (2010).
- We combined the two observations because there were no substantial changes in the fluxes seen in the two epochs. The basic steps for the data processing of the PACS and SPIRE data are described by (24). Despite the subtraction of the baseline, residual striping remained along the scan direction of the PACS maps. To mitigate the stripes, we subtracted from the PACS images an image of the striping, produced by unambiguously identifying the power spectrum associated with the stripes in Fourier space. The HERITAGE astrometric reference frame is tied to the Two Micron All Sky Survey (2MASS) (44), which has an accuracy of 0.3" (1 SD), through the Spitzer mid-infrared LMC data (45).
- T. M. Potter *et al.*, *Astrophys. J.* **705**, 261 (2009).
- We used the Starfinder software (46), which employs PSF fitting on complex backgrounds. The SPIRE PSFs are empirical models derived from scan-map data for Neptune (47). For PACS, we used empirical model PSFs, based on maps of Vesta (48).
- R. Sturm, F. Haberl, B. Aschenbach, G. Hasinger, *Astron. Astrophys.* **515**, A5 (2010).
- G. Zandaro *et al.*, *Astrophys. J.* **710**, 1515 (2010).
- We generated 1000 Monte Carlo realization simulations of the spectra. The flux errors are accounted as the normal distribution to account for the maximum and

minimum dust masses. We used three different dust candidates: astronomical silicates, amorphous carbon, and pure iron, whose refractive indices we took from (49–51) and dust densities from (52–54). We calculated the dust mass absorption coefficients ( $\kappa \text{ cm}^2 \text{ g}^{-1}$ ) using Mie theory. In the far-infrared/submm wavelength regime,  $\kappa$  is independent of grain radius up to  $\sim 5 \mu\text{m}$ , except for metallic iron, for which the absorption efficiency depends on the relative sizes of the skin depth and grain radius. The entries for metallic iron were calculated for a nominal radius of 0.1  $\mu\text{m}$  and a radius of 0.5  $\mu\text{m}$ , for which the value of  $\kappa$  is a maximum in the Herschel wavelength range. The iron mass for this radius thus represents a minimum required mass to fit the observations.

- We assumed that the silicon, iron, and magnesium elemental abundances before the explosion were the same as those of the ISM and that these abundances limited the dust mass. We adopted an ISM gas-to-dust mass ratio of 300 because it is higher in the LMC than in the Milky Way (55), where it is typically 100.
- J. L. Racusin *et al.*, *Astrophys. J.* **703**, 1752 (2009).
- These dust temperatures have some dependence on the grain size and the grain compositions.
- A. Jerkstrand, C. Fransson, C. Kozma, *Astron. Astrophys.* **530**, A45 (2011).
- S. E. Woosley, P. A. Pinto, T. A. Weaver, *Proc. Astron. Soc. Aust.* **7**, 355 (1988).
- J. Larsson *et al.*, *Nature* **474**, 484 (2011).
- D. Maoz, C. Badenes, *Mon. Not. R. Astron. Soc.* **407**, 1314 (2010).
- M. Matsuura *et al.*, *Mon. Not. R. Astron. Soc.* **396**, 918 (2009).
- S. Srinivasan *et al.*, *Astron. J.* **137**, 4810 (2009).
- A. P. Jones, A. G. G. M. Tielens, D. J. Hollenbach, *Astrophys. J.* **469**, 740 (1996).
- J.-P. Bernard *et al.*, *Astron. J.* **136**, 919 (2008).
- H. L. Morgan, M. G. Edmunds, *Mon. Not. R. Astron. Soc.* **343**, 427 (2003).
- E. Dwek, I. Cherchneff, *Astrophys. J.* **727**, 63 (2011).
- M. F. Skrutskie *et al.*, *Astron. J.* **131**, 1163 (2006).
- M. Meixner *et al.*, *Astron. J.* **132**, 2268 (2006).
- E. Diolaiti *et al.*, *Astron. Astrophys.* **147** (suppl.), 335 (2000).
- B. M. Swinyard *et al.*, *Astron. Astrophys.* **518**, L4 (2010).
- PACS Observer's Manual; [http://herschel.esac.esa.int/Docs/PACS/html/pacs\\_om.html](http://herschel.esac.esa.int/Docs/PACS/html/pacs_om.html).
- J. C. Weingartner, B. T. Draine, *Astrophys. J.* **548**, 296 (2001).
- V. G. Zubko, V. Mennella, L. Colangeli, E. Bussoletti, *Mon. Not. R. Astron. Soc.* **282**, 1321 (1996).
- D. Semenov, T. Henning, C. Helling, M. Ilgner, E. Sedlmayr, *Astron. Astrophys.* **410**, 611 (2003).
- A. Laor, B. T. Draine, *Astrophys. J.* **402**, 441 (1993).
- F. Rouleau, P. G. Martin, *Astrophys. J.* **377**, 526 (1991).
- T. Nozawa, T. Kozasa, A. Habe, *Astrophys. J.* **648**, 435 (2006).
- K. D. Gordon, G. C. Clayton, K. A. Misselt, A. U. Landolt, M. J. Wolff, *Astrophys. J.* **594**, 279 (2003).
- N. Panagia, M. Romaniello, S. Scuderi, R. P. Kirshner, *Astrophys. J.* **539**, 197 (2000).
- K. France *et al.*, *Science* **329**, 1624 (2010).

**Acknowledgments:** Herschel is a European Space Agency (ESA) space observatory with science instruments provided by European-led principal investigator consortia and with important participation from NASA. We acknowledge financial support from the NASA Herschel Science Center, JPL contracts 1381522 and 1381650. We acknowledge the contributions and support of the ESA, the PACS and SPIRE teams, the Herschel Science Centre, the NASA Herschel Science Center, and the PACS and SPIRE instrument control centers, without which none of this work would have been possible. We also thank the HERITAGE team for various inputs on the data reduction. M.M. appreciates many discussions and inputs from H. Hirashita, I. Sakon, R. Wesson, K. Nomoto, and B. M. Swinyard. M.M. acknowledges the award of a University College London Institute of Origins Fellowship, and M.L. acknowledges a European Southern Observatory-Keele studentship.

23 March 2011; accepted 21 June 2011  
Published online 7 July 2011;  
10.1126/science.1205983