

# Star Formation<sup>1</sup>

Kohji Tomisaka  
National Astronomical Observatory Japan

version 0.5: August 5, 2001  
version 1.0: August 30, 2001  
version 2.0: November 8, 2003

<sup>1</sup> [http://yso.mtk.nao.ac.jp/~tomisaka/Lecture\\_Notes/StarFormation/2.pdf](http://yso.mtk.nao.ac.jp/~tomisaka/Lecture_Notes/StarFormation/2.pdf)

# Contents

<b>1</b>	<b>Introduction</b>	<b>3</b>
1.1	Interstellar Matter . . . . .	3
1.2	Case Study — Taurus Molecular Clouds . . . . .	4
1.3	T-Tauri Stars . . . . .	6
1.4	Spectral Energy Distribution (SED) . . . . .	6
1.5	Protostars . . . . .	10
1.5.1	B335 . . . . .	10
1.5.2	L1551 IRS 5 . . . . .	14
1.6	L 1544: Pre-protostellar Cores . . . . .	17
1.7	Magnetic Fields . . . . .	17
1.7.1	Cores with Protostars . . . . .	19
1.8	Density Distribution . . . . .	23
1.9	Mass Spectrum . . . . .	26
<b>2</b>	<b>Physical Background</b>	<b>29</b>
2.1	Basic Equations of Hydrodynamics . . . . .	29
2.2	The Poisson Equation of the Self-Gravity . . . . .	29
2.3	Free-fall Time . . . . .	30
2.3.1	Accretion Rate . . . . .	32
2.4	Gravitational Instability . . . . .	33
2.4.1	Sound Wave . . . . .	34
2.5	Jeans Instability . . . . .	35
2.6	Gravitational Instability of Thin Disk . . . . .	36
2.7	Super- and Subsonic Flow . . . . .	38
2.7.1	Flow in the Laval Nozzle . . . . .	38
2.7.2	Steady State Flow under an Influence of External Fields . . . . .	39
2.7.3	Stellar Wind — Parker Wind Theory . . . . .	40
<b>3</b>	<b>Galactic Scale Star Formation</b>	<b>45</b>
3.1	Schmidt Law . . . . .	45
3.1.1	Global Star Formation . . . . .	45
3.1.2	Local Star Formation Rate . . . . .	46
3.2	Gravitational Instability of Rotating Thin Disk . . . . .	47
3.2.1	Tightly Wound Spirals . . . . .	49
3.2.2	Toomre’s $Q$ Value . . . . .	50
3.3	Spiral Structure . . . . .	51
3.4	Density Wave Theory . . . . .	52

3.5	Galactic Shock . . . . .	55
<b>4</b>	<b>Local Star Formation Process</b>	<b>61</b>
4.1	Hydrostatic Balance . . . . .	61
4.1.1	Bonnor-Ebert Mass . . . . .	62
4.1.2	Equilibria of Cylindrical Cloud . . . . .	63
4.2	Virial Analysis . . . . .	64
4.2.1	Magnatohydrostatic Clouds . . . . .	65
4.3	Subcritical Cloud vs Supercritical Cloud . . . . .	67
4.4	Ambipolar Diffusion . . . . .	67
4.4.1	Ionization Rate . . . . .	67
4.4.2	Ambipolar Diffusion . . . . .	68
4.5	Dynamical Collapse . . . . .	71
4.5.1	Protostellar Evolution of Supercritical Clouds . . . . .	76
4.6	Accretion Rate . . . . .	77
4.7	Outflow . . . . .	79
4.7.1	Magneto-driven Model . . . . .	79
4.7.2	Entrainment Model . . . . .	83
4.8	Evolution to Star . . . . .	83
4.9	Example of Numerical Simulation . . . . .	86
<b>A</b>	<b>Basic Equation of Fluid Dynamics</b>	<b>97</b>
A.1	What is fluid? . . . . .	97
A.2	Equation of Motion . . . . .	97
A.3	Lagrangian and Euler Equation . . . . .	97
A.4	Continuity Equation . . . . .	98
A.5	Energy Equation . . . . .	99
A.5.1	Polytropic Relation . . . . .	99
A.5.2	Energy Equation from the First Law of Thermodynamics . . . . .	99
A.6	Shock Wave . . . . .	100
A.6.1	Rankine-Hugoniot Relation . . . . .	100
<b>B</b>	<b>Basic Equations of Magnetohydrodynamics</b>	<b>101</b>
B.1	Magnetohydrodynamics . . . . .	101
B.1.1	Flux Freezing . . . . .	101
B.1.2	Basic Equations of Idel MHD . . . . .	102
B.1.3	Axisymmetric Case . . . . .	102
<b>C</b>	<b>Hydrostatic Equilibrium</b>	<b>105</b>
C.1	Polytrope . . . . .	105
C.2	Magnetohydrostatic Configuration . . . . .	107
<b>D</b>	<b>Basic Equations for Radiative Hydrodynamics</b>	<b>109</b>
D.1	Radiative Hydrodynamics . . . . .	109



# Chapter 1

## Introduction

### 1.1 Interstellar Matter




Figure 1.1: Multiphases of the interstellar medium. The temperature and number density of gaseous objects of the interstellar medium in our Galaxy are summarized. Originally made by Myers (1978), reconstructed by Saigo (2000).

Figure 1.1 shows the temperature and number density of gaseous objects in our Galaxy. Cold interstellar medium forms molecular clouds ( $T \sim 10\text{K}$ ) and diffuse clouds ( $T \sim 100\text{K}$ ). Warm interstellar medium  $10^3\text{K} \lesssim T \lesssim 10^4\text{K}$  are thought to be pervasive (wide-spread). HII regions are ionized by the Ly continuum photons from the early-type stars. There are coronal (hot but tenuous) gases with  $T \sim 10^6\text{K}$  in the Galaxy, which are heated by the shock fronts of supernova remnants. Pressures of these gases are in the range of  $10^2\text{K cm}^{-3} \lesssim p \lesssim 10^4\text{K cm}^{-3}$ , except for the HII regions. This may suggest that the gases are in a pressure equilibrium.

In this figure, a theoretical path from the molecular cloud core to the star is also shown. We will see the evolution more closely in Chapter 4.

Globally, the molecular form of Hydrogen is much abundant inside the Solar circle, while the atomic hydrogen is more abundant than molecular  $H_2$  in the outer galaxy. In Figure 1.2 (left), the radial distributions of molecular and atomic gas are shown. The right panel shows the distributions for four typical external galaxies (M51, M101, NGC6946, and IC342). This indicates these distributions are similar with each other. HI is distributed uniformly, while  $H_2$  density increases greatly reaching the galaxy center. In other words, only in the region where the total (HI+ $H_2$ ) density exceeds some critical value,  $H_2$  molecules are distributed.




Figure 1.2: Radial distribution of  $H_2$  (solid line) and HI (dashed line) gas density. (Left:) our galaxy. Converting from CO antenna temperature to  $H_2$  column density,  $n(\text{CO})/n(\text{H}_2) = 6 \times 10^{-5}$  is assumed. Taken from Gordon & Burton (1976). (Right:) Radial distribution of  $H_2$  and HI gas for external galaxies. The conversion factor is assumed constant  $X(\text{H}_2/\text{CO}) = 3 \times 10^{20} \text{H}_2/\text{K km s}^{-1}$ . Taken from Honma et al (1995).

## 1.2 Case Study — Taurus Molecular Clouds

Figure 1.3 (left) shows the  $^{13}\text{CO}$  total column density map of the Taurus molecular cloud (Mizuno et al 1995) whose distance is 140 pc far from the Sun. Since  $^{13}\text{CO}$  contains  $^{13}\text{C}$ , a rare isotope of C, the abundance of  $^{13}\text{CO}$  is much smaller than that of  $^{12}\text{CO}$ . Owing to the low abundance, the emission lines of  $^{13}\text{CO}$  are relatively optically thinner than that of  $^{12}\text{CO}$ . Using  $^{13}\text{CO}$  line, we can see deep inside of the molecular cloud. The distributions of T-Tauri stars and  $^{13}\text{CO}$  column density coincide with each other. Since T-Tauri stars are young pre-main-sequence stars with  $M \sim 1M_{\odot}$ , which are in the Kelvin contraction stage and do not reach the main sequence Hydrogen burning stage, it is shown that stars are newly formed in molecular clouds.




Figure 1.3: (*Left*)  $^{13}\text{CO}$  total column density map of the Taurus molecular cloud (Mizuno et al 1995). Taken from their home page with a url of <http://www.a.phys.nagoya-u.ac.jp/nanten/taurus.html> (in Japanese). T Tauri stars, which are thought to be pre-main-sequence stars in the Kelvin contraction stage, are indicated by bright spots. (*Right*)  $\text{C}^{18}\text{O}$  map of Heiles cloud 2 region in the Taurus molecular cloud (Onishi et al. 1996). This shows clearly that the cloud is composed of a number of high-density regions.

Since  $^{18}\text{O}$  is much more rare isotope ( $^{18}\text{O}/^{16}\text{O} \ll ^{13}\text{C}/^{12}\text{C}$ ), the distribution of much higher-density gases is explored using  $\text{C}^{18}\text{O}$  lines. Figure 1.3 (right) shows  $\text{C}^{18}\text{O}$  map of Heiles cloud 2 in the Taurus molecular cloud by Onishi et al (1996). This shows us that there are many molecular cloud cores which have much higher density than the average. Many of these molecular cloud cores are associated with IRAS sources and T-Tauri stars. It is shown that star formation occurs in the molecular cloud cores in the molecular cloud. They found 40 such cores in the Taurus molecular cloud. Typical size of the core is  $\sim 0.1$  pc and the average density of the core is as large as  $\sim 10^4 \text{cm}^{-3}$ . The mass of the  $\text{C}^{18}\text{O}$  cores is estimated as  $\sim 1 - 80 M_{\odot}$ .

$\text{H}^{13}\text{CO}^+$  ions are excited only after the density is much higher than the density at which CO molecules are excited.  $\text{H}^{13}\text{CO}^+$  ions are used to explore the region with higher density than that observed by  $\text{C}^{18}\text{O}$ . Figure 1.4 shows the map of cores observed by  $\text{H}^{13}\text{CO}^+$  ions. The cores shown in the lower panels are accompanied with infrared sources. The energy source of the stellar IR radiation is thought to be maintained by the accretion energy. That is, since the gravitational potential energy at the surface of a protostar with a radius  $r_*$  and a mass  $M_*$  is equal to  $\Phi \simeq -GM_*/r_*$ , the kinetic energy of the gas accreting on the stellar surface is approximately equal to  $\sim GM_*/r_*$ . The energy inflow rate owing to the accretion is  $(\sim GM_*/r_*) \times \dot{M} \sim (GM_*/r_*) \times A(c_s^3/G)$ , where  $\dot{M} = A(c_s^3/G)$  is the mass accretion rate. In the upper panels, the cores without IR sources are shown. This core does not show accretion but collapse. That is, before a protostar is formed, the core itself contract owing to the gravity.

In Figure 1.4,  $\text{H}^{13}\text{CO}^+$  total column density maps of the  $\text{C}^{18}\text{O}$  cores are shown. Cores in the lower panels have associated IRAS sources, while the cores in the upper panels have no IRAS sources. Since the IRAS sources are thought to be protostars or objects in later stage, the core seems to evolve from that without an IRAS sources to that with an IRAS source. From this, the core with an IRAS source is called **protostellar core**, which means that the cores contain protostars. On the other hand, the core without IRAS source is called pre-protostellar core or, in short, **pre-stellar core**.

Figure 1.4 shows that the prestellar cores are less dense and more extended than the protostellar core. This seems to suggest the density distribution around the density peak changes between before

FIG. 1 Contour maps of  $\text{H}^{13}\text{CO}^+$  ( $J=1-0$ ) total intensity of typical dense cores obtained with the Nobeyama 45-m telescope. The number on the map corresponds to the core number listed in Table 1. Upper panels are maps of dense cores without IRAS emission, except for core (7), which is the lower left core in panel (2), (7). Lower panels are dense cores with IRAS emission. The beam size is 20 arcsec at 3 mm wavelength, corresponding to 0.014 pc at the distance of Taurus. Typical r.m.s. noise of the spectra is  $\sim 0.15$  K at the velocity resolution of  $0.13 \text{ km s}^{-1}$ . Contours are from  $0.14 \text{ K km s}^{-1}$  with a  $0.07 \text{ K km s}^{-1}$  step. The positions of IRAS sources are indicated as crosses. The reference positions (given as RA(1950), dec.(1950)) of offsets are as follows: core (1), 4 h 15 min 5.4 s,  $27^\circ 27' 43''$ ; core (2) and core (7), 4 h 16 min 35.5 s,  $27^\circ 27' 43''$ ; core (3), 4 h 25 min 34.5 s,  $26^\circ 45' 4''$ ; core (8), 4 h 16 min 53.8 s,  $27^\circ 2' 52''$ ; core (11), 4 h 28 min 40.2 s,  $18^\circ 1' 42''$ ; core (15), 4 h 36 min 49.3 s,  $25^\circ 57' 16''$ . The half-power beam width (HPBW) is indicated as a circle in panel (15).




Figure 1.4: Pre-protostellar vs protostellar cores ( $\text{H}^{13}\text{CO}^+$  map). Upper panel shows the  $\text{C}^{18}\text{O}$  cores without associated IRAS sources. Lower panel shows the cores with IRAS sources. Taken from Fig.1 of Mizuno et al. (1994)

and after the protostar formation.

### 1.3 T-Tauri Stars

T-Tauri stars are observationally late-type stars with strong emission-lines and irregular light variations associated with dark or bright nebulosities. T-Tauri stars are thought to be low-mass pre-main-sequence stars, which are younger than the main-sequence stars. Since these stars are connecting between protostars and main-sequence stars, they attract attention today. More massive counterparts are Herbig Ae-Be stars. They are doing the Kelvin contraction in which the own gravitational energy released as it contracts gradually is the energy source of the luminosity. Many emission lines are found in the spectra of T-Tauri stars. WTTS (Weak Emission T-Tauri Stars) and CTTS (Classical T-Tauri Stars) are classified by their equivalent widths of emission lines. That is, the objects with an EW of  $\text{H}\alpha$  emission  $< 1 \text{ nm} = 10 \text{ \AA}$  is usually termed a WTTS. Figure 1.5 is the HR diagram ( $T_{\text{eff}} - L_{\text{bol}}$ ) of T-Tauri stars in Taurus-Auriga region (Kenyon & Hartman 1995). WTTSs distribute near the main-sequence and CTTSs are found even far from the main-sequence. A number of theoretical evolutionary tracks for pre-main-sequence stars with  $M \sim 0.1M_{\odot} - 2.5M_{\odot}$  are shown in a solid line, while the isochrones for ages of  $10^5 \text{ yr}$ ,  $10^6 \text{ yr}$ , and  $10^7 \text{ yr}$  are plotted in a dashed line. Vertical evolutionary paths are the Hayashi convective track. Since  $\text{D}=\text{}^2\text{H}$  has a much lower critical temperature (and density) for a fusion nuclear reaction to make He than  $\text{}^1\text{H}$ , Deuterium begins to burn before reaching the zero-age-main sequence. This occurs near the isochrone for the age of  $10^5 \text{ yr}$  and some activities related to the ignition of Deuterium seem to make the central star visible (Stahler 1983).

### 1.4 Spectral Energy Distribution (SED)

A tool to know the process of star formation is provided by the spectral energy distribution (SED) mainly in the near- and mid-infrared light. T-Tauri stars and protostars have typical respective SEDs. IR SEDs of T-Tauri stars were classified into three as Class I, Class II, and Class III, from a stand-






Fig. 1.2. HR diagram positions of young stars lying within the Taurus–Auriga molecular cloud complex (Figure 1.1). For comparison, theoretical evolutionary tracks for pre-main-sequence stars of masses 2.5, 2.0, 1.5, 1.0, 0.5, 0.3, and 0.1  $M_{\odot}$  are shown. The dashed lines are isochrones for ages of  $10^5$ ,  $10^6$ , and  $10^7$  yr (0.1, 1, and 10 Myr), with the hydrogen-fusion ‘zero-age main sequence’ or ZAMS shown as the lowest line running from upper left to lower right. The open circles refer to weak-emission T Tauri stars (WTTS; see text), while the filled circles denote the positions of the classical T Tauri stars (CTTS). Stellar properties taken from Kenyon & Hartmann (1995); evolutionary tracks are from D’Antona & Mazzitelli (1994).

Figure 1.5: HR diagram of T-Tauri stars. Many emission lines are found in the spectra of T-Tauri stars. WTTS (Weak Emission T-Tauri Stars) and CTTS (Classical T-Tauri Stars) are classified by the equivalent widths of emission lines. That is, the objects with an EW of  $H\alpha$  emission  $< 1\text{nm}$  is usually termed a WTTS. Taken from Fig.1.2 of Hartmann (1998).

point of relative importance of the radiation from a dust disk to the stellar black-body radiation. Today, the classification is extended to the protostars, which is precedence of the T-Tauri stars, and they are called Class 0 objects. (Unfortunately, there is no zero in Roman numerals.) In Figure 1.6, typical SEDs and models of emission regions are shown.

1. Class III is well fitted by a black-body spectrum, which shows the energy mainly comes from a central star. This SED is observed in the weak-line T-Tauri stars. Although T-Tauri stars show emission lines of such as Hydrogen Balmer sequence, the weak-line T-Tauri stars do not show prominent emission lines, which indicates the amount of gas just outside the star (this seems to be supplied by the accretion process) is small. In this stage, a disk has been disappeared or an extremely less-massive disk is still alive.
2. Class II SED is fitted by a single-temperature black-body plus excess IR emission. This shows that there is a dust disk around a pre-main sequence star and it is heated by the radiation from a central star. The width of the spectrum of the disk component is much wider than that expected from a single-temperature black-body radiation. Thus, the disk has a temperature gradient which decreases with increasing the distance from the central star. In this stage, the dust disk is more massive than that of Class III sources. Classical T-Tauri stars have such SEDs.
3. In Class I SED, the mid infrared radiation which seems to come from the dust envelope is predominant over the stellar black-body radiation. Since the stellar black-body radiation seems to escape at least partially from the dust envelope, a relatively large solid angle is expected for a region where the dust envelope does not intervene.
4. Class 0 SED seems to be emitted by isothermal dust with  $\sim 30\text{K}$ . The protostar seems to be completely covered by gas and dust and is obscured with a large optical depth by the dust envelope. No contribution can be reached from the stellar-black body radiation.

The reason why the emission from the disk becomes wide in the spectral range is understood (Fig.1.7) as follows: Temperature of the disk is determined by a balance of heating and cooling. Assuming the disk is geometrically thin but optically thick, the cooling per unit area is given by the equation of the black-body Planck radiation. Therefore, the temperature is determined by the heating predominantly by viscous heating and extra heating by the radiation from the central star. The flux density emitted by the disk is given by

$$\nu F_\nu \sim \int \nu \pi B_\nu [T(r)] 2\pi r dr \sim r (T \text{ or } \nu)^2 \nu B_\nu. \quad (1.1)$$

Assuming the radial distribution of temperature as

$$T = T_0 \left( \frac{R}{R_0} \right)^{-q}, \quad (1.2)$$

( $q = 3/4$  for the standard accretion disk) and taking notice that each temperature in the disk radiates at a characteristic frequency  $\nu \propto T$  (the Wien's law for black-body radiation)

$$\nu F_\nu \sim r^2 \nu B_\nu \propto \nu^4 T^{-2/q} \propto \nu^{4-2/q}, \quad (1.3)$$

where we used the fact that the peak value of  $B_\nu \propto \nu^3$ . Therefore, it is shown that

$$\nu F_\nu \propto \nu^n; \quad n = 4 - \frac{2}{q}. \quad (1.4)$$




Figure 1.6: Spectral Energy Distribution (SED) of young stellar objects (YSOs) and their models. (Left):  $\nu - \nu F_\nu$  plot taken from Lada (1999). (Right):  $\lambda - \lambda F_\lambda$  plot taken from André (1994)




Figure 1.7: Explanation for the spectral index of the emission from a geometrically thin but optically thick disk. Taken from Fig.16 of Lada (1999).

As shown in the previous section, we have no young **stellar** objects found by IR before a protostar is formed. These kind of objects (pre-protostellar core) are often called Class -1. The classification was originally based on the SED and did not exactly mean an evolution sequence. However, today YSOs are considered to evolve as the sequence of the classes: Class -1  $\rightarrow$  Class 0  $\rightarrow$  Class I  $\rightarrow$  Class II  $\rightarrow$  Class III  $\rightarrow$  main-sequence star.

## 1.5 Protostars

### 1.5.1 B335

B335 is a dark cloud (Fig.1.8) with a distance of  $D \simeq 250$ pc. Inside the dark cloud, a Class 0 IR source is found. The object is famous for the discovery of gas infall motion. In Figure 1.9, the




Figure 1.8: Near infrared images of B335, which is a Class 0 source.




FIG. 4a

Figure 1.9: Line profile of CS  $J = 2 - 1$  line radio emission. Model spectra illustrated in a dashed line (Zhou 1995) are overlaid on to the observed spectra in a solid line (Zhou et al 1993).




Figure 1.10: Explanation of blue-red asymmetry when we observe a spherical symmetric inflow motion. An isovelocity curve for the red-shifted gas is plotted in a solid line. That for the blue-shifted gas is plotted in a dashed line. Taken from Fig.14 of Lada (1999).

line profiles of CS  $J = 2 - 1$  line emissions are shown (Zhou et al. 1993). The relative position of the profiles correspond to the position of the beam. (9,9) represents the offset of (9'',9'') from the center. At the center (0,0), the spectrum shows two peaks and the blue-shifted peak is brighter than the red-shifted one. This is believed to be a sign of gas infall motion. The blue-red asymmetry is explained as follow:

1. Considering a spherical symmetric inflow of gas, whose inflow velocity  $v_r$  increases with reaching the center (a decreasing function of  $r$ )
2. Considering a gas element at  $\mathbf{r}$  moving at a speed of  $v_r(\mathbf{r}) < 0$ , the velocity projected on a line-of-sight is equal to

$$v_{\text{line-of-sight}} = v_{\text{systemic}} + v_r \cos \theta, \quad (1.5)$$

where  $v_{\text{systemic}}$  represents the systemic velocity of the cloud (line-of-sight velocity of the cloud center) and  $\theta$  is the angle between the line-of-sight and the position vector of the gas element. The isovector lines, the line which connect the positions whose procession/recession velocities are the same, become like an ellipse shown in Fig.1.10.

3. An isovelocity curve for the red-shifted gas is plotted in a solid line. That for the blue-shifted gas is plotted in a dashed line. If the gas is optically thin, the blue-shifted and red-shifted gases contribute equally to the observed spectrum and the blue- and red-shifted peaks of the emission line should be the same.
4. In the case that the gas has a finite optical depth, for the red-shifted emission line a cold gas in the fore side absorbs effectively the emission coming from the hot interior. On the other hand, for the blue-shifted emission line, the emission made by the hot interior gas escapes from the cloud without absorbed by the cold gas (there is no cold blue-shifted gas).




Figure 1.11:  $\text{C}^{18}\text{O}$  total column density map (left) and  $\text{H}^{13}\text{CO}^+$  channel map (right) of B335 along with the position-velocity maps along the major and minor axes. Taken from Fig.3 of Saito et al (1999).

5. As a result, the blue peak of the emission line becomes more prominent than that of the red-shifted emission. This is the explanation of the blue-red asymmetry.

In Figure 1.9, model spectra calculated with the Sobolev approximation (Zhou 1995) are shown. These show the blue-red asymmetry (the blue line > the red line).

Many bipolar molecular outflows are found in star forming regions. B335 is also an outflow source. In Figure 1.11, distributions of high density gases traced by the  $\text{C}^{18}\text{O}$  and  $\text{H}^{13}\text{CO}^+$  lines are shown as well as the bipolar outflow whose outline is indicated by a shadow (Hirano et al 1988). Comparing left and right panels, it is shown that the distribution of  $\text{C}^{18}\text{O}$  gas is more extended than that of  $\text{H}^{13}\text{CO}^+$  which traces higher-density gas. And the distribution of the  $\text{H}^{13}\text{CO}^+$  is more compact and the projected surface density seems to show the the actual distribution is spherical. And the molecular outflow seems to be ejected in the direction of the minor axis of the high-density gas. It may suggest that (1) a molecular outflow is focused or collimated by the effect of density distribution or that (2) collimation is made by the magnetic fields which run preferentially perpendicularly to the gas disk. This gas disk is observed by these high density tracers.

Combining the  $\text{C}^{18}\text{O}$  and  $\text{H}^{13}\text{CO}^+$  distributions, the surface density distribution along the major axis is obtained by Saito et al (1999). From the lower panel of Figure 1.12, the column density distribution is well fitted in the range from 7000 to 42,000 AU in radius,

$$\Sigma(r) = 6.3 \times 10^{21} \text{cm}^{-2} \left( \frac{r}{10^4 \text{AU}} \right)^{-0.95}, \quad (1.6)$$

where they omitted the data of  $r \lesssim 7000$  since the beam size is not negligible. Similar power-law density distributions are found by the far IR thermal dust emission.




Figure 1.12: Column density distribution  $N_{\text{H}}(r)$  derived from the  $\text{H}^{13}\text{CO}^+$  and  $\text{C}^{18}\text{O}$  data taken by the Nobeyama 45 m telescope. Taken from Fig.9 of Saito et al (1999).

### 1.5.2 L1551 IRS 5




Figure 1.13: (*Left:*)  $^{13}\text{CO}$  column density distribution. The contour lines represent the distribution of  $^{13}\text{CO}$  column density.  $2.2\ \mu\text{m}$  infra-red reflection nebula is shown in grey scale which was observed by Hoddap (1994). (*Right:*) Schematic view of L1551 IRS5 region.

L1551 IRS 5 is one of the most well studied objects. This has an infra-red emission nebosity (Fig.1.13). It is believed that there is a hole perpendicular to the high-density disk and the emission from the central star escapes through the hole and irradiate the nebosity. In this sense this is a reflection nebula. L1551 IRS 5 has an elongated structure of dense gas similar to that observed in B335. The gas is extending in the direction from north-west to south-east [Fig.1.13 (left)]. Since the opposite side of the nebosity is not observed, the opposite side of nebosity seems to be located beyond the high-density disk and be obscured by the disk. This is possible if we see the south surface of the high-density disk as in Figure 1.13 (right).






Figure 1.14: IsovLOCITY contours measured by the  $^{13}\text{CO } J = 1 - 0$  line. It should be noticed that the isovelocity lines run parallelly to the major axis. The north-eastern side shows a red-shift and the south-western side shows a blue-shift.

### Infall Motion

The inflow motion is measured. Figure 1.14 shows the isovelocity contours measured by the  $^{13}\text{CO } J = 1 - 0$  observation (Ohashi et al 1996). It should be noticed that the isovelocity lines run parallelly to the major axis. The north-eastern side shows a red-shift and the south-west side shows a blue-shift. Considering the configuration of the gas disk shown in Fig.1.13 (right), this pattern of isovelocity contours indicates not outflow but inflow. That is, the north-east side is a near side of the disk and the south-west side is a far side. Since a red-shifted motion is observed in the near side and a blue-shifted motion is observed in the far side, it should be concluded that the gas disk of the L1551 IRS5 is now infalling.

### Optical Jet

HST found two optical jets emanating from L1551 IRS5. This has been observed by SUBARU telescope jet emission is dominated by [FeII] lines in the J- and H-bands. The jet extends to the south-western direction and disappears at  $\sim 10'' \simeq 1400\text{AU}$  from the IRS5. The width-to-length ratio is very small  $\lesssim 1/10$  or less, while the bipolar molecular outflow shows a less collimated flow. As for the origin of the two jets, these two jets might be ejected from a single source. However, since there are at least two radio continuum sources in IRS5 within the mutual separation of  $\sim 0.''5$  [see Fig.1.15 (right)], these jets seem to be ejected from the two sources independently.




Figure 1.15: (*Left:*) Infrared image (J- and K-band) of the IR reflection nebula around L1551 IRS5 by SUBARU telescope. Taken from Fig.1 of Itoh et al. (2000). (A jpeg file is available from the following url: [http://SubaruTelescope.org/Science/press\\_release/9908/L1551.jpg](http://SubaruTelescope.org/Science/press_release/9908/L1551.jpg)). (*Right:*) Central 100 AU region map of L1551 IRS5. This is taken by the  $\lambda = 2.7\text{cm}$  radio continuum observation. Deconvolved map (lower-left) shows clearly that IRS5 consists of two sources. Taken from Looney et al. (1997).

Although the lengths of these jets are restricted to  $10''$ , Herbig-Haro jets, which are much larger than the jets in L1551 IRS5, have been found. HH30 has a  $\sim 500$  AU-scale jet whose emission is mainly from the shock-excited emission lines. One of the largest ones is HH111, which is a member of the Orion star forming region and whose distance is as large as  $D \sim 400$ pc, and a jet with a length of  $\sim 4$ pc is observed. Source of HH111 system is thought to consist of at least binary stars or possibly triple stars [Reipurth et al (1999)]. Star A, which coincides with a  $\lambda = 3.6$  cm radio continuum source (VLA 1), shows an elongation in the VLA map whose direction is parallel to the axis of the jet. Therefore, star A is considered to be a source of the jet. Since the VLA map of star A shows another elongated structure perpendicular to the jet axis, star A may be a binary composed by two outflow sources.

## 1.6 L 1544: Pre-protostellar Cores

L1544 is known as a pre-protostellar core (Taffala et al 1998). That shows an infall motion but this contains no IR protostars. In Figure 1.17(*left*), CCS total column density map is shown, which shows an elongated structure. Ohashi et al (1999) have found both rotation and infall motion in the cloud. Position-velocity (PV) diagram along the minor axis shows the infall motion. That along the major axis indicates a rotational motion, which is shown by a velocity gradient. Due to a finite size of the beam, a contraction motion is also shown in the PV diagram along the major axis.

## 1.7 Magnetic Fields

Directions of B-Field are studied by (1) measuring the polarization of light which is suffered from interstellar absorption. In this case the direction of magnetic field is parallel to the polarization vector. The reason is explained in Figure 1.18. In the magnetic fields, the dusts are aligned in a way that their major axes are perpendicular to the magnetic field lines. Such aligned dusts absorb selectively the radiation whose E-vector is parallel to their major axes. As a result, the detected light has a polarization parallel to the magnetic field lines.

However, the polarization measurement in the near IR wavelength limited to the region with low gas density, because background stars suffer severe absorption and becomes hard to be observed if we want to measure the polarization of the high-density region. More direct method is (2) the measurement of the linear polarization of the thermal emission from dusts in the mm wavelengths; in this case the direction of magnetic field is perpendicular to the polarization vector. The mechanism is explained in Figure 1.18b. The aligned dusts, whose major axes are perpendicular to the magnetic field lines, emit the radiation whose E-vector is parallel to the major axes. Since the absorption does not have a severe effect in this mm wavelengths, this gives information the magnetic fields deep inside the clouds.

### Prestellar Core

Figure 1.19 illustrates the polarization maps of three prestellar cores (L1544, L183, and L43) done in the  $850 \mu\text{m}$  band by JCMT-SCUBA. In L1544 and L183 the mean magnetic fields are at an angle of  $30$  deg to the minor axes of the cores. L43 is not a simple object; there is a T Tauri star located in the second core which extends to south-western side of the core (an edge of this core is seen near the western SCUBA frame boundary). And a molecular outflow from the source seems to affect the core. The magnetic field as well as the gas are swept by the molecular outflow. L43 seems an exception.




Figure 1.16: A mosaic image of HH 111 based on HST NICMOS images (bottom) and WFPC2 images (top). Taken from Fig.1 of Reipurth et al (1999).




Figure 1.17: CCS image of prestellar core L1544. (*Left:*) Total intensity map. (*Right:*) PV diagrams along the minor axis (left) and along the major axis (right).

The fact that the mean magnetic fields are parallel to the minor axis of the gas distribution seems the gas contracts preferentially in the direction parallel to the magnetic fields.

### 1.7.1 Cores with Protostars

B-field at the position of protostars and T-Tauri stars are measured for IRAS 16293-2422, L1551 IRS5, NGC1333 IRAS 4A, and HL Tau (Tamura et al. 1995). Although HL Tau is a T-Tauri star, it has a gas disk. Thus this is a Class I source. The others are believed to be in protostellar phase (Class 0 sources). It is known that IRAS 16293-2422, L1551 IRS5, and HL Tau have disks with the radii of 1500-4000 AU from radio observations of molecular lines. Further, near infrared observations have shown that these objects have 300-1000 AU dust disks. Figure 1.20 shows the E-vector of polarized light. If this is the dust thermal radiation, the direction of B-fields is perpendicular to the polarization E-vector. Figure shows the B-fields run almost perpendicular to the elongation of the gas disk. Global directions of B-field outside the gas disk and the direction of CO outflows are also shown in the figure by arrows. It is noteworthy that the directions of local B-fields, global B-fields, and outflows coincide with each other within  $\sim 30$  deg.




Figure 1.18: Explanation how the polarized radiation forms. Taken from Weintraub et al.(2000).




Figure 1.19: Directions of B-Field are shown from the linear polarization observation of  $850 \mu\text{m}$  thermal emission from dusts by JCMT-SCUBA. L 1544 and L183, the magnetic field and the minor axis of the molecular gas distribution coincide with each other within  $\sim 30\text{deg}$ . Taken from Ward-Thompson et al (2000).




FIG. 1.—(a) IRAS 16293–2422: Geometrical relationship among the electric vector of the 1.1 mm continuum polarization ( $19''$  beam), the embedding cloud magnetic field from optical or near-infrared polarimetry, the envelope/disk structure of gas or dust, and the directions of blue/red lobes of high-velocity CO outflows (this work; Vrba et al. 1976; Tamura 1988; Mundt et al. 1990; Mizuno et al. 1990). Note that the direction of the envelope/disk magnetic field inferred from the 1 mm polarization is perpendicular to the electric vector. (b) Same as (a) except for L1551 IRS 5 (this work; Vrba et al. 1976; Sargent et al. 1988; Snell et al. 1980). (c) Same as (a) except for NGC 1333 IRAS 4A (this work; Sandell et al. 1991; Sandell & Aspin 1933). (d) Same as (a) except for HL Tau. The optical jet is also shown (this work; Hayashi et al. 1993; Vrba et al. 1976; Mundt, Ray, & Bührke 1988).

Figure 1.20: Polarization of the radio continuum  $\lambda = 1\text{mm}$ ,  $\lambda = 0.8\text{mm}$ . IRAS 16293-2422 (*upper-left*), L1551 IRS5 (*upper-right*), NGC1333 IRAS 4A (*lower-left*), and HL Tau (*lower-right*). Taken from Tamura et al (1995).



## 1.8 Density Distribution

Motte & André (2001) made  $\lambda = 1.3$  mm continuum mapping survey of the embedded young stellar objects (YSOs) in the Taurus molecular cloud. Their maps include several isolated Bok globules, as well as protostellar objects in the Perseus cluster. For the protostellar envelopes mapped in Taurus, the results are roughly consistent with the predictions of the self-similar inside-out collapse model of Shu and collaborators (section 4.5). The envelopes observed in Bok globules are also qualitatively consistent with these predictions, providing the effects of magnetic pressure are included in the model. By contrast, the envelopes of Class 0 protostars in Perseus have finite radii  $\lesssim 10000$  AU and are a factor of 3 to 12 denser than is predicted by the standard model.

Another method to measure the density distribution is to use the near IR extinction. From  $(H - K)$  colors of background stars, the local value of  $A_V$  in a dark cloud can be obtained using a standard reddening law,

$$A_V = 15.87E(H - K) \quad (1.7)$$

if the intrinsic colors of background stars are known. Here, the color excess is defined as the difference between the observed color and the intrinsic color:  $E(H - K) \equiv (H - K)_{\text{obs}} - (H - K)_{\text{intrinsic}}$ . We can convert the extinction to the column density assuming the gas/dust ratio is constant

$$N(\text{H} + \text{H}_2) = 2 \times 10^{21} \text{cm}^{-2} \text{mag}^{-1} A_V. \quad (1.8)$$

This is a standard method to obtain the local column density of the dark cloud using the near IR photometry.

See Figure 1.21. If the density distribution is expressed as

$$\rho(r) = \rho_0 \left( \frac{r}{r_0} \right)^{-\alpha}, \quad (1.9)$$

where  $r$  is a physical distance from the center. The column density for the projected distance of the line-of-sight from the center of the cloud is given

$$N_\rho(p) = 2 \int_0^{(R^2 - p^2)^{1/2}} \rho \left[ (s^2 + p^2)^{1/2} \right] ds, \quad (1.10)$$

where  $R$  represents the outer radius of the cloud. Using equation (1.8), this yields  $A_V$  distribution

$$A_V(p) = 10^{-23} \rho_0 r_0^\alpha \int_0^{(R^2 - p^2)^{1/2}} (s^2 + p^2)^{-\alpha/2} ds. \quad (1.11)$$

If background stars are uniformly distributed, the number of stars with  $A_V|_{\text{obs}}$  is proportional to the area which satisfies  $A_V|_{\text{obs}} = A_V(p)$ . That is, if we plot  $A_V(p)$  against  $2\pi p dp$ , this gives the number distribution of background stars with  $A_V$ . Figure 1.23 shows the result of L977 dark cloud by Alves et al (1998).

Recently, Alves et al (2001) derived directly the radial distribution of  $N_H$  by comparing the  $N_H(p)$  model distribution for B68. They obtained a distribution is well fitted by the Bonner-Ebert sphere in which a hydrostatic balance between the self-gravity and the pressure force is achieved (lower panels of Fig.1.23) (see section 4.1.1).

In this fields, we should pay attention to the density distribution in cylindrical clouds. As seen in the Taurus molecular cloud, there are many filamentary structures in a molecular cloud. In §4.1, we will give the distribution for a hydrostatic spherical symmetric and cylindrical cloud. The former is proportional to  $\rho \propto r^{-2}$  and the latter is  $\rho \propto r^{-4}$ . Therefore, the distribution  $\rho \propto r^{-4}$  was expected



Figure 1.21: Schematic view to explain an  $A_V$  distribution.

Figure 1.22: (Left:) Radial intensity profiles of the environment of 7 embedded YSOs (a-g) and 1 starless core (h). (Right:) Same as left panel but for 4 isolated globules (a-d) and 4 Perseus protostars (e-h). Taken from Motte &amp; Andre (2001).




Figure 1.23: Density distribution of L977 (top) and B68 (bottom) dark clouds. (Top-left:) L977 dark cloud dust extinction map derived from the infrared (H-K) observations. (Top-right:) Observed




Figure 1.24: A structure of magnetic fields in the L1641 region. Polarization of light from embedded stars (Vrba et al. 1988) is shown by a bar. The direction of B-fields in the line-of-sight is observed using the HI Zeeman splitting, which is shown by a circle and cross (Heiles 1989).

for cylindrical cloud. From near IR extinctions observation (Alves et al 1998), even if a cloud is rather elongated [Fig.1.23 (top-left)], the power of the density distribution is equal to not -4 but  $\simeq -2$ . Fiege, & Pudritz (2001) proposed an idea that a toroidal component of the magnetic field,  $B_\phi$ , plays an important role in the hydrostatic balance of the cylindrical cloud (Fig.1.24).

## 1.9 Mass Spectrum

We have seen that a molecular cloud consists in many molecular cloud cores. For many years, there are attempts to determine the mass spectrum of the cores.

From a radio molecular line survey, a mass of each cloud core is determined. Plotting a histogram number of cores against the mass, we have found that a mass spectrum can be fitted by a power law as

$$\frac{dN}{dM} = M^n \quad (1.12)$$

where  $dN/dM$  represents the number of cores per unit mass interval. Many observation indicate that  $n \sim -1.5$  as Table 1.1.

Figure 1.25 (Motte et al 2001) shows the cumulative mass spectrum ( $N(> M)$  vs.  $M$ ) of the 70 starless condensations identified in NGC 2068/2071. The mass spectrum for the 30 condensations of the NGC 2068 sub-region is very similar in shape. The best-fit power-law is  $N(> M) \propto M^{n+1} \propto M^{-1.1}$  above  $M \gtrsim 0.8M_\odot$ . That is,  $n = -2.1$ . This power derived from the dust thermal emission is different from that derived by the radio molecular emission lines. The power  $n + 1 = -1.1$  which is close to the Salpeter's IMF for new born stars,  $N(> M_*) \propto M_*^{-1.35}$  might be meaningful. The reason of the difference is not clear. For example, Tothill et al. (2002) observed the Lagoon nebula

Table 1.1: Mass spectrum indices derived with molecular line surveys.

Paper	$n$	Region	Mass range
Loren (1989)	-1.1	$\rho$ Oph	$10M_{\odot} \lesssim M \lesssim 300M_{\odot}$
Stutzki & Guesten (1990)	$-1.7 \pm 0.15$	M17 SW	a few $M_{\odot} \lesssim M \lesssim$ a few $10^3 M_{\odot}$
Lada et al (1991)	-1.6	L1630	$M \gtrsim 20M_{\odot}$
Nozawa et al (1991)	-1.7	$\rho$ Oph North	$3M_{\odot} \lesssim M \lesssim 160M_{\odot}$
Tatematsu et al. (1993)	$-1.6 \pm 0.3$	Orion A	$M \gtrsim 50M_{\odot}$
Dobashi et al (1996)	-1.6	Cygnus	$M \gtrsim 100M_{\odot}$
Onishi et al (1996)	$-0.9 \pm 0.2$	Taurus	$3M_{\odot} \lesssim M \lesssim 80M_{\odot}$
Kramer et al.(1998)	$-1.6 \sim -1.8$	L1457 etc*	$10^{-4}M_{\odot} \lesssim M \lesssim 10^4 M_{\odot}$
Heithausen et al (2000)	-1.84	MCLD123.5+24.9,Polaris Flare	$M_J \lesssim M \lesssim 10M_{\odot}$
Onishi et al. (2002)	-2.5	Yaurus $\text{H}^{13}\text{CO}^+$	$3.5M_{\odot} \lesssim M \lesssim 20.1M_{\odot}$

\* MCLD126.6+24.5, NGC 1499 SW, Orion B South, S140, M17 SW, and NGC 7538




Figure 1.25: Cumulative mass distribution of the 70 pre-stellar condensations of NGC 2068/2071. The dotted and dashed lines are power-laws corresponding to the mass spectrum of CO clumps (Kramer et al. 1996) and to the IMF of Salpeter (1955), respectively. Taken from Fig.3 of Motte et al (2001).

around the edge of the HII region M8. From the continuum emission  $\lambda = 450\mu\text{m}$ ,  $850\mu\text{m}$ ,  $1.3\text{mm}$ , they obtained index of  $-1.7 \pm 0.6$ , which is consistent with other molecular line studies.

## Chapter 2

# Physical Background

### 2.1 Basic Equations of Hydrodynamics

The basic equation of hydrodynamics are (1) the continuity equation of the density [equation (A.11)],

$$\frac{\partial \rho}{\partial t} + \operatorname{div}(\rho \mathbf{v}) = 0, \quad (2.1)$$

(2) the equation of motion [equation (A.7)]

$$\rho \left[ \frac{\partial \mathbf{v}}{\partial t} + (\mathbf{v} \cdot \nabla) \mathbf{v} \right] = -\nabla p + \rho \mathbf{g}, \quad (2.2)$$

and (3) the equation of energy [equation (A.18)]

$$\frac{\partial \epsilon}{\partial t} + \operatorname{div}(\epsilon + p) \mathbf{v} = \rho \mathbf{v} \cdot \mathbf{g}. \quad (2.3)$$

Occasionally barotropic relation  $p = P(\rho)$  substitutes the energy equation (2.3). Especially polytropic relation  $p = K\rho^\Gamma$  is often used on behalf of the energy equation. In the case that the gas is isothermal or isentropic, the polytropic relations of

$$p = c_{is}^2 \rho \quad (\text{isothermal}) \quad (2.4)$$

and

$$p = c_s^2 \rho^\gamma \quad (\text{isentropic}) \quad (2.5)$$

are substitution to equation (2.3). [We can replace equation (2.3) with equations (2.4) and (2.5).]

### 2.2 The Poisson Equation of the Self-Gravity

In this section, we will show the basic equation describing how the gravity works. First, compare the gravity and the static electric force. Consider the electric field formed by a point charge  $Q$  at a distance  $r$  from the point source as

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}, \quad (2.6)$$

where  $\epsilon_0$  is the electric permittivity of the vacuum. On the other hand, the gravitational acceleration by the point mass of  $M$  at the distance  $r$  from the point mass is written down as

$$g = -G \frac{M}{r^2}, \quad (2.7)$$

where  $G = 6.67 \times 10^{-8} \text{kg}^{-1} \text{m}^3 \text{s}^{-2}$  is the gravitational constant. Comparing these two, replacing  $Q$  with  $M$  and at the same time  $1/4\pi\epsilon_0$  to  $-G$  these equations (2.6) and (2.7) are identical with each other.

The Gauss theorem for electrostatic field as

$$\text{div } \mathbf{E} = \frac{\rho_e}{\epsilon_0}, \quad (2.8)$$

and another expression using the electrostatic potential  $\phi_e$  as

$$\nabla^2 \phi_e = -\frac{\rho_e}{\epsilon_0}, \quad (2.9)$$

lead to the equations for the gravity as

$$\text{div } \mathbf{g} = -4\pi G\rho, \quad (2.10)$$

and

$$\nabla^2 \phi = 4\pi G\rho, \quad (2.11)$$

where  $\rho_e$  and  $\rho$  represent the electric charge density and the mass density. Equation (2.11) is called the Poisson equation for the gravitational potential and describes how the potential  $\phi$  is determined from the mass density distribution  $\rho$ .

### Problem

Consider a spherical symmetric density distribution. Using the Poisson equation, obtain the potential ( $\phi$ ) and the gravitational acceleration ( $g$ ) for a density distribution shown below.

$$\rho \begin{cases} = \rho_0 & \text{for } r < R \\ = 0 & \text{for } r \geq R \end{cases}$$

Hint: The Poisson equation (2.11) for the spherical symmetry is

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{\partial \phi}{\partial r} \right) = 4\pi G\rho.$$

## 2.3 Free-fall Time

If the pressure force can be neglected in the equation of motion (A.1), the gravitational one remains. Assuming the spherical symmetry, consider the gravity  $g_r(r)$  at the position of radial distance from the center being equal to  $r$ . Using the Gauss' theorem,  $g_r$  is related to the mass inside of  $r$ , which is expressed by the equation

$$M_r = \int_0^r \rho 4\pi r^2 dr, \quad (2.12)$$

and  $g_r$  is written as

$$g_r(r) = -\frac{GM_r}{r^2}. \quad (2.13)$$

This leads to the equation motion for a cold gas under the control of the self-gravity is written

$$\frac{d^2 r}{dt^2} = -\frac{GM_r}{r^2}. \quad (2.14)$$



Analyzing the equation is straightforward, multiplying  $dr/dt$  gives

$$\frac{dv^2/2}{dt} = +\frac{d}{dt} \left( \frac{GM_r}{r} \right), \quad (2.15)$$

which leads to the conservation of mechanical energy as

$$\frac{1}{2} \left( \frac{dr}{dt} \right)^2 - \frac{GM_r}{r} = E, \quad (2.16)$$

in which  $E$  represents the total energy of the pressureless gas element and it is fixed from the initial condition. If the gas is static initially at the distance  $R$ , the energy is negative as

$$E = -\frac{GM_r(R)}{R}, \quad (2.17)$$

because at  $t = 0$ ,  $r = R$  and  $dr/dt = 0$ .

The solutions of equation (2.16) are well known as follows:

1. the case of negative energy  $E < 0$ . Considering the case that the gas sphere is inflowing  $v < 0$ , equation (2.16) becomes

$$\frac{dr}{dt} = -[2GM_r(R)]^{1/2} \left( \frac{1}{r} - \frac{1}{R} \right)^{1/2}, \quad (2.18)$$

where we assumed initially  $dr/dt = 0$  at  $r = R$ . Using a parameter  $\eta(t)$ , the radius of the gas element at some epoch  $t$  is written

$$r = R \cos^2 \eta. \quad (2.19)$$

In this case, equation (2.18) reduces to

$$\cos^2 \eta \frac{d\eta}{dt} = \left( \frac{GM_r(R)}{2R^3} \right)^{1/2}. \quad (2.20)$$

This gives us the expression of  $t$  as

$$t = \left( \frac{R^3}{2GM_r(R)} \right)^{1/2} \left( \eta + \frac{\sin 2\eta}{2} \right). \quad (2.21)$$

This corresponds to the closed universe in the cosmic expansion ( $\Omega_0 < 1$ ).

2. if the energy is equal to zero, the solution of equation (2.16) is written as

$$\left( r^{3/2} - R^{3/2} \right)^{2/3} = \left( \frac{9GM_r(R)}{2} \right)^{1/3} t^{2/3}, \quad (2.22)$$

where  $R = r(t = 0)$ .

### Problem

Solve equation (2.16) and obtain (2.22).




Figure 2.1: Free-fall. x-axis and y-axis represent  $\cos^2 \eta$  and  $\eta + \sin 2\eta/2$ .

3. If  $E > 0$ , the expansion law is given by

$$t = \left( \frac{R^2}{2E} \right)^{1/2} \left( \frac{\dot{R}^2 R}{2GM_r(R)} \right)^{-1} \left( \frac{\sinh 2\eta}{2} - \eta \right) \quad (2.23)$$

and

$$x = \left( \frac{E}{GM_r(R)} \right) r = \sinh^2 \eta, \quad (2.24)$$

where  $E$  represents the total energy

$$E = \frac{\dot{R}^2}{2} - \frac{GM_r(R)}{R} > 0. \quad (2.25)$$

### Problem

Solve equation (2.16) and obtain equations (2.23) and (2.24).

In the present case, at  $t = 0$ , since  $dr/dt = 0$  the energy is negative. Equation (2.19) shows us  $r$  becomes equal to zero (the gas collapses) if  $\eta = \pi/2$  as well as  $\eta = 0$  at  $t = 0$ . Equation (2.21) indicates it occurs at the epoch of

$$\begin{aligned} t = t_{\text{ff}} &= \left( \frac{R^3}{2GM_r(R)} \right)^{1/2} \frac{\pi}{2}, \\ &= \left( \frac{3\pi}{32G\bar{\rho}} \right)^{1/2}, \end{aligned} \quad (2.26)$$

where  $\bar{\rho}$  represents the average density inside of  $r$ , that is  $M_r/(4\pi r^3/3)$ . This is called “**free-fall time**” of the gas. This gives the time-scale for the gas with density  $\bar{\rho}$  to collapse. In the actual interstellar space, the gas pressure is not negligible. However,  $t_{\text{ff}}$  gives a typical time-scale for a gas cloud to collapse and to form stars in it.

### 2.3.1 Accretion Rate

Equation 2.26 indicates that the gas shell with a large  $\bar{\rho}$  reaches the center earlier than that with a small  $\bar{\rho}$ . Immediately, this means a spherical cloud with a uniform density  $\rho_0$  contracts uniformly and

all the mass reaches the center at  $t = t_{\text{ff}} = (3\pi/32G\rho_0)^{1/2}$ . In this case, the mass accretion rate to a central source becomes infinity. In contrast, consider a cloud whose density gradually decreases outwardly. In this case, the outer mass shell has smaller  $\bar{\rho}$  than the inner mass shell. Therefore even when the inner mass shell collapses and reaches the center, the outer mass shell are contracting and does not reach the center. This gives a smaller mass accretion rate than a uniform cloud. If the gas pressure is neglected, the accretion rate is determined by the initial spatial distribution of the density. We will compare the accretion rate derived here with results of hydrodynamical calculation in §4.6

## 2.4 Gravitational Instability

Here, we will study a typical size where the self-gravity play an important role and form density inhomogeneities — the Jean wavelength.

### Linear Analysis

Consider a uniform gas with density  $\rho_0$  and pressure  $p_0$  without motion  $\mathbf{u}_0 = 0$ . In this uniform gas distribution, we assume small perturbations. As a result, the distributions of the density, the pressure and the velocity are perturbed from the uniform ones as

$$\rho = \rho_0 + \delta\rho, \quad (2.27)$$

$$p = p_0 + \delta p, \quad (2.28)$$

and

$$\mathbf{u} = \mathbf{u}_0 + \delta\mathbf{u} = \delta\mathbf{u}, \quad (2.29)$$

where the amplitudes of perturbations are assumed much small, that is,  $|\delta\rho|/\rho_0 \ll 1$ ,  $|\delta p|/p_0 \ll 1$  and  $|\delta\mathbf{u}|/c_s \ll 1$ . We assume the variables changes only in the  $x$ -direction. In this case the basic equations for isothermal gas are

$$\frac{\partial\rho}{\partial t} + \frac{\partial\rho u}{\partial x} = 0, \quad (2.30)$$

$$\rho \left( \frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} \right) = -\frac{\partial p}{\partial x} + \rho g_x, \quad (2.31)$$

and

$$p = c_{is}^2 \rho, \quad (2.32)$$

where  $u$  and  $g_x$  represent the  $x$ -component of the velocity and that of the gravity, respectively.

Using equations (2.27), (2.28), and (2.29), equation (2.30) becomes

$$\frac{\partial\rho_0 + \delta\rho}{\partial t} + \frac{\partial(\rho_0 + \delta\rho)(u_0 + \delta u)}{\partial x} = 0. \quad (2.33)$$

Noticing that the amplitudes of variables with and without  $\delta$  are completely different, two equations are obtained from equation(2.33) as

$$\frac{\partial\rho_0}{\partial t} + \frac{\partial\rho_0 u_0}{\partial x} = 0, \quad (2.34)$$

$$\frac{\partial\delta\rho}{\partial t} + \frac{\partial\rho_0\delta u + \delta\rho u_0}{\partial x} = 0, \quad (2.35)$$

where the above is the equation for unperturbed state and the lower describes the relation between the quantities with  $\delta$ . Equation (2.34) is automatically satisfied by the assumption of uniform distribution. In equation (2.35) the last term is equal to zero. Equation of motion

$$(\rho_0 + \delta\rho) \left( \frac{\partial u_0 + \delta u}{\partial t} + (u_0 + \delta u) \frac{\partial u_0 + \delta u}{\partial x} \right) = -\frac{\partial p_0 + \delta p}{\partial x} + (\rho_0 + \delta\rho) \frac{\partial \phi_0 + \delta \phi}{\partial x}, \quad (2.36)$$

gives the relationship between the terms containing only **one** variable with  $\delta$  as follows:

$$\rho_0 \frac{\partial \delta u}{\partial t} = -\frac{\partial \delta p}{\partial x} - \rho_0 \frac{\partial \delta \phi}{\partial x}. \quad (2.37)$$

The perturbations of pressure and density are related with each other as follows: for the isothermal gas

$$\frac{\delta p}{\delta \rho} = \left( \frac{\partial p}{\partial \rho} \right)_T = \frac{p_0}{\rho_0} = c_{is}^2, \quad (2.38)$$

and for isentropic gas

$$\frac{\delta p}{\delta \rho} = \left( \frac{\partial p}{\partial \rho} \right)_{ad} = \gamma \frac{p_0}{\rho_0} = c_s^2. \quad (2.39)$$

### 2.4.1 Sound Wave

If the self-gravity is ignorable, equations(2.35)

$$\frac{\partial \delta \rho}{\partial t} + \rho_0 \frac{\partial \delta u}{\partial x} = 0, \quad (2.40)$$

and equations(2.37)

$$\rho_0 \frac{\partial \delta u}{\partial t} = -c_{is}^2 \frac{\partial \delta \rho}{\partial x}, \quad (2.41)$$

where we assumed the gas is isothermal. These two equations describe the propagation and growth of perturbations. If the gas acts adiabatically, replace  $c_{is}$  with  $c_s$ .

Making  $\frac{\partial}{\partial x} \times (2.40)$  and  $\frac{\partial}{\partial t} \times (2.41)$  vanishes  $\delta \rho$  and we obtain

$$\frac{\partial^2 \delta u}{\partial t^2} - c_{is}^2 \frac{\partial^2 \delta u}{\partial x^2} = 0. \quad (2.42)$$

Since this leads to

$$\frac{\partial \delta u}{\partial t} - c_{is} \frac{\partial \delta u}{\partial x} = 0, \quad (2.43)$$

$$\frac{\partial \delta u}{\partial t} + c_{is} \frac{\partial \delta u}{\partial x} = 0, \quad (2.44)$$

equation (2.42) has a solution that a wave propagates with a phase velocity of  $\pm c_s$ . Since the displacement ( $\propto \delta u$ ) is parallel to the propagation direction  $x$ , and the restoring force comes from the pressure, this seems **the sound wave**.

### Problem

Interstellar gas contains mainly Hydrogen and Helium, whose number ratio is approximately 10:1. Obtain the value of average molecular weight for the fully ionized interstellar gas with temperature  $T = 10^4 \text{K}$  (components are ionized  $\text{H}^+$  (HII) and  $\text{He}^{+2}$  (HeIII) and electron  $e^{-1}$ ). How about the molecular gas ( $T = 10 \text{K}$ ) containing molecular  $\text{H}_2$ , neutral He (HeI) and no electron.

## 2.5 Jeans Instability

Sound wave seems to be modified in the medium where the self-gravity is important. Beside the continuity equation (2.35)

$$\frac{\partial \delta \rho}{\partial t} + \rho_0 \frac{\partial \delta u}{\partial x} = 0, \quad (2.45)$$

and the equation of motion (2.37)

$$\rho_0 \frac{\partial \delta u}{\partial t} = -c_{is}^2 \frac{\partial \delta \rho}{\partial x} - \rho_0 \frac{\partial \delta \phi}{\partial x}, \quad (2.46)$$

the linearized Poisson equation

$$\frac{\partial^2 \delta \phi}{\partial x^2} = 4\pi G \delta \rho, \quad (2.47)$$

should be included.  $\frac{\partial}{\partial x} \times \text{eq.}(2.46)$  gives

$$\rho_0 \left( \frac{\partial^2 \delta u}{\partial x \partial t} \right) = -c_{is}^2 \frac{\partial^2 \delta \rho}{\partial x^2} - 4\pi G \rho_0 \delta \rho. \quad (2.48)$$

where we used equation (2.47) to eliminate  $\delta \phi$ . This yields

$$\frac{\partial^2 \delta \rho}{\partial t^2} = c_{is}^2 \frac{\partial^2 \delta \rho}{\partial x^2} + 4\pi G \rho_0 \delta \rho. \quad (2.49)$$

where we used  $\frac{\partial}{\partial t} \times \text{eq.}(2.45)$ .

This is the equation which characterizes the growth of density perturbation owing to the self-gravity. Here we consider the perturbation are expressed by the linear combination of plane waves as

$$\delta \rho(x, t) = \sum A(\omega, k) \exp(i\omega t - ikx), \quad (2.50)$$

where  $k$  and  $\omega$  represent the wavenumber and the angular frequency of the wave. Picking up a plane wave and putting into equation (2.49), we obtain **the dispersion relation** for the gravitational instability as

$$\omega^2 = c_{is}^2 k^2 - 4\pi G \rho_0. \quad (2.51)$$

Reducing the density to zero, the equation gives us the same dispersion relation as that of sound wave as  $\omega/k = c_{is}$ . For short waves ( $k \gg k_J = (4\pi G \rho_0)^{1/2}/c_{is}$ ), since  $\omega^2 > 0$  the wave is ordinary oscillatory wave. Increasing the wavelength (decreasing the wavenumber),  $\omega^2$  becomes negative for  $k < k_J = (4\pi G \rho_0)^{1/2}/c_{is}$ . For negative  $\omega^2$ ,  $\omega$  can be written  $\omega = \pm i\alpha$  using a positive real  $\alpha$ . In this case, since  $\exp(i\omega t) = \exp(\mp \alpha t)$ , the wave which has  $\omega = -i\alpha$  increases its amplitude exponentially. This means that even if there are density inhomogeneities only with small amplitudes, they grow in a time scale of  $1/\alpha$  and form the density inhomogeneities with large amplitudes.

The critical wavenumber

$$k_J = (4\pi G \rho_0)^{1/2}/c_{is} \quad (2.52)$$

corresponds to the wavelength

$$\lambda_J = \frac{2\pi}{k_J} = \left( \frac{\pi c_{is}^2}{G \rho_0} \right)^{1/2}, \quad (2.53)$$

which is called the Jeans wavelength. Ignoring a numerical factor of the order of unity, it is shown that the Jeans wavelength is approximately equal to the free-fall time scale (eq.2.26) times the sound




Figure 2.2: Dispersion Relation




Figure 2.3: Thin disk.

speed. The short wave with  $\lambda \ll \lambda_J$  does not suffer from the self-gravity. For such a scale, the analysis we did in the preceding section is valid.

Typical values in molecular clouds, such as  $c_{is} = 200\text{m s}^{-1}$ ,  $\rho_0 = 2 \times 10^{-20}\text{g cm}^{-3}$ , give us the Jeans wavelength as  $\lambda_J = 1.7 \times 10^{18}\text{cm} = 0.56\text{pc}$ . The mass contained in a sphere with a radius  $r = \lambda_J/2$  is often called Jeans mass, which gives a typical mass scale above which the gas collapses. Typical value of the Jeans mass is as follows

$$M_J \simeq \frac{4\pi}{3} \rho_0 \left(\frac{\lambda_J}{2}\right)^3 = \frac{\pi}{6} \left(\frac{\pi}{G\rho_0}\right)^{3/2} c_{is}^3 \rho_0. \quad (2.54)$$

Using again the above typical values in the molecular clouds,  $c_{is} = 200\text{m s}^{-1}$ ,  $\rho_0 = 2 \times 10^{-20}\text{g cm}^{-3}$ , the Jeans mass of this gas is equal to  $M_J \simeq 27M_\odot$ .

## 2.6 Gravitational Instability of Thin Disk

Disks are common in the Universe. Spiral and barred spiral galaxies have disks where stars are formed. In more small scale, gas and dust disks are often found around protostars. Moreover, the

disk may become a proto-planetary disk. It is valuable to study how the self-gravity works in such thin structures. Here, we assume a thin disk extending in  $x$ - and  $y$ -directions whose surface density is equal to  $\sigma = \int_{-\infty}^{\infty} \rho dz$ , in other word the density is written using the Dirac's delta function  $\delta(z)$  as

$$\rho(x, y, z) = \sigma(x, y)\delta(z). \quad (2.55)$$

Integrating along the  $z$ -direction basic equations (2.45), (2.46), and (2.47), the linearized basic equations for the thin disk are as follows:

$$\frac{\partial \delta \sigma}{\partial t} + \sigma_0 \frac{\partial \delta u}{\partial x} = 0, \quad (2.56)$$

$$\sigma_0 \frac{\partial \delta u}{\partial t} = -c_{is}^2 \frac{\partial \delta \sigma}{\partial x} - \sigma_0 \frac{\partial \delta \phi}{\partial x}, \quad (2.57)$$

$$\frac{\partial^2 \delta \phi}{\partial x^2} + \frac{\partial^2 \delta \phi}{\partial z^2} = 4\pi G \delta \sigma \delta(z), \quad (2.58)$$

where we assumed  $\sigma = \sigma_0 + \delta \sigma$ ,  $u = \delta u$ ,  $\phi = \phi_0 + \delta \phi$  and took the first order terms (those contain only one  $\delta$ ).

Outside the disk, the rhs of equation (2.58) is equal to zero. It reduces to the Laplace equation

$$\frac{\partial^2 \delta \phi}{\partial x^2} + \frac{\partial^2 \delta \phi}{\partial z^2} = 0. \quad (2.59)$$

Taking a plane wave of

$$\delta X(x, t) = \delta A \exp(i\omega t - ikx), \quad (2.60)$$

equation (2.59) is reduced to

$$\frac{\partial^2 \delta \phi}{\partial z^2} - k^2 \delta \phi = 0. \quad (2.61)$$

This has a solution which does not diverge at the infinity  $z = \pm\infty$  as

$$\delta \phi = \delta \phi(z=0) \exp(-k|z|). \quad (2.62)$$

On the other hand, integrating equation (2.58) from  $z = -0$  to  $z = +0$  or in other word, applying the Gauss' theorem to the region containing the  $z = 0$  surface, it is shown that the gravity  $\delta g_z = -\partial \delta \phi / \partial z$  has a jump crossing the  $z = 0$  surface as

$$\left. \frac{\partial \delta \phi}{\partial z} \right|_{z=+0} - \left. \frac{\partial \delta \phi}{\partial z} \right|_{z=-0} = 4\pi G \delta \sigma. \quad (2.63)$$

Equations (2.62) and (2.63) lead a final form of the potential as

$$\delta \phi = -\frac{2\pi G \delta \sigma}{k} \exp(-k|z|). \quad (2.64)$$

Putting this to equations (2.57), and using equations (2.56) and (2.57), we obtain the dispersion relation for the gravitational instability in a thin disk as

$$\omega^2 = c_{is}^2 k^2 - 2\pi G \sigma_0 k. \quad (2.65)$$

This reduces to the dispersion relation of the sound wave for the short wave  $k \gg 2\pi G \sigma_0 / c_{is}^2$ . While for a longer wave than  $\lambda_{cr} = c_{is}^2 / G \sigma_0$ , an exponential growth of  $\delta \sigma$  is expected. The dispersion relation is shown in Fig.2.2.




Figure 2.4: *Left:* Explanation of Laval nozzle. *Right:* The relation between the cross-section  $S(x)$  and the flow velocity  $v_x$ .

## 2.7 Super- and Subsonic Flow

Flow whose velocity is faster than the sound speed is called supersonic, while that slower than the sound speed is called subsonic. The subsonic and supersonic flows are completely different.

### 2.7.1 Flow in the Laval Nozzle

Consider a tube whose cross-section,  $S(x)$ , changes gradually, which is called Laval nozzle. Assuming the flow is steady  $\partial/\partial t = 0$  and is essentially one-dimensional, the continuity equation (2.1) is rewritten as

$$\rho u S = \text{constant}, \quad (2.66)$$

or

$$\frac{1}{\rho} \frac{\partial \rho}{\partial x} + \frac{1}{u} \frac{\partial u}{\partial x} + \frac{1}{S} \frac{\partial S}{\partial x} = 0. \quad (2.67)$$

Equation of motion (2.2) becomes

$$u \frac{\partial u}{\partial x} = -\frac{1}{\rho} \frac{\partial p}{\partial x} = -\frac{c_s^2}{\rho} \frac{\partial \rho}{\partial x}, \quad (2.68)$$

where we used the relationship of

$$\frac{\partial p}{\partial x} = \left( \frac{\partial p}{\partial \rho} \right)_s \frac{\partial \rho}{\partial x} = c_s^2 \frac{\partial \rho}{\partial x}. \quad (2.69)$$

When the flow is isothermal, use the isothermal sound speed  $c_{is}^2$  instead of the adiabatic one. From equations (2.67) and (2.68), we obtain

$$\left( \frac{u^2}{c_s^2} - 1 \right) \frac{1}{u} \frac{\partial u}{\partial x} = \frac{1}{S} \frac{\partial S}{\partial x}, \quad (2.70)$$

where the factor  $\mathcal{M} = u/c_s$  is called **the Mach number**. For supersonic flow  $\mathcal{M} > 1$ , while  $\mathcal{M} < 1$  for subsonic flow.

In the supersonic regime  $\mathcal{M} > 1$ , the factor in the parenthesis of the lhs of equation (2.69) is positive. This leads to the fact that the velocity increases ( $du/dx > 0$ ) as long as the cross-section






Figure 2.5: *Left:* External field potential. *Right:* Velocity and density variations. Gas flows in the external field whose potential is shown in the left panel. The upper panel represents a subsonic flow. The lower panel does a supersonic flow.

increases ( $dS/dx > 0$ ). On the other hand, in the subsonic regime, the velocity decreases ( $du/dx < 0$ ) while the cross-section increases ( $dS/dx > 0$ ). See right panel of 2.4.

If  $\mathcal{M} = 1$  at the point of minimum cross-section (throat), two curves for  $\mathcal{M} < 1$  and  $\mathcal{M} > 1$  have an intersection. In this case, gas can be accelerated through the Laval nozzle. First, a subsonic flow is accelerated to the sonic speed at the throat of the nozzle. After passing the throat, the gas follows the path of a supersonic flow, where the velocity is accelerated as long as the cross-section increases.

### 2.7.2 Steady State Flow under an Influence of External Fields

Consider the flow under the force exerted on the gas whose strength varies spatially. Let  $g(x)$  represent the force working per unit mass. Assuming the cross-section is constant

$$\rho u = \text{constant}, \quad (2.71)$$

immediately we have

$$\frac{1}{\rho} \frac{\partial \rho}{\partial x} + \frac{1}{u} \frac{\partial u}{\partial x} = 0. \quad (2.72)$$

On the other hand, the equation motion is

$$u \frac{\partial u}{\partial x} = -\frac{c_s^2}{\rho} \frac{\partial \rho}{\partial x} + g(x), \quad (2.73)$$

From equations (2.72) and (2.73), we obtain

$$\left( \frac{u^2}{c_s^2} - 1 \right) \frac{1}{u} \frac{\partial u}{\partial x} = \frac{g(x)}{c_s^2}. \quad (2.74)$$

Consider an external field whose potential is shown in Fig.2.5(*Left*). (1) For subsonic flow, the factor in the parenthesis is negative. Before the potential minimum, since  $g(x) > 0$ ,  $u$  is decelerated. On the other hand, after the potential minimum,  $u$  is accelerated owing to  $g(x) < 0$ . Using equation (2.71), this leads to a density distribution in which density peaks near the potential minimum. (2) For supersonic flow, the factor is positive. In the region of  $g(x) > 0$ ,  $u$  is accelerated. After passing the potential minimum,  $u$  is decelerated. The velocity and the density distribution is shown in Fig.2.5(right-lower panel).

The density distribution of the subsonic flow in an external potential is similar to that of hydrostatic one. That is, considering the hydrostatic state in an external potential, the gas density peaks at the potential minimum. On the other hand, The density distribution of the supersonic flow looks like that made by ballistic particles which are moving freely in the potential. Owing to the conservation of the total energy (kinetic + potential energies), the velocity peaks at the density minimum. And the condition of mass conservation leads to the distribution in which the density decreases near the potential minimum.

### 2.7.3 Stellar Wind — Parker Wind Theory

Stellar winds are observed around various type of stars. Early type (massive) stars have large luminosities; the photon absorbed by the bound-bound transition transfers its outward momentum to the gas. This line-driven mechanism seems to work around the early type stars. On the other hand, acceleration mechanism of less massive stars are thought to be related to the coronal activity or dust driven mechanism (dusts absorb the emission and obtain outward momentum from the emission).

Here, we will see the identical mechanism in §2.7.1 and §2.7.2 works to accelerate the wind from a star. Consider a steady state and ignore  $\partial/\partial t = 0$ . The continuity equation (2.1) gives

$$r^2 \rho u = \text{const}, \quad (2.75)$$

where we used

$$\text{div} \rho \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho u). \quad (2.76)$$

This leads to

$$\frac{2}{r} + \frac{1}{r} \frac{\partial \rho}{\partial r} + \frac{1}{u} \frac{\partial u}{\partial r} = 0. \quad (2.77)$$

The equation of motion (2.2) is as follows:

$$u \frac{\partial u}{\partial r} = -\frac{c_s^2}{\rho} \frac{\partial \rho}{\partial r} - \frac{GM_*}{r^2}, \quad (2.78)$$

where we used  $g = -GM_*/r^2$  ( $M_*$  means the mass of the central star). From these two equations (2.77) and (2.78) we obtain

$$\left( \mathcal{M}^2 - 1 \right) \frac{1}{u} \frac{\partial u}{\partial r} = \frac{2}{r} - \frac{GM_*}{c_s^2} \frac{1}{r^2}, \quad (2.79)$$




Figure 2.6: Right-hand side of equation (2.79) is plotted against the distance from the center  $r$ .

where  $\mathcal{M}$  represents the Mach number of the radial velocity. Take notice that this is similar to equations (2.70) and (2.74). That is, the fact that the rhs of equation (2.79) is positive corresponds to increasing the cross-section  $dS/dx > 0$ . On the contrary, when the rhs is negative, the fluid acts as the cross-section  $S$  is decreasing.

For simplicity, we assume the gas is isothermal. The rhs of equation (2.79) varies shown in Figure 2.6. Therefore, near to the star, the flow acts as the cross-section of nozzle is decreasing and far from the star it does as the cross-section is increasing. This is the same situation that the gas flows through the Laval nozzle.

Using a normalized distance  $x \equiv r/(GM_*/2c_{is}^2)$ , equation (2.79) becomes

$$(M^2 - 1) \frac{1}{\mathcal{M}} \frac{\partial \mathcal{M}}{\partial x} = \frac{2}{x} - \frac{2}{x^2}. \quad (2.80)$$

This is rewritten as

$$\frac{d}{dx} \left( \frac{\mathcal{M}^2}{2} - \log \mathcal{M} - 2 \log x - \frac{2}{x} \right) = 0, \quad (2.81)$$

we obtain the solution of equation (2.80) as

$$\mathcal{M}^2 - 2 \log \mathcal{M} = 4 \log x + \frac{4}{x} + C. \quad (2.82)$$

This gives how the Mach number  $\mathcal{M}$  varies changing  $x$ . To explore this, we define two functions:

$$f(\mathcal{M}) = 2\mathcal{M}^2 - 2 \log \mathcal{M} \quad (2.83)$$

is a function only depending on  $\mathcal{M}$  and

$$g(x) = 4 \log x + \frac{4}{x} + C \quad (2.84)$$

is a function only depending on  $x$ .

Since the minima of  $f(\mathcal{M})$  and  $g(x)$  are respectively 1 and  $C + 4$ , the permitted region in  $(x, \mathcal{M})$  changes for values of  $C$ .




Figure 2.7:  $f(\mathcal{M})$  (left) and  $g(x)$  for  $C = -3$  (right)

1. If  $C = -3$ , for all values of  $x > 0$  there exist  $\mathcal{M}$  which satisfies  $f(\mathcal{M}) = g(x)$ . This corresponds to the two curves which pass through a critical X-point of  $(x, \mathcal{M}) = (1, 1)$  in Figure 2.8.
2. If  $C < -3$ , the minimum of  $g(x)$  is smaller than that of  $f(\mathcal{M})$ . In this case, for  $x$  where  $g(x) < 1 = \min(f(\mathcal{M}))$ , there is no solution. Thus,  $f(\mathcal{M}) = g(x)$  has solutions for  $x < x_1$  and  $x > x_2$ , where  $x_1 < x_2$ , and  $g(x_1) = g(x_2) = 1$ . This corresponds to the curves running perpendicularly in Figure 2.8.
3. If  $C > -3$ , the minimum of  $f(\mathcal{M})$  is smaller than that of  $g(x)$ . In this case,  $f(\mathcal{M}) = g(x)$  has solutions for  $\mathcal{M} < \mathcal{M}_1$  and  $\mathcal{M} > \mathcal{M}_2$ , where  $\mathcal{M}_1 < \mathcal{M}_2$ , and  $f(\mathcal{M}_1) = f(\mathcal{M}_2) = C + 4$ . This corresponds to curves running horizontally in Figure 2.8.

Out of the two solutions of  $C = -3$ , a wind solution is one with increasing  $\mathcal{M}$  while departing from the star. This shows us the outflow speed is slow near the star but it is accelerated and a supersonic wind blows after passing the critical point. Since the equations are unchanged even if we replace  $u$  with  $-u$ , the above solution is valid for an accreting flow  $u < 0$ . Considering such a flow, the solution running from the lower-right corner to the upper-left corner represents the accretion flow, in which the inflow velocity is accelerated reaching the star and finally accretes on the star surface with a supersonic speed.




Figure 2.8:  $\mathcal{M}$  vs  $x$  (the  $x$ -axis is the normalized distance from the center  $x \equiv r/(GM_*/2c_{is}^2)$  and the  $y$ -axis is the Mach number  $\mathcal{M}$ .)



## Chapter 3

# Galactic Scale Star Formation

### 3.1 Schmidt Law

Schmidt (1959) speculated that the star formation rate is proportional to a power of the surface density of the interstellar medium

$$\dot{\Sigma}_{\text{SF}} \propto \Sigma_{\text{gas}}^n, \quad (3.1)$$

where the power  $n$  seems between 1 and 2 around the solar vicinity. If  $n = 2$ , the star formation rate is thought to be determined by the collision rate of interstellar clouds. At that time Schmidt showed us  $n \simeq 2$ . On the other hand, if the gas passing through the galactic arms forms stars, the star formation rate seems proportional to the gas surface density and the arm-to-arm period. Thus this predicts  $n = 1$ .

#### 3.1.1 Global Star Formation

The star formation rate is estimated by the intensity of H $\alpha$  emission (Kennicutt, Tamblyn, & Congdon 1994) as

$$\text{SFR}(M_{\odot}\text{yr}^{-1}) = \frac{L(\text{H}\alpha)}{1.26 \times 10^{41} \text{erg s}^{-1}}, \quad (3.2)$$

which is used for normal galaxies. While in **the starburst galaxies** which show much larger star formation rate than the normal galaxies, FIR luminosity seems a better indicator of star formation rate

$$\text{SFR}(M_{\odot}\text{yr}^{-1}) = \frac{L(\text{FIR})}{2.2 \times 10^{43} \text{erg s}^{-1}} = \frac{L(\text{FIR})}{5.8 \times 10^9 L_{\odot}}. \quad (3.3)$$

Kennicutt (1998) summarized the relation between SFRs and the surface gas densities [Fig.3.1 (left)] for 61 normal spiral and 36 infrared-selected starburst galaxies. As seen in Fig.3.1, the star formation rate averaged over a galaxy ( $\Sigma_{\text{SFR}}(M_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2})$ ), which is called the global star formation rate, is well correlated to the average gas surface density  $\Sigma_{\text{gas}}(M_{\odot} \text{ pc}^{-2})$ . He gave the power of the global Schmidt law as  $n = 1.4 \pm 0.15$ . That is,

$$\Sigma_{\text{SFR}} \simeq (1.5 \pm 0.7) \times 10^{-4} \left( \frac{\Sigma_{\text{gas}}}{1 M_{\odot} \text{ kpc}^{-2}} \right)^{1.4 \pm 0.15} M_{\odot} \text{ yr}^{-1} \text{ pc}^{-2}. \quad (3.4)$$

The fact that the power is not far from 3/2 seems to be explained as follows: Star formation rate should be proportional to the gas density ( $\Sigma_{\text{gas}}$ ) but it should also be inversely proportional to the time scale of forming stars in respective gas clouds, which is essentially the free-fall time scale.




FIG. 6.—Composite star formation law for the normal disk (*filled circles*) and starburst (*squares*) samples. Open circles show the SFRs and gas densities for the centers of the normal disk galaxies. The line is a least-squares fit with index  $N = 1.40$ . The short, diagonal line shows the effect of changing the scaling radius by a factor of 2.




FIG. 7.—Relation between the SFR for the normal disk and starburst samples and the ratio of the gas density to the disk orbital timescale, as described in the text. The symbols are the same as in Fig. 6. The line is a median fit to the normal disk sample, with the slope fixed at unity as predicted by equation (7).

Figure 3.1: Taken from Figs.6 and 7 of Kennicutt (1998). Left: The  $x$ -axis means the total ( $\text{HI}+\text{H}_2$ ) gas density and the  $y$ -axis does the global star formation rate. Right: The  $x$ -axis means the total ( $\text{HI}+\text{H}_2$ ) gas density divided by the orbital time-scale. The  $y$ -axis is the same.

Remember the fact that the free-fall time given in equation (2.26) is proportional to  $\tau_{\text{ff}} \propto 1/(G\rho)^{1/2}$ . Therefore

$$\rho_{\text{SFR}} \propto \rho_{\text{gas}} \times (G\rho_{\text{gas}})^{1/2} \propto \rho_{\text{gas}}^{3/2}, \quad (3.5)$$

where  $\rho_{\text{gas}}$  and  $\rho_{\text{SFR}}$  are the volume densities of gas and star formation.

He found another correlation between the quantity of gas surface density divided by the orbital period of galactic rotation and the star formation rate [Fig.3.1 (right)]. Although the actual slope is equal to 0.9 instead of 1, the correlation in Fig.3.1(right) is well expressed as

$$\Sigma_{\text{SFR}} \simeq 0.017 \Sigma_{\text{gas}} \Omega_{\text{gas}} = 0.21 \frac{\Sigma_{\text{gas}}}{\tau_{\text{arm-to-arm}}}, \quad (3.6)$$

where  $\Omega_{\text{gas}}$  represents the angular speed of galactic rotation. This means that 21 % of the gas mass is processed to form stars per orbit. These two correlations [eqs (3.4) and (3.6)] implicitly ask another relation of  $\Omega_{\text{gas}} \propto \Sigma_{\text{gas}}^{1/2}$ .

### 3.1.2 Local Star Formation Rate

In Figure 3.2 (left), the correlation between star formation rate and gas density is plotted for specific galaxies (NGC 4254 and NGC 2841). This shows us that  $\text{H}\alpha$  surface brightness (star formation







Figure 3.2: Distributions of  $\Sigma_{\text{SF}}$  and  $\Sigma_{\text{gas}}$ . (Left:) Relation between  $\Sigma_{\text{SF}}$  and  $\Sigma_{\text{gas}}$  for an Sc galaxy NGC 4254 and an Sb galaxy NGC 2841. (Right:) Relation between  $\Sigma_{\text{SF}}$  and  $\Sigma_{\text{gas}}$  for various galaxies. These are taken from Figs.7 and 8 of Kennicutt (1989).

rate) and the gas column density are well correlated each other. Figure 3.2 (left) also indicates that there seems a critical gas density below which star formation is not observed. The value of this threshold column density is approximately  $\simeq 4M_{\odot}\text{pc}^{-2}$  for both galaxies in Figure 3.2 (left). The same correlation is seen in other spiral galaxies [Fig.3.2(right)]. Fitting the correlation with a power-law, he obtained

$$\Sigma_{\text{SFR}} \propto \Sigma_{\text{gas}}^{1.3 \pm 0.3}, \quad (3.7)$$

for the region active in star formation. Take notice that this power is very close to that of the global Schmidt law [eq.(3.4)] The threshold surface gas density ranges from  $1 M_{\odot}\text{pc}^{-2}$  to  $10M_{\odot}\text{pc}^{-2}$  ( $10^{20} - 10^{21}\text{H cm}^{-2}$ ). Therefore, theory of star formation must explain (1) the Schmidt law (clear correlation between star formation rate and the gas surface density) above the threshold column density and (2) the fact that there is no evidence for star formation in the gas deficient region below the threshold column density.

### 3.2 Gravitational Instability of Rotating Thin Disk

Here, we will derive the dispersion relation for the gravitational instability of a rotating thin disk. We will see the spatial variation of Toomre's  $Q$  parameter, which determines the stability of the rotating disk, explains the nonlinearity of star formation rate, that is, there is a threshold density and no stars are formed in the low density region.

Use the cylindrical coordinate  $(R, Z, \phi)$  and the basic equations for thin disk in §2.6. In linear analysis, we assume  $\Sigma(R, \phi) = \Sigma_0(R) + \delta\Sigma(R, \phi)$ ,  $u(R, \phi) = 0 + \delta u(R, \phi)$ ,  $v(R, \phi) = v_0(R) + \delta v(R, \phi)$ , where  $u$  and  $v$  represent the radial and azimuthal component of the velocity. Linearized continuity equation is

$$\frac{\partial \delta \Sigma}{\partial t} + \frac{1}{R} \frac{\partial}{\partial R} (R \Sigma_0 \delta u) + \Omega \frac{\partial \delta \Sigma}{\partial \phi} + \frac{\Sigma_0}{R} \frac{\partial \delta v}{\partial \phi} = 0, \quad (3.8)$$

where  $\Omega = v_0/R$ .

Linearized equations of motion are

$$\left( \frac{\partial}{\partial t} + \Omega \frac{\partial}{\partial \phi} \right) \delta u - 2\Omega \delta v = -\frac{\partial}{\partial R} (\delta \Phi + \delta h), \quad (3.9)$$

and

$$\left( \frac{\partial}{\partial t} + \Omega \frac{\partial}{\partial \phi} \right) \delta v + \frac{\kappa^2}{2\Omega} \delta u = -\frac{1}{R} \frac{\partial}{\partial \phi} (\delta \Phi + \delta h), \quad (3.10)$$

where  $h$  is a specific enthalpy as  $dh = dp/\Sigma$  and

$$\kappa^2 = 4\Omega^2 + R \frac{d\Omega^2}{dR} \quad (3.11)$$

is **the epicyclic frequency**.

We assume any solution of equations (3.8), (3.9) and (3.10) can be written as a sum of terms of the form

$$\delta u = u_a \exp[i(m\phi - \omega t)], \quad (3.12)$$

$$\delta v = v_a \exp[i(m\phi - \omega t)], \quad (3.13)$$

$$\delta \Sigma = \Sigma_a \exp[i(m\phi - \omega t)], \quad (3.14)$$

$$\delta h = h_a \exp[i(m\phi - \omega t)], \quad (3.15)$$

$$\delta \Phi = \Phi_a \exp[i(m\phi - \omega t)]. \quad (3.16)$$

Using the equation of state of  $p = K\Sigma^\gamma$ ,

$$h_a = c_s^2 \Sigma_a / \Sigma_0. \quad (3.17)$$

Using equations (3.12)-(3.16), equations (3.8), (3.9), and (3.10) are rewritten as

$$i(m\Omega - \omega)\Sigma_a + \frac{1}{R} \frac{\partial}{\partial R} (R \Sigma_0 u_a) + im \frac{\Sigma_0 v_a}{R} = 0, \quad (3.18)$$

$$u_a [\kappa^2 - (m\Omega - \omega)^2] = -i \left[ (m\Omega - \omega) \frac{d}{dR} (\Phi_a + h_a) + 2m\Omega \frac{(\Phi_a + h_a)}{R} \right], \quad (3.19)$$

and

$$v_a [\kappa^2 - (m\Omega - \omega)^2] = \left[ \frac{\kappa^2}{2\Omega} \frac{d}{dR} (\Phi_a + h_a) + m(m\Omega - \omega) \frac{(\Phi_a + h_a)}{R} \right], \quad (3.20)$$




Figure 3.3: Tightly wound (left) vs loosely wound (right) spirals.

### 3.2.1 Tightly Wound Spirals

We assume the wave driven by the self-gravity has a form of tightly-wound spiral [Fig.3.3(left)]. When we move radially, the density  $\delta\Sigma$  varies rapidly. While, it changes its amplitude slowly in the azimuthal direction. In a mathematical expression, if we write the density perturbation  $\delta\Sigma$  as

$$\delta\Sigma = A(R, t) \exp[im\phi + i f(R, t)], \quad (3.21)$$

where the amplitude of spiral  $A(R, t)$  is a slowly varying function of  $R$ , a **tightly wound spiral** means the shape function varies fast (the radial wavenumber  $k \simeq df/dR$  is large enough). We consider the gravitational force from the vicinity of  $(R_0, \phi_0)$ , since the  $\delta\Sigma$  oscillates and cancels even if we integrate over large region. Thus,

$$\delta\Sigma(R, \phi, t) \simeq \Sigma_a \exp[ik(R_0, t)(R - R_0)], \quad (3.22)$$

where

$$\Sigma_a = A(R_0, t) \exp[im\phi_0 + f(R_0, t)]. \quad (3.23)$$

Notice that the density perturbation [eq.(3.22)] is similar to that studied in §2.6. The potential should be expressed in a similar form to equation (2.64) as

$$\delta\Phi \simeq -\frac{2\pi G \Sigma_a}{|k|} \exp[ik(R_0, t)(R - R_0)], \quad (3.24)$$

which simply means

$$\Phi_a = -\frac{2\pi G \Sigma_a}{|k|}. \quad (3.25)$$

If we set  $R = R_0$ , we obtain our final result for the potential due to the surface density perturbation

$$\delta\Phi(R, \phi, t) \simeq -\frac{2\pi G}{|k|} A(R, t) \exp[im\phi + f(R, t)]. \quad (3.26)$$

Differentiating this equation with  $R$  and ignoring the term  $dA(R, t)/dR$  compared to that of  $df(R, t)/dR = k$ , we obtain

$$\delta\Sigma(R, \phi, t) = i \frac{\text{sign}(k)}{2\pi G} \frac{d\delta\Phi(R, \phi, t)}{dR}, \quad (3.27)$$

Table 3.1: Epicyclic frequency vs rotation law.

Rotation	$\kappa$
Rigid-body rotation $\Omega = \text{const.}$	$2\Omega$
Flat rotation $v_\phi = \text{const.}$	$\sqrt{2}\Omega$
Kepler rotation $v_\phi \propto r^{-1/2}$	$\Omega$

Neglecting the terms  $\propto 1/R$  compared to the terms containing  $\partial/\partial R$ , equations (3.18), (3.19), and (3.20) are rewritten as

$$i(m\Omega - \omega)\Sigma_a + ik\Sigma_0 u_a = 0, \quad (3.28)$$

$$u_a[\kappa^2 - (m\Omega - \omega)^2] = (m\Omega - \omega)k(\Phi_a + h_a), \quad (3.29)$$

and

$$v_a[\kappa^2 - (m\Omega - \omega)^2] = i\frac{\kappa^2}{2\Omega}k(\Phi_a + h_a), \quad (3.30)$$

Using these equations [(3.28), (3.29), and (3.30)],  $\Phi_a = -2\pi G\Sigma_a/|k|$ , and  $h_a = c_s^2\Sigma_a/\Sigma_0$ , we obtain the dispersion relation for the self-gravitating instability of the rotating gaseous thin disk

$$(m\Omega - \omega)^2 = k^2 c_s^2 - 2\pi G\Sigma_0|k| + \kappa^2. \quad (3.31)$$

Generally speaking, the epicyclic frequency depends on the rotation law but is in the range of  $\Omega \lesssim \kappa \lesssim 2\Omega$  (see Table 3.1 for  $\kappa$  for typical rotation laws). It is shown that the system is stabilized due to the the epicyclic frequency compared with a nonrotating thin disk [eq.(3.38)].

### 3.2.2 Toomre's $Q$ Value

Consider the case of  $m = 0$  axisymmetric perturbations. Equation (3.31) becomes

$$\omega^2 = k^2 c_s^2 - 2\pi G\Sigma_0|k| + \kappa^2 = c_s^2 \left( k - \frac{\pi G\Sigma_0}{c_s^2} \right)^2 + \kappa^2 - \left( \frac{\pi G\Sigma_0}{c_s} \right)^2. \quad (3.32)$$

If  $\omega^2 > 0$  the system is stable against the axisymmetric perturbation, while if  $\omega^2 < 0$  the system is unstable. Defining

$$Q = \frac{\kappa c_s}{\pi G\Sigma_0}, \quad (3.33)$$

if  $Q > 1$ ,  $\omega^2 > 0$  for all wavenumbers  $k$ . On the other hand, if  $Q < 1$ ,  $\omega^2$  becomes negative for some wavenumbers  $k_1 < k < k_2$ . Therefore, the Toomre's  $Q$  number gives us a criterion whether the system is unstable or not for the axisymmetric perturbation. [Recommendation for a reference book of this section: Binney & Tremaine (1988).]

The condition is expressed as

$$\Sigma_0 > \Sigma_{\text{cr}} = \frac{\kappa c_s}{\pi G} \quad (Q < 1). \quad (3.34)$$

Kennicutt plotted  $\Sigma_0/\Sigma_{\text{cr}}$  against the normalized radius as  $R/R_{\text{HII}}$  for various galaxies, where  $R_{\text{HII}}$  represents the maximum distance of HII regions from the center (Fig.3.4). Since  $\Sigma_0/\Sigma_{\text{cr}} = Q$ , Figure shows that HII regions are observed mainly in the region with  $Q < 1$  but those are seldom seen in the outer low-density  $Q > 1$  region. This seems the gravitational instability plays an important role.




Figure 3.4:  $\Sigma_0/\Sigma_{cr}$  vs  $R/R_{HII}$ .  $R_{HII}$  represents the maximum distance of HII regions from the center. The sound speed is assumed constant  $c_s = 6\text{km s}^{-1}$ . Taken from Fig.11 of Kennicutt (1989).

The above discussion is for the gaseous disk. The Toomre's  $Q$  value is also defined for stellar system as

$$Q = \frac{\sigma_R \kappa}{3.36 G \Sigma_0}, \quad (3.35)$$


where  $\sigma_R$  represents the radial velocity dispersion.

For non-axisymmetric waves, even if  $1 \lesssim Q \lesssim 2$  the instability grows. To explain this, **the swing amplification** mechanism is proposed (Toomre 1981). If there is a leading spiral perturbation in the disk with  $1 \lesssim Q \lesssim 2$ , the wave unwinds and finally becomes a trailing spiral pattern. At the same time, the amplitude of the wave (perturbations) is amplified (see Fig.3.5).

### 3.3 Spiral Structure

Fig.3.6 shows the  $B$ - (left) and  $I$ -band images of M51.  $B$  light which originates from the massive early type stars. Although the image taken in  $B$ -band shows a number of spiral arms, that of  $I$ -band shows clearly two arms. The  $I$ -band light seems to come from mainly less-massive long-lived stars, while the  $B$ -band light is essentially coming from the massive short-lived stars which are formed in the spiral arm. On the contrary, the less-massive stars are not necessarily born in the spiral arm. This suggests that there are two kinds of spiral patterns: one is made by stars (mainly less-massive) and the other is the gaseous spiral arm where massive stars are born and contribute to the  $B$ -band image.

In this section, first, we will briefly describe the **density wave theory** which explains the former spiral pattern in stellar component. You will find the amplitude of the spiral pattern in stellar component is not so large. However, the response of gaseous components (HI and  $H_2$  gas) to the spiral density wave potential with small amplitude is much more nonlinear than that of stellar component and a high-contrast spiral pattern appears in gaseous component .



**Figure 6-17.** Evolution of a packet of leading waves in a stellar Mestel disk with  $Q = 1.5$  and  $f = 1$ . Contours represent fixed fractional excess surface densities; since the calculations are based on linear perturbation theory, the amplitude normalization is arbitrary. Contours in regions of depleted surface density are not shown to minimize confusion. The time interval between diagrams is one-half of a rotation period at corotation. From Toomre (1981), by permission of Cambridge University Press.

Figure 3.5: Numerical simulation of the swing amplification mechanism. The number attached each panel shows the time sequence. This is obtained by the time-dependent linear analysis. First, perturbation with leading spiral pattern is added to the Mestel disk with  $Q = 1.5$ . The leading spiral gradually unwinds and become a trailing spiral. Loosely wound spiral pattern winds gradually and the last panel shows a tightly wound leading spiral pattern. The final amplitude is  $\sim 100$  times larger than that of the initial state.

### 3.4 Density Wave Theory

We have derived the dispersion relation of the gravitational instability in the rotating thin disk as

$$(m\Omega - \omega)^2 = k^2 c_s^2 - 2\pi G \Sigma_0 |k| + \kappa^2, \quad (3.36)$$

where  $m$  represents the number of spiral arms. Although the stability of the stellar system is a little different, we assume this is valid for the stellar system after  $c_s$  is replaced to the velocity dispersion. Since

$$(m\Omega - \omega)^2 = k^2 c_s^2 - 2\pi G \Sigma_0 |k| + \kappa^2 = c_s^2 \left( k - \frac{\pi G \Sigma_0}{c_s^2} \right)^2 + \kappa^2 - \left( \frac{\pi G \Sigma_0}{c_s} \right)^2, \quad (3.37)$$

we obtain

$$|k| = k_T \frac{2}{Q^2} \left[ 1 \pm \sqrt{1 - Q^2(1 - \nu^2)} \right], \quad (3.38)$$

where  $\Omega_P = \omega/m$  is a pattern speed,  $\nu = m(\Omega_P - \Omega)/\kappa$  is the normalized frequency,  $k_T = \kappa^2/2\pi G \Sigma_0$  is the Toomre's critical wavenumber for a cold ( $c_s = 0$ ) system.  $\nu = \pm 1$ , which leads to  $|k| = 0$ ,




Figure 3.6:  $B$ - (left), and  $I$ -band (right) images of M51. Taken from Fig. 3 of Elmegreen et al. (1989).




Figure 3.7: (*Left:*) A plot of the dispersion relation eq.(3.38).  $x$ - and  $y$ -axes are  $\nu$  and  $k/k_T$ . Respective lines are for  $Q = 1$  (straight lines),  $Q = 1.2$ ,  $Q = 1.5$ , and  $Q = 2$ . The trailing part  $k > 0$  is only plotted. Points  $\nu = -1$ ,  $\nu = 0$ , and  $\nu = 1$  correspond respectively to Inner Lindblad Resonance (ILR), Corotation Resonance (CR), and Outer Lindblad Resonance (OLR). The relation is symmetric against the  $x$ -axis and the curve of  $k < 0$  represents the leading wave. (*Right:*) Leading vs trailing spiral.

represents **the Lindblad resonance** and is rewritten as

$$\Omega_{\text{P}} = \frac{\omega}{m} = \Omega \pm \frac{\kappa}{2}. \quad (3.39)$$

Assuming  $m = 2$ , the resonance when  $\Omega_{\text{P}} = \Omega + \kappa/2$  is called outer Lindblad resonance while that of  $\Omega_{\text{P}} = \Omega - \kappa/2$  is called inner Lindblad resonance.  $\nu = 0$  means **the co-rotation resonance**  $\Omega_{\text{P}} = \Omega$ .

Writing down equation (3.38) [Fig.3.7(left)], it is shown that, in the case of  $Q = 1$ , the wavenumber exists for all  $\nu$ . Since  $\nu = -1, 0$ , and  $+1$  correspond to the points of ILR, CR, and OLR and these three resonance points appear in accordance with the radial distance, the  $x$ -axis of Figure 3.7(left) seems to correspond to the radial distance from the center. In the case of  $Q > 1$ , it is shown that a forbidden region appears around the co-rotation resonance point. Waves cannot propagate into the region. Figure 3.7(left) shows that the  $k/k_{\text{T}}$  has two possible wavenumbers in the permitted region. The waves with larger  $k$  and smaller  $k$  are called short waves and long waves.

Consider a wave expressed by  $\Sigma \propto \exp[im\phi + ikr]$ . If  $k < 0$ , moving from a point  $(R_0, \phi_0)$  in the direction  $\Delta\phi > 0$  and  $\Delta r > 0$  the phase difference between the two points  $[m(\phi_0 + \Delta\phi) + k(R_0 + \Delta r)] - [m\phi_0 + kR_0]$  can be equal to zero. That is, in the case of  $k < 0$  the wave is leading. On the other hand, if  $k > 0$ , moving in the direction  $\Delta\phi < 0$  and  $\Delta r > 0$  the phase will be unchanged. In this case the wave pattern is trailing. Since the dispersion relation is symmetric for  $k > 0$  and  $k < 0$ , there are two waves, trailing waves and leading waves. Therefore there are four waves: a short trailing wave, a long trailing wave, a short leading wave, and a long leading wave.

### Group Velocity

The wave transfers the energy with the group velocity. Whether the group velocity is positive or negative is quite important considering the energy transfer. Using the dispersion relation, equation (3.37), The group velocity

$$v_g(R) = \frac{d\omega}{dk} = \text{sign}(k) \frac{|k|c_s^2 - \pi G\Sigma_0}{\omega - m\Omega}. \quad (3.40)$$

For a region  $R > R_{\text{CR}}$ ,  $\omega - m\Omega > 0$  or  $\nu > 0$ . On the other hand, for a region  $R < R_{\text{CR}}$ ,  $\omega - m\Omega < 0$  or  $\nu < 0$ . Consider first the trailing wave. In the region  $R > R_{\text{CR}}$ , long-waves propagate inwardly to the CR, since  $|k|c_s^2 < \pi G\Sigma_0$  for long-waves and  $\nu > 0$ . Short-waves propagate outwardly from the CR, since  $|k|c_s^2 > \pi G\Sigma_0$  for short-waves and  $\nu > 0$ . In the region  $R < R_{\text{CR}}$ , long-waves propagate outwardly to the CR since  $|k|c_s^2 < \pi G\Sigma_0$  and  $\nu < 0$ . Short-waves propagate inwardly from the CR, since  $|k|c_s^2 > \pi G\Sigma_0$  and  $\nu < 0$ . As a result, it is concluded that the long-wave propagates toward the CR and the short-wave does away from the CR. As for the leading wave, the short-wave propagates toward the CR and the long-wave does away from the CR.

Assuming that the wave packet is made near the Lindblad resonance points, (1) the long-trailing waves propagate toward the co-rotation resonance points; (2) they are reflected by the  $Q$ -barrier; (3) they change their character to short-waves and propagate away from the co-rotation resonance points; and finally (4) the waves are absorbed at the center or propagate away to the infinity. (1) the short-leading waves propagate toward the co-rotation resonance points; (2) they are reflected by the  $Q$ -barrier; (3) they change their character to long-waves and propagate away from the co-rotation resonance points; and finally (4) they reach the Lindblad resonance points and the energy may be converted to the long-trailing waves there.

The wave obtains its energy at the resonance points. The density wave transfer the energy to the co-rotation points. Therefore, the density wave theory predicts the galactic stellar disk has spiral density pattern between the inner Lindblad resonance points and the outer Lindblad resonance point if  $Q \gtrsim 1$ .






Figure 3.8: Spiral coordinate. Along a curve of constant  $\eta$ ,  $r \propto \exp(\theta \tan i)$ .

### 3.5 Galactic Shock

In the preceding section, we have seen that a spiral pattern of density perturbation is made by the gravitational instability in the stellar system with  $Q \gtrsim 1$ . This forms a grand design spiral observed in the *I*-band images which represent the mass distribution which consists essentially in less-massive stars. The amplitude of the pattern is smaller than that observed in *B*-band images. In this section we will see how the distribution early type stars is explained. [Recommendation for a reference book of this section: Spitzer (1978).]

Gas flowing through a spiral gravitational potential, even if its amplitude is relatively small, acts rather nonlinearly. Consider the gravitational potential in the sum of an axisymmetric term,  $\Phi_0(R)$  and a spiral term  $\Phi_1(R)$ . The spiral gravitational field is rotating with a pattern speed  $\Omega_P$ . We use a reference frame rotating with  $\Omega_P$ .  $\mathbf{u} = (u, v)$ , where  $u$  and  $v$  are radial ( $R$ ) and azimuthal ( $\phi$ ) component of the flow velocity. Consider the reference system rotating with the angular rotation speed  $\Omega_P$ . We assume the variables with suffix 0 represent those without the spiral potential  $\Phi_1$  and variables with suffix 1 are used to express the difference between before and after  $\Phi_1$  is added.  $u = u_0 + u_1 = 0 + u_1$ ,  $v = v_0 + v_1$ , and

$$v_0 = R(\Omega - \Omega_P) = R \left[ \left( \frac{1}{R} \frac{\partial \Phi_0}{\partial R} \right)^{1/2} - \Omega_P \right], \quad (3.41)$$

where  $\Omega$  means the circular rotation speed. The equation of motion in the  $r$ -direction is

$$\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial r} + \frac{v}{r} \frac{\partial u}{\partial \theta} - \frac{v_1^2}{r} = -\frac{c_s^2}{\Sigma} \frac{\partial \Sigma}{\partial r} - \frac{\partial \Phi_1}{\partial r} + 2\Omega v_1, \quad (3.42)$$

and that in the  $\phi$  direction is

$$\frac{\partial v}{\partial t} + u \left( \frac{\partial v_1}{\partial r} + \frac{v_1}{r} \right) + \frac{v}{r} \frac{\partial v_1}{\partial \phi} = -\frac{c_s^2}{\Sigma r} \frac{\partial \Sigma}{\partial \phi} - \frac{1}{r} \frac{\partial \Phi_1}{\partial \phi} - \frac{\kappa^2}{2\Omega} u, \quad (3.43)$$

and the continuity equation is

$$\frac{\partial \Sigma R u}{\partial R} + \frac{\partial \Sigma v}{\partial \phi} = 0. \quad (3.44)$$

As seen in Figure 3.8, we introduce the spiral coordinate  $(\eta, \xi)$  in which  $\xi$ -axis is parallel to the spiral pattern which has a pitch angle  $i$  and  $\eta$ -axis is perpendicular to the  $\xi$ -axis.

$$rd\eta = dr \cos i + rd\phi \sin i, \quad (3.45)$$

$$rd\xi = -dr \sin i + rd\phi \cos i. \quad (3.46)$$

$$v_\eta = u \cos i + v \sin i, \quad (3.47)$$

$$v_\xi = -u \sin i + v \cos i \quad (3.48)$$

Assuming that  $i \ll 1$  (tightly wound spiral), equations (3.44) (3.42) and (3.43) become

$$\frac{\partial v_\eta}{\partial t} + v_\eta \frac{\partial v_\eta}{\partial \eta} = -\frac{c_s^2}{\Sigma} \frac{\partial \Sigma}{\partial \eta} - \frac{\partial \Phi_1}{\partial \eta} + 2\Omega(v_\xi - v_{\xi 0}), \quad (3.49)$$

$$\frac{\partial v_\xi}{\partial t} + v_\eta \frac{\partial v_\xi}{\partial \eta} = -\frac{\kappa^2}{2\Omega}(v_\eta - v_{\eta 0}), \quad (3.50)$$

$$\frac{\partial \Sigma}{\partial t} + \frac{\partial \Sigma v_\eta}{\partial \eta} = 0. \quad (3.51)$$

Similar to §2.7 we look for a steady state solution. Equation (3.51) becomes

$$\frac{d\Sigma}{d\eta} = -\frac{\Sigma}{v_\eta} \frac{dv_\eta}{d\eta}. \quad (3.52)$$

Using this, equation (3.49) reduces to

$$(v_\eta^2 - c_s^2) \frac{dv_\eta}{d\eta} = 2\Omega(v_\xi - v_{\xi 0}) - \frac{d\Phi_1}{d\eta}. \quad (3.53)$$

Equation (3.50) becomes

$$v_\eta \frac{dv_\xi}{d\eta} = -\frac{\kappa^2}{2\Omega}(v_\eta - v_{\eta 0}), \quad (3.54)$$

In these equations we used following quantities:

$$v_\eta = v_{\eta 0} + v_{\eta 1}, \quad (3.55)$$

$$v_\xi = v_{\xi 0} + v_{\xi 1}, \quad (3.56)$$

$$v_{\eta 0} = v_0 \sin i, \quad (3.57)$$

$$v_{\xi 0} = v_0 \cos i, \quad (3.58)$$

$v_0$  is given in equation (3.41).

Equations (3.52), (3.53), and (3.54) are solved under the periodic boundary condition:  $X(\eta = \text{right end}) = X(\eta = \text{left end})$ . Since equation (3.53) is similar to the equations in §2.7, you may think the solution seems like Figure 2.5. However, it contains a term which expresses the effect of Coriolis force  $2\Omega(v_\xi - v_{\xi 0})$ , the flow becomes much complicated.

Taking care of the point that a shock front exists for a range of parameters, the solution of the above equations are shown in Figure 3.9. Numerical hydrodynamical calculations which solve equations (3.49), (3.50), and (3.51) was done and steady state solutions are obtained (Woodward 1975). It is shown that  $F \leq 0.7\%$ , the velocity ( $v_\eta$ : velocity perpendicular to the wave) does not show any discontinuity. In contrast, for  $F > 2\%$ , a shock wave appears. Steady state solution is obtained from the ordinary differential equations (3.52) (3.53), and (3.54) by Shu, Milione, &

Roberts (1973). Inside the CR,  $v_0 > 0$  (Gas has a faster rotation speed than the spiral pattern). As long as  $v_\eta > c_s$  there is no shock. Increasing the amplitude of the spiral force  $F$ , an amplitude of the variation in  $v_\eta$  increases and finally  $v_\eta$  becomes subsonic partially. When the flow changes its nature from supersonic to subsonic, it is accompanied with a shock. (An inverse process, that is, changing from subsonic to supersonic is not accompanied with a shock.)

In the outer galaxy (still inside CR) since  $\Omega - \Omega_P$  decreases with the the distance from the galactic center,  $v_{\eta 0} = R(\Omega - \Omega_P) \sin i$  decreases. In this region,  $v_{\eta 0} < c_{is}$  and the flow is subsonic if there is no spiral gravitational force,  $F = 0$ . In such a circumstance, increasing  $F$  amplifies the variation in  $v_\eta$  and  $v_\eta$  finally reaches the sound speed. Transonic flow shows again a shock.

Summary of this section is:

1. There exists a spiral density pattern of the stellar component if the Toomre's  $Q$  parameter is  $1 \lesssim Q \lesssim 2$ .
2. This is driven by the self-gravity of the rotating thin disk.
3. If the amplitude of the non-axisymmetric force is as large as  $\sim 1\%$  of the axisymmetric one, interstellar gas whose sound speed is as large as  $c_{is} \sim 8 \text{ km s}^{-1}$  forms the galactic shock. This is observed when the flow is transonic.
4. The amplitude of the **gas** density fluctuation is much larger than that in the stellar density.
5. If stars are formed preferentially in the postshock region of the spiral arm, we expect clear spiral arms made by early-type stars, which are massive and short-lived, as seen in the  $B$  band images of spiral galaxies.




Figure 3.9: Galactic shocks. The velocity perpendicular to the spiral wave front is plotted against the phase of the spiral wave. The spiral potential takes its minimum at  $\theta = 90$ .  $c_s = 8.6\text{km s}^{-1}$ ,  $i = 6.7$  deg,  $R = 10\text{kpc}$ ,  $R\Omega = 250\text{km s}^{-1}$ ,  $R\kappa = 313\text{km s}^{-1}$ ,  $R\Omega_P = 135\text{km s}^{-1}$ . The amplitude of spiral gravity force is taken  $F = 0.4\%$ ,  $0.7\%$ ,  $2\%$ , and  $5\%$  of the axisymmetric force  $\partial\Phi_0/\partial R$ . Taken from Fig.13.3 of Spizer (1978) [originally Woodward (1975)].




FIG. 1.—Entirely supersonic flow in the velocity plane for  $\omega = 10$  kpc with  $a = 8$  km s<sup>-1</sup>. The unperturbed velocity is marked by a cross, and the flow proceeds in a counterclockwise direction. For  $F = 0.97\%$ , the flow contains a cusp at the location of the minimum of the spiral potential  $\eta = 0$ . FIG. 2.—Transonic flow in the velocity plane for  $\omega = 10$  kpc with  $a = 8$  km s<sup>-1</sup>. The unperturbed velocity is marked by a cross, and the flow proceeds in a counterclockwise direction. For  $F > F_{sup} = 0.97\%$ , the flow contains a shock located at the indicated value of  $\eta$ .




FIG. 4

Figure 3.10: Galactic shock. (Left:)  $v_\eta$  vs  $v_\xi$  plot. Increasing  $F$  from 0.5% to 0.9%, the variation in  $v_\eta$  increases. And at  $F = 0.97\%$ , the minimum speed reaches the sound speed  $8\text{ km s}^{-1}$ . Further increasing  $F$ , a subsonic region appears ( $v_\eta < c_{is}$  for  $F = 1.05\%$  and  $5\%$ ). When flow changes its nature from supersonic to subsonic, a shock appears.  $R = 10\text{ kpc}$ ,  $c_s = 8\text{ km s}^{-1}$ ,  $i = 6.7$  deg. Other parameters are the same as Fig.3.9. (Right:) Comparison of two density distributions:  $F = 0.97$  and  $F = 5\%$  (shock). Taken from Shu, Milione, & Roberts (1973).



## Chapter 4

# Local Star Formation Process

### 4.1 Hydrostatic Balance

Consider a hydrostatic balance of isothermal cloud. By the gas density,  $\rho$ , the isothermal sound speed,  $c_{\text{is}}$ , and the gravitational potential,  $\Phi$ , the force balance is written as

$$-\frac{c_{\text{is}}^2}{\rho} \frac{d\rho}{dr} - \frac{d\Phi}{dr} = 0, \quad (4.1)$$

and the gravity is calculated from a density distribution as

$$-\frac{d\Phi}{dr} = -\frac{GM_r}{r^2} = -\frac{4\pi G}{r^2} \int_0^r \rho r^2 dr, \quad (4.2)$$

for a spherical symmetric cloud, where  $M_r$  represents the mass contained inside the radius  $r$ . The expression for a cylindrical cloud is

$$-\frac{d\Phi}{dr} = -\frac{G\lambda_r}{r} = -\frac{2\pi G}{r} \int_0^r \rho r dr, \quad (4.3)$$

where  $\lambda_r$  represents the mass per unit length within a cylinder of radius being  $r$ .

For the spherical symmetric case, the equation becomes the Lane-Emden equation with the polytropic index of  $\infty$  (see Appendix C.1). This has no analytic solutions. However, the numerical integration gives us a solution shown in Figure 4.1 (left). Only in a limiting case with the infinite central density, the solution is expressed as

$$\rho(r) = \frac{c_{\text{is}}^2}{2\pi G} r^{-2}. \quad (4.4)$$

Increasing the central density, the solution reaches the above Singular Isothermal Sphere (SIS) solution.

On the other hand, a cylindrical cloud has an analytic solution (Ostriker 1964) as

$$\rho(r) = \rho_c \left(1 + \frac{r^2}{8H^2}\right)^{-2}, \quad (4.5)$$

where

$$H^2 = c_{\text{is}}^2 / 4\pi G \rho_c. \quad (4.6)$$




Figure 4.1: Radial density distribution. A spherical cloud (left) and a cylindrical cloud (right). In the right panel, solutions for polytropic gases with  $\Gamma = 1.1$  (relatively compact) and  $\Gamma = 0.9$  (relatively extended) are plotted as well as the isothermal one.

Far from the cloud symmetric axis, the distribution of equation (4.5) gives

$$\rho(r) \propto r^{-4}, \quad (4.7)$$

while the spherical symmetric cloud has

$$\rho(r) \propto r^{-2} \quad (4.8)$$

distribution.

### Problem 1

Show that the SIS is a solution of equations (4.1) and (4.2).

### Problem 2

Show that the density distribution of [equation (4.5)] is a solution of equations (4.1) and (4.3).

#### 4.1.1 Bonnor-Ebert Mass

In the preceding section [Fig.4.1 (left)], we have seen the radial density distribution of a hydrostatic configuration of an isothermal gas. Consider a circumstance that such kind of cloud is immersed in a low-density medium with a pressure  $p_0$ . To establish a pressure equilibrium, the pressure at the surface  $c_{\text{is}}^2 \rho(R)$  must equal to  $p_0$ . This means that the density at the surface is constant  $\rho(R) = p_0/c_{\text{is}}^2$ .

Figure 4.2 (left) shows three models of density distribution,  $\rho_c = \rho(r=0) = 10\rho_s$ ,  $10^2\rho_s$ , and  $10^3\rho_s$ . Comparing these three models, it should be noticed that the cloud size (radius) decreases with increasing the central density  $\rho_c$ . The mass of the cloud is obtained by integrating the distribution, which is illustrated against the central-to-surface density ratio  $\rho_c/\rho_s$  in Figure 4.2 (right). The  $y$ -axis represents a normalized mass as  $m = M/[4\pi\rho_s(c_{\text{is}}/\sqrt{4\pi G\rho_s})^3]$ . The maximum value of  $m = 4.026$  means

$$M_{\text{max}} \simeq 1.14 \frac{c_{\text{is}}^2}{G^{3/2} p_0^{1/2}}. \quad (4.9)$$



This is the maximum mass which is supported against the self-gravity by the thermal pressure with an isothermal sound speed of  $c_{\text{is}}$ , when the cloud is immersed in the pressure  $p_0$ . This is called Bonnor-Ebert mass [Bonnor (1956), Ebert (1955)]. It is to be noticed that the critical state  $M_{\text{cl}} = M_{\text{max}}$  is achieved when the density contrast is rather low  $\rho_c \simeq 16\rho_s \equiv \rho_{\text{cr}}$ .

Another important result from Figure 4.2 (right) is the stability of an isothermal cloud. Even for a cloud with  $M_{\text{cl}} < M_{\text{max}}$ , any clouds on the part of  $\partial M_{\text{cl}}/\partial \rho_c < 0$  are unstable, whose clouds are distributed on the branch with  $\rho_c > \rho_{\text{cr}}$ . This is understood as follows: For a hydrostatic cloud the mass should be expressed with the external pressure and the central density [Fig.4.2 (right)] as

$$M_{\text{cl}} = M_{\text{cl}}(p_{\text{ext}}, \rho_c). \quad (4.10)$$

In this case, a relation between the partial derivatives such as

$$\left(\frac{\partial M_{\text{cl}}}{\partial p_{\text{ext}}}\right)_{\rho_c} \cdot \left(\frac{\partial p_{\text{ext}}}{\partial \rho_c}\right)_{M_{\text{cl}}} \cdot \left(\frac{\partial \rho_c}{\partial M_{\text{cl}}}\right)_{p_{\text{ext}}} = -1, \quad (4.11)$$

is satisfied, unless each term is equal to zero. Figure 4.1 (left) shows that the cloud mass  $M_{\text{cl}}$  is a decreasing function of the external pressure  $p_{\text{ext}} = \rho_s c_{\text{is}}^2$ , if the central density is fixed. Since this means

$$\left(\frac{\partial M_{\text{cl}}}{\partial p_{\text{ext}}}\right)_{\rho_c} < 0, \quad (4.12)$$

equation (4.11) gives us

$$\left(\frac{\partial p_{\text{ext}}}{\partial \rho_c}\right)_{M_{\text{cl}}} \cdot \left(\frac{\partial \rho_c}{\partial M_{\text{cl}}}\right)_{p_{\text{ext}}} > 0. \quad (4.13)$$

For a cloud with  $\rho_c < \rho_{\text{cr}} = 16p_{\text{ext}}/c_{\text{is}}^2$  the mass is an increasing function of the central density as

$$\left(\frac{\partial M_{\text{cl}}}{\partial \rho_c}\right)_{p_{\text{ext}}} > 0. \quad (4.14)$$

Thus, equation (4.13) leads to the relation

$$\left(\frac{\partial \rho_c}{\partial p_{\text{ext}}}\right)_{M_{\text{cl}}} > 0, \quad (4.15)$$

for  $\rho < \rho_{\text{cr}}$ . This means that gas cloud contracts (the central density and pressure increase), when the external pressure increases. This is an ordinary reaction of a stable gas.

On the other hand, the cloud on the part of  $(\partial M_{\text{cl}}/\partial \rho_c)_{p_{\text{ext}}} < 0$  (for  $16 \lesssim \rho_c/(p_{\text{ext}}/c_{\text{is}}^2) \lesssim 2000$ ) behaves

$$\left(\frac{\partial \rho_c}{\partial p_{\text{ext}}}\right)_{M_{\text{cl}}} < 0, \quad (4.16)$$

and this represents that an extra external pressure decreases the central density and thus the pressure. Pressure encourages expansion. This reaction is unstable.

#### 4.1.2 Equilibria of Cylindrical Cloud

In Figure 4.1 (right) we plotted the structure for a polytropic cloud. Inner structure is not dependent of  $\Gamma$ , it is clear the slope of the outer envelope is dependent on  $\Gamma$ .




Figure 4.2: (Left) radial density distribution. Each curve has different  $\rho_c$ . The x-axis denotes the radial distance normalized by a scale-length as  $c_{\text{is}}^2/\sqrt{4\pi G p_{\text{ext}}}$ . It is shown that the radius increases with decreasing  $\rho_c$  in this range ( $\rho_c = (10, 10^2, 10^3) \times p_{\text{ext}}/c_{\text{is}}^2$ ). (Right) The relation between mass ( $M_{\text{cl}}$ ) and the central density ( $\rho_c$ ) is plotted, under the condition of constant external pressure. The x-axis represents the central density normalized by  $\rho_s \equiv p_{\text{ext}}/c_{\text{is}}^2$ . The y-axis represents the cloud mass normalized by  $4\pi\rho_s(c_{\text{is}}/\sqrt{4\pi G\rho_s})^3$ .

1. In the case of the spherically symmetric, consider a polytrope ( $p \propto \rho^\Gamma$ ) with  $\Gamma < 6/5$  (at least the envelope of  $\Gamma = 6/5$  cloud extends to  $\infty$ ), in which gas extends to  $\infty$ . if  $\rho \propto r^{-p}$ , the mass inside of  $r$  is proportional to  $M_r \propto r^{3-p}$ . Thus, the gravity per unit volume at a radius  $r$ ,  $GM\rho/r^2$ , is proportional to  $GM\rho/r^2 \propto r^{1-2p}$ . On the other hand the pressure force is  $|(\partial p/\partial r)| = (\partial p/\partial \rho)|(\partial \rho/\partial r)| \propto (r^{-p})^{\Gamma-1} r^{-p-1} \propto r^{-p\Gamma-1}$ . These two powers become the same, only if  $p = 2/(2 - \Gamma)$ .
2. In the case of cylindrical cloud with  $\Gamma \leq 1$ , the mass per unit length  $\lambda \propto r^{2-p}$ . The gravity at  $r$ ,  $G\lambda\rho/r \propto r^{1-2p}$ . Note that the power is the same as the spherical case. Since the power of the pressure force should be the same as the spherical case, the resultant  $p$  should be the same  $p = 2/(2 - \Gamma)$ .

The case of  $\Gamma = 0.9$ , an envelope extending to a large radius indicates the power-law distribution much shallower than that of the isothermal  $\Gamma = 1$  one.

## 4.2 Virial Analysis

Hydrodynamic equation of motion using the Lagrangean time derivative [eq.(A.3)] is

$$\rho \left( \frac{d\mathbf{v}}{dt} \right) = -\nabla p - \rho \nabla \Phi. \quad (4.17)$$

Multiplying the position vector  $\mathbf{r}$  and integrate over a volume of a cloud, we obtain the Virial relation as

$$\frac{1}{2} \frac{d^2 I}{dt^2} = 2(T - T_0) + W, \quad (4.18)$$

where

$$I = \int \rho r^2 dV, \quad (4.19)$$

is an inertia of the cloud,

$$T = \int \left( \frac{3}{2} p_{\text{th}} + \frac{1}{2} \rho v^2 \right) dV = \frac{3}{2} \bar{p} V_{\text{cl}}, \quad (4.20)$$

is a term corresponding to the thermal pressure plus turbulent pressure,

$$T_0 = \int_S P_{\text{th}} \mathbf{r} \cdot \mathbf{n} dS = \frac{3}{2} P_0 V_{\text{cl}} \quad (4.21)$$

comes from a surface pressure, and

$$W = - \int \rho \mathbf{r} \cdot \nabla \Phi dV = - \frac{3}{5} \frac{GM^2}{R} \quad (4.22)$$

is a gravitational energy. To derive the last expression in each equation, we have assumed the cloud is spherical and uniform. Here we use a standard notation as the radius  $R$ , the volume  $V_{\text{cl}} = 4\pi R^3/3$ , the average pressure  $\bar{P}$ , and the mass  $M$ .

To obtain a condition of mechanical equilibrium, we assume  $d^2 I/dt^2 = 0$ . Equation (4.18) becomes

$$4\pi \bar{p} R^3 - 4\pi p_0 R^3 - \frac{3}{5} \frac{GM^2}{R} = 0. \quad (4.23)$$

Assuming the gas is isothermal  $p = c_{\text{is}}^2 \rho$ , the average pressure is written as

$$\bar{p} = c_{\text{is}}^2 \bar{\rho} = c_{\text{is}}^2 \frac{3M}{4\pi R^3}. \quad (4.24)$$

Using equation (4.24) to eliminate  $\bar{p}$  from equation (4.18), the external pressure is related to the mass and the radius as

$$p_0 = \frac{3c_{\text{is}}^2 M}{4\pi R^3} - \frac{3GM^2}{20\pi R^4}. \quad (4.25)$$

Keeping  $M$  constant and increasing  $R$  from zero,  $p_0$  increases first, but it takes a maximum,  $p_{0,\text{max}} = 3.15c_{\text{is}}^8/(G^3 M^2)$ , and finally declines. This indicates that the surface pressure must be smaller than  $p_{0,\text{max}}$  for the cloud to be in an equilibrium. In other words, keeping  $p_0$  and changing  $R$ , it is shown that  $M$  has a maximum value to have a solution. The maximum mass is equal to

$$M_{\text{max}} = 1.77 \frac{c_{\text{is}}^4}{G^{3/2} p_0^{1/2}}. \quad (4.26)$$

The cloud massive than  $M_{\text{max}}$  cannot be supported against the self-gravity. This corresponds to the Bonnor-Ebert mass [eq.(4.9)], although the numerical factors are slightly different.

### 4.2.1 Magnatohydrostatic Clouds

Consider here the effect of the magnetic field. In the magnetized medium, the Lorentz force

$$\mathbf{F} = \frac{1}{4\pi} (\nabla \times \mathbf{B}) \times \mathbf{B} = -\frac{1}{8\pi} \nabla B^2 + \frac{1}{4\pi} (\mathbf{B} \cdot \nabla) \mathbf{B} \quad (4.27)$$

works in the ionized medium. The first term of equation (4.27), which is called the **magnetic pressure**, has an effect to support the cloud against the self-gravity.

The virial analysis is also applicable to the magnetohydrostatic clouds. The terms related to the magnetic fields are

$$\begin{aligned} M &= \int \frac{B^2}{8\pi} dV + \int_S (\mathbf{r} \cdot \mathbf{B}) \mathbf{B} \cdot \mathbf{n} dS - \int_S \frac{B^2}{8\pi} \mathbf{r} \cdot \mathbf{n} dS \\ &\simeq \int \frac{B^2 - B_0^2}{8\pi} dV \simeq \frac{1}{6\pi^2} \left( \frac{\Phi_B^2}{R} - \frac{\Phi_B^2}{R_0} \right), \end{aligned} \quad (4.28)$$

where  $\Phi_B$  represents a magnetic flux and it is assumed to be conserved if we change the radius,  $R$ , that is  $\Phi_B = \pi B_0 R_0^2 = \pi B R^2$ . Equation (4.23) becomes

$$4\pi\bar{p}R^3 - 4\pi p_0 R^3 - \frac{3GM^2}{5R} + \frac{1}{6\pi^2} \frac{\Phi_B^2}{R} = 0, \quad (4.29)$$

where we ignored the term  $\frac{1}{6\pi^2} \frac{\Phi_B^2}{R_0}$ . The last two terms are rewritten as

$$\frac{3G}{5R} (M^2 - M_\Phi^2), \quad (4.30)$$

where  $M_\Phi$  is defined as  $3GM_\Phi^2/5 = \Phi_B^2/6\pi^2$ .

This shows the effects of the magnetic fields:

1. B-fields effectively reduce the gravitational mass as  $M^2 - M_\Phi^2 = M^2 - 5\Phi_B^2/(18\pi^2G)$ . This plays a part to support a cloud.
2. However, even a cloud contracts (decreasing its radius from  $R_0$  to  $R$ ), the ratio of the gravitational to the magnetic terms keeps constant since these two terms are proportional to  $\propto R^{-1}$ . Thus, if the magnetic term does not work initially, the gravitational term continues to predominate over the magnetic term.

If  $M < M_\Phi$ , a sum of last two terms in equation (4.29) is positive. Since the second term of rhs of equation (4.25) is positive, there is one  $R$  which satisfies equation (4.29) irrespective of the external pressure  $p_0$ . While, if  $M > M_\Phi$ , there is a maximum allowable external pressure  $p_0$ . Therefore,  $M = M_\Phi$  gives a criterion whether the magnetic fields work to support the cloud or not. More realistic calculation [Mouschovias (1976a,1976b), Tomisaka et al (1988)] gives us a criterion

$$G^{1/2} \frac{dm}{d\Phi_B} = \frac{G^{1/2}\sigma}{B} = 0.17 \simeq \frac{1}{2\pi}, \quad (4.31)$$

where  $\sigma$  and  $B$  means the column density and the magnetic flux density. A cloud with a mass

$$M > \frac{\Phi_B}{2\pi G^{1/2}} \quad (4.32)$$

is sometimes called magnetically supercritical, while that with

$$M < \frac{\Phi_B}{2\pi G^{1/2}} \quad (4.33)$$

is subcritical.

More precisely speaking, the criterion showed in equations (4.32) and (4.33) should be applied for a cloud which has a much larger mass than the Bonnor-Ebert mass. That is, even without magnetic fields, the cloud less-massive than the Bonnor-Ebert mass has a hydrostatic configuration shown in

Figure 4.2 (left). The cloud with central density of  $\rho_c = 10$  has a stable density distribution. To fit the numerical results, Tomisaka et al (1988) obtained an expression for the critical mass when the cloud has a mass-to-flux ratio  $dm/d\Phi_B$ , the isothermal sound speed  $c_{is}$ , and the external pressure  $p_0$  as

$$M_{cr} = 1.3 \left\{ 1 - \left[ \frac{1/2\pi}{G^{1/2} dm/d\Phi_B|_{r=0}} \right]^2 \right\}^{-3/2} \frac{c_s^4}{p_0^{1/2} G^{3/2}}. \quad (4.34)$$

This shows that the critical mass is a decreasing function of the mass-to-flux ratio or increasing function of the magnetic flux. And the critical mass becomes much larger than the Bonnor-Ebert mass  $\simeq c_s^4/(p_0^{1/2} G^{3/2})$  only when the mass-to-flux ratio at the center of the cloud is reaching  $1/2\pi$  at which the term in the curly bracket goes to zero. Hereafter, we call here the cloud/cloud core with mass larger than the critical mass  $M_{cr}$  a supercritical cloud/cloud core. The cloud/cloud core less-massive than the critical mass is subcritical.

### 4.3 Subcritical Cloud vs Supercritical Cloud

We have seen there is a critical mass above which the cloud has no (magneto)hydrostatic configuration but below which the cloud has at least an equilibrium state. This gives us an idea that there are two kind of clouds/cloud cores: that with a mass larger than the critical mass which has to collapse dynamically and that with a mass smaller than the critical mass which is in an equilibrium state. In the density range of  $10^4 \text{cm}^{-3} \lesssim n \lesssim 10^{10} \text{cm}^{-3}$ , the interstellar gas is essentially isothermal. In this region a major cooling agent is dusts; that is, the dust is heated by the collision of molecules. The excess energy liberated at the collision increases the dust temperature. Finally the thermal emission from the dust cools down the dust again. By this process, the thermal energy of the gas is reduced. Therefore, we consider the cloud/cloud core is isothermal and study the collapse of the isothermal cloud.

Since the supercritical cloud has no hydrostatic configuration, it must evolve in a dynamical way. On the other hand, since the subcritical cloud is in a static state, it evolves in much longer time-scale of the free-fall time. Such cloud evolves by the effect called the ambipolar diffusion.

## 4.4 Ambipolar Diffusion

### 4.4.1 Ionization Rate

In the dense clouds, the ionization fraction is low. Since the uv/optical radiations from stars can not reach the cloud center, potential ionization comes from the cosmic ray particles. In this case the rate of ionization per volume is given as  $\zeta n_n$ , where  $\zeta \sim 10^{-17} \text{s}^{-1}$ . In Figure 4.3, the ionization fraction for various density is shown (Nakano, Nishi, & Umebayashi 2002). This clearly shows that the fraction of ions decrease approximately in proportion to  $n_H^{-1/2}$  for  $n_H \lesssim 10^8 \text{cm}^{-3}$ . This is understood as follows: Equilibrium balance between one kind of ion  $m^+$  and its neutral  $m^0$

$$m^0 \leftrightarrow m^+ + e^-, \quad (4.35)$$

is considered. The recombination (reaction from the right to left) rate per unit volume is expressed  $\alpha n_e n_{m^+}$ , while the ionization rate (left to right) per unit volume is  $\zeta n_{m^0}$ , where  $\alpha$  means the recombination rate coefficient. Nakano (1984) obtained in the range of  $10^2 \text{cm}^{-3} \lesssim n \lesssim 10^8 \text{cm}^{-3}$

$$\rho_i = C \rho_n^{1/2}, \quad (4.36)$$

where the numerical factor  $C = 4.46 \times 10^{-16} \text{g}^{1/2} \text{cm}^{-3/2}$ .




Figure 4.3: Ionization fractions  $X/n_{\text{H}}$  are plotted against the number density  $n_{\text{H}}$ . The ionization rate of an  $\text{H}_2$  molecule by cosmic rays outside the cloud is taken  $\zeta_0 = 1 \times 10^{-17} \text{s}^{-1}$ . 20% of C and O and 2% of metallic elements are assumed to remain in the gas phase and the rest in grains.

#### 4.4.2 Ambipolar Diffusion

The ionization fraction of the gas in the cloud is low. Neutral gas and the ions are coupled via mutual collisions and ions are coupled with the magnetic field. Thus, the neutral molecules, a major component of the gas, are coupled with the magnetic field indirectly via ionized ions. Equation of motion for the molecular gas is

$$\rho \frac{d\mathbf{u}_n}{dt} = -c_s^2 \nabla \rho - \rho \nabla \psi + \rho r \omega^2 \mathbf{e}_r + \mathbf{F}, \quad (4.37)$$

where the forces appeared in the rhs represent, respectively, the pressure force, the self-gravity, the centrifugal force, and the force exerted on the neutral component per unit volume through two-body collision with ions. The friction force has a form

$$\mathbf{F} = \rho_i \rho_n \gamma (\mathbf{v}_i - \mathbf{v}_n). \quad (4.38)$$

We have assumed  $\rho \simeq \rho_n$  since the mass density of the charged component is much smaller than that of the neutral one. On the other hand, that for ions is

$$\rho_i \frac{d\mathbf{u}}{dt} = \frac{1}{4\pi} (\nabla \times \mathbf{B}) \times \mathbf{B} - \mathbf{F}, \quad (4.39)$$

where we ignored the pressure and self-gravity forces compared with the Lorentz force. If the inertia of the ions are ignored [lhs of eq.(4.39)=0], equations (4.37) and (4.39) give the equation of motion similar to that of the one-fluid as follows:

$$\rho \frac{d\mathbf{u}}{dt} = -c_s^2 \nabla \rho - \rho \nabla \psi + \rho r \omega^2 \mathbf{e}_r + \frac{1}{4\pi} (\nabla \times \mathbf{B}) \times \mathbf{B}. \quad (4.40)$$




Figure 4.4: Relation central density  $\rho_c$  against mass-to-flux ratio  $M/\Phi$ . Plane-parallel and axisymmetric models are illustrated in the left and right panels, respectively.

Ignoring the inertia of the ions, equation (4.39) indicates that the friction force should be balanced with the Lorentz force as

$$\mathbf{F} = \frac{1}{4\pi}(\nabla \times \mathbf{B}) \times \mathbf{B}. \quad (4.41)$$

Using equations (4.36) and (4.38), this gives the drift velocity of ions against the neutrals as

$$\begin{aligned} \mathbf{v}_D &\equiv \mathbf{v}_i - \mathbf{v}_n, \\ &= \frac{\mathbf{F}}{\rho_i \rho_n \gamma}, \\ &= \frac{1}{4\pi \rho_i \rho_n \gamma} (\nabla \times \mathbf{B}) \times \mathbf{B}, \\ &= \frac{1}{4\pi C \gamma \rho_n^{3/2}} (\nabla \times \mathbf{B}) \times \mathbf{B}. \end{aligned} \quad (4.42)$$

Since the Lorentz force works outwardly, the drift velocity  $\mathbf{v}_D$  also points outwardly. Viewing from the ions and magnetic field lines,  $\mathbf{v}_n - \mathbf{v}_i$  points inwardly. Thus, the mass in a flux tube varies with time; that of the cloud center increases while that of the cloud surface decreases. The ambipolar diffusion is a process to change the mass distribution against the magnetic flux tube.

How is the structure affected by the increase of mass/flux ratio at the cloud center? Here, we will derive the relation between the mass-to-flux ratio and central density and answer the question. Numerical calculation is necessary for the axisymmetric (realistic) cloud. However, this can be obtained for the case of a plane-parallel disk in which the magnetic field is running parallel to the disk. For the plane-parallel disk the hydrostatic balance is achieved as

$$\frac{\partial}{\partial z} \left( p + \frac{B^2}{8\pi} \right) = -\rho \frac{\partial \psi}{\partial z} = -2\pi G \rho \sigma, \quad (4.43)$$

where  $\sigma = \int_{-z}^z \rho dz$ . Equation (4.43) is integrated to obtain the relation between the variables at the

center  $z = 0$  as

$$p_c + \frac{B_c^2}{8\pi} - \left( p_s + \frac{B_s^2}{8\pi} \right) = \frac{\pi G}{2} \sigma_s^2, \quad (4.44)$$

where we used  $d\sigma/dz = 2\rho$ . This is rewritten as

$$\frac{\rho_c}{\rho_s} + \frac{1}{8\pi} \left( \frac{B_c}{\rho_c} \right)^2 \frac{\rho_s}{c_s^2} \left( \frac{\rho_c}{\rho_s} \right)^2 - \frac{1}{8\pi} \left( \frac{B_s}{\rho_s} \right)^2 \frac{\rho_s}{c_s^2} = \frac{\pi G \sigma_s^2}{2\rho_s c_s^2} + 1. \quad (4.45)$$

In the second term of lhs,  $y_c \equiv (\rho_c/B_c)(c_s/\rho_s^{1/2})$  is a nondimensional mass flux ratio at the center and this is related to the center-to-surface density ratio  $\rho'_c \equiv \rho_c/\rho_s$ . Using these nondimensional variables, equation (4.45) is expressed as

$$\rho'_c + \frac{(\rho'_c)^2}{8\pi y_c^2} - \frac{1}{8\pi y_s^2} = \bar{\rho}'_c, \quad (4.46)$$

where  $\bar{\rho}'_c \equiv \pi G \sigma_s^2 / (2\rho_s c_s^2) + 1$  represents the central density necessary for the disk to be supported without a magnetic field ( $y \rightarrow \infty$ ). Figure 4.4 (left) illustrates this relation. This shows that if the central mass-to-flux ratio increases (moving upward), the central density increases. In low density,  $\rho'_c \ll \bar{\rho}'_c$ , equation (4.46) indicates that  $\rho'_c \propto y_c \propto \rho_c/B_c$ . This means that on the low density regime, the density increases but the strength of magnetic field does not increase. Equation (4.46) also indicates that  $y_c$  increase much as  $y_c \rightarrow \bar{\rho}'_c y_s$  when  $\rho'_c \rightarrow \bar{\rho}'_c$ . This relation between mass-to-flux ratio and the central density is well fitted to the numerical calculation of plane-parallel disk cloud driven by the ambipolar diffusion by Mouschovias, Paleologou, & Fiedler (1985). Does this indicate that the mass-to-flux ratio increases much when the central density increases, for example, from  $10^4 \text{cm}^{-3}$  to  $10^6 \text{cm}^{-3}$ ?

From this point, quasistatic evolution of the magnetized cloud was first studied by Nakano (1979) adopting a method seeking for magnetohydrostatic configuration (Mouschovias 1976a, 1976b). He obtained that the magnetic flux to mass ratio near the center of the cloud does not decrease much below a critical density above which magnetohydrostatic configurations no more exist. Paleologou & Mouschovias (1983), Shu (1983), and Mouschovias, Paleologou, & Fiedler (1985) gave a completely different result. That is, the magnetic flux to mass ratio near the center decreases much when the density becomes larger than  $n \gtrsim 10^6 \text{cm}^{-3}$ . Critical discussion has been done between them<sup>1</sup>. However, we have now realized that the difference comes from the geometry assumed.

Relation of the mass-to-flux ratio to the central density for the axisymmetric realistic (not plane-parallel disk) cloud is plotted in Figure 4.4 (right) (Tomisaka, Ikeuchi, & Nakamura 1990). The central mass-to-flux ratio increases by the ambipolar diffusion. This increase the central density. Lines with arrows indicate the evolutionary paths driven by the ambipolar diffusion. It is to be noticed that the relation is completely different from that of the plane-parallel disk. Increase of the mass-flux ratio is small although the central density increases much. This indicates that in the quasistatic evolution driven by the ambipolar diffusion the mass-flux ratio does not increase much in contrast to the plane-parallel case.

In fully ionized plasma, the magnetic field is coupled to the matter. However, in the low ionized gas the magnetic field is coupled only to the ions. Thus the induction equation of magnetic field should be written

$$\frac{\partial \mathbf{B}}{\partial t} + \nabla \times (\mathbf{v}_i \times \mathbf{B}) = 0. \quad (4.47)$$

<sup>1</sup> One of the reasons why large increase in the mass/flux ratio is favored is understood as follows: There have been a long-standing ‘‘magnetic flux problem of stars’’ in which the magnetic flux of, say,  $1M_\odot$  main-sequence star  $\Phi \sim XXG \text{cm}^2$  is much smaller than that of the parent cloud  $\Phi \sim XXG \text{cm}^2$ . The magnetic flux must be reduced in the star formation process. If the mass-flux ratio increases much in these density range, this might resolve the magnetic flux problem of stars.



This leads to the final expression as

$$\frac{\partial \mathbf{B}}{\partial t} + \nabla \times (\mathbf{v}_n \times \mathbf{B}) + \nabla \times \left\{ \frac{1}{4\pi C \gamma \rho_n^{3/2}} [(\nabla \times \mathbf{B}) \times \mathbf{B}] \times \mathbf{B} \right\} = 0, \quad (4.48)$$

where we used the expression of the drift velocity [equation (4.42)]. In the isothermal cloud, there is a maximum mass supported by the magnetic field. Since in the subcritical cloud  $M < M_{cr}$  the magnetic flux escapes from the center, the critical mass decreases with time. After the critical mass becomes smaller than the actual mass ( $M_{cl} > M_{cr}$ ), there is no hydrostatic configuration. The cloud evolves into the supercritical cloud region and experiences dynamical collapse.

## 4.5 Dynamical Collapse




Fig. 1. Radial profiles of (a) density and (b) velocity for the  $\alpha = 4$  isothermal model (model A2). The solid and dashed curves denote the physical quantities before and after core formation, respectively. The number attached to each curve denotes the evolution time from the initial state. In this model, we set  $\rho_{\text{sink}} = 10^6 \rho_{\text{eq}}$ .

Figure 4.5: Evolution of isothermal clouds massive than the Bonnor-Ebert mass. Density (*left*) and velocity (*right*) distributions are illustrated. Solid lines show the cores of the preprotostellar phase (prestellar core) and dashed lines show those of protostellar phase (protostellar core). The evolution of the protostellar phase is studied by the sink-cell method, where we assume the gas that entered in the sink-cells is removed from finite-difference grids and add to the point mass sitting at the center of the sink-cells which corresponds to a protostar.

In 1969, Larson (1969) and Penston (1969) found a self-similar solution which is suited for the dynamical contraction. Figure 4.5 is a radial density distribution for a spherical collapse of an isothermal cloud, where the cloud has a four-times larger mass than that of the Bonnor-Ebert mass. Although the figure is taken from a recent numerical study by Ogino et al (1999), a similar solution was obtained in Larson (1969). We can see that the solution has several characteristic points as follows:

1. The cloud evolves in a self-similar way. That is, the spatial distribution of the density (left) at  $t = 0.326$  is well fitted by that at  $t = 0.293$  after shifting in the  $-x$  and the  $+y$  directions. As for the infall velocity spatial distribution, only a shift in the  $-x$  direction is needed.

2. The density distribution in the envelope, which is fitted by  $\propto r^{-2}$ , is almost unchanged. Only the central part of the cloud (high-density region) contracts.
3. The time before the core formation epoch (the core formation time  $t_0$  is defined as the time at which the central density increases greatly) is a good indicator to know how high the central density is. That is, reading from the figure, at  $t - t_0 = 0.003$  ( $t = 0.326$ ) the central density reaches  $\rho_c \simeq 10^4$  and at  $t - t_0 = 0.036$  ( $t = 0.293$ ) the density is equal to  $\rho_c \simeq 10^2$ . This shows the maximum (central) density is approximately proportional to  $(t - t_0)^2$ , which is reasonable from the description of the free-fall time  $\propto \rho^{-1/2}$ .

The basic equations of spherical symmetric isothermal flow are

$$\frac{\partial \rho}{\partial t} + \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \rho v) = 0, \quad (4.49)$$

$$\frac{\partial v}{\partial t} + v \frac{\partial v}{\partial r} + \frac{c_{\text{is}}^2}{\rho} \frac{\partial \rho}{\partial r} + \frac{GM}{r^2} = 0, \quad (4.50)$$

$$M(r, t) = M(0, t) + \int_0^r 4\pi r'^2 \rho(r', t) dr', \quad (4.51)$$

where  $M(r, t)$  represents the mass included in the radius  $r$  and  $M(0, t)$  denotes the mass of the protostar. A self-similar solution which has a form

$$\rho(r, t) = \frac{\Omega(\xi)}{4\pi G(t - t_0)^2}, \quad (4.52)$$

$$v(r, t) = c_{\text{is}} V(\xi), \quad (4.53)$$

$$M(r, t) = \frac{c_{\text{is}}^3 |t - t_0|}{G} m(\xi) \quad (4.54)$$

$$\xi = \frac{r}{c_{\text{is}} |t - t_0|}, \quad (4.55)$$

should be found, where  $\Omega$  and  $V$  are functions only on  $\xi$ . For example, equation (4.52) asks the shape of the density distribution is the same after resizing of equation (4.55)  $r \rightarrow r/c_{\text{is}}|t - t_0|$  and re-normalizing  $\rho \rightarrow \rho \cdot 4\pi G(t - t_0)^2$  as  $\Omega(\xi)$ . Since

$$\frac{\partial}{\partial t} = \frac{d}{d\xi} \left( \frac{\partial \xi}{\partial t} \right)_r = \frac{r}{c_{\text{is}} |t - t_0|^2} \frac{d}{d\xi}, \quad (4.56)$$

and

$$\frac{\partial}{\partial r} = \frac{d}{d\xi} \left( \frac{\partial \xi}{\partial r} \right)_t = \frac{1}{c_{\text{is}} |t - t_0|} \frac{d}{d\xi}, \quad (4.57)$$

the basic equations for the spherical symmetric model yield

$$m = (\xi - V)\xi^2 \Omega, \quad (4.58)$$

$$\left[ (\xi - V)^2 - 1 \right] \frac{dV}{d\xi} = \left[ \Omega(\xi - V) - \frac{2}{\xi} \right] (\xi - V), \quad (4.59)$$

$$\frac{(\xi - V)^2 - 1}{\Omega} \frac{d\Omega}{d\xi} = \left[ \Omega - \frac{2}{\xi}(\xi - V) \right] (\xi - V), \quad (4.60)$$

Equations (4.59) and (4.60) have a singular point at which  $(\xi - V)^2 - 1 = 0$  or  $V = \xi \pm 1$ . Since the point of  $\xi = \text{const}$  moves with  $c_{\text{is}}$ , the flow velocity relative to this  $\xi = \text{const}$  is equal to  $v - c_{\text{is}} \xi = (V - \xi)c_{\text{is}}$ .

Thus the singular point  $\xi_*$  at which  $V = \xi \pm 1$  corresponds to a sonic point. Therefore, since the flow has to pass the sonic point smoothly, the rhs of equations (4.59) and (4.60) have to be equal to zero at the singular point  $\xi_*$ . This gives at the sonic point  $\pm\xi_* > 0$ ,

$$\Omega - \frac{2(\xi - V)}{\xi} = \Omega(\xi - V) - \frac{2}{\xi} = 0, \quad (4.61)$$

which leads to

$$V_* = \xi_* \mp 1, \quad (4.62)$$

$$\Omega_* = \pm \frac{2}{\xi_*}. \quad (4.63)$$

These equations (4.58), (4.59) and (4.60) have an analytic solution

$$V(\xi) = 0, \quad \Omega = \frac{2}{\xi^2}, \quad m = 2\xi, \quad -\infty < \xi < \infty \quad (4.64)$$

This is a solution which agrees with the Chandrasekhar's SIS. Generally, solutions are obtained only by numerical integration.  $|\xi \rightarrow \infty|$  the solution have to converge to an asymptotic form of

$$V(\xi) = V_\infty - \frac{A-2}{\xi} + \frac{V_\infty}{\xi^2} + \frac{4V_\infty + (A-2)(A-6)}{6\xi^3} + \mathcal{O}(\xi^{-4}), \quad (4.65)$$

$$\Omega(\xi) = \frac{A}{\xi^2} - \frac{\Omega_\infty(A-2)}{2\xi^4} + \mathcal{O}(\xi^{-6}), \quad (4.66)$$

This shows that for sufficiently large radius the gas flows with a constant inflow velocity  $V_\infty c_{\text{is}}$ .

This has a solution in which the density and the infall velocity should be regular with reaching the center ( $\xi \ll 1$ ). Such kind of solution is plotted in Figure 4.6 (left) and Figure 4.6 (right) with  $t < 0$ . This time evolution is expected from the self-similar solution. This shows that

$$\rho \begin{cases} \simeq \rho_c & (\text{in the central region}) \\ \propto r^{-2} & (\text{in the outer envelope}) \end{cases} \quad (4.67)$$

$$v \begin{cases} \propto r & (\text{in the central region}) \\ \simeq 3.28c_{\text{is}} & (\text{in the outer envelope}) \end{cases} \quad (4.68)$$

Reaching the outer boundary the numerical solution (Fig.4.5) differs from the self-similar solution (Fig.4.6). For example,  $v$  is reduced to zero in the numerical simulations, while it reaches a finite value 3.28 in the self-similar solution. And as for the density distribution,  $\rho$  drops near the outer boundary in the numerical simulations while it decreases proportional to  $\propto r^{-2}$ . However, in the region except for the vicinity of the outer boundary the self-similar solution expresses well the dynamical collapse of the spherical isothermal cloud. This solution gives the evolution of a pre-protostellar core formed in a supercritical cloud/cloud core.

### Inside-out Collapse Solution

In 1977 Shu found another self-similar solution which is realized after a central protostar with infinitesimal mass is formed in the singular isothermal sphere solution. The gas begins to accrete to the protostar. Outside the region where the accretion occurs, the initial SIS is kept unchanged, since the SIS is a hydrostatic solution. And the front of accretion expands radially outward in time. Since the inflow region expand outwardly, he called it **the inside-out collapse solution**. In Figure 4.7, the




Figure 4.6: A self-similar solution indicating a dynamical collapse of isothermal spherical cloud (Larson-Penston solution). Spatial distribution of the density and inflow velocity which are expected from the self-similar solution are plotted. Dashed lines show the evolution prestellar core and solid lines show that of protostellar core. Taken from Hanawa (1999).

evolution is shown for density and inflow velocity. This solution gives the evolution of a protostellar core formed in a subcritical cloud/cloud core.

Using equations (4.65) and (4.66) and assuming the inflow velocity should reduce at large radius, we obtain

$$V(\xi) = -\frac{A-2}{\xi} - \frac{(A-2)(A-6)}{6\xi^3} + \dots, \quad (4.69)$$

$$\Omega(\xi) = \frac{A}{\xi^2} - \frac{A(A-2)}{2\xi^4} + \dots. \quad (4.70)$$

Since  $\xi \rightarrow \infty$  means  $t \rightarrow t_0$  (if  $r$  is finite),  $\Omega(\xi) \rightarrow A/\xi^2$  means that

$$\rho(r, t_0) = \frac{Ac_{\text{is}}^2}{4\pi Gr^2}. \quad (4.71)$$

Comparing with the SIS, when  $A = 2$  this gives the SIS and when  $A > 2$  this gives a density distribution in which the pressure is inefficient and the cloud is contracting. The solution with  $A > 2$  is obtained by a procedure as (1) at a sufficiently large radius  $\xi_1$ , calculate  $V(\xi_1)$  and  $\Omega(\xi_1)$ . (2) from these values, integrate equations (4.59) and (4.60) inwardly. Figure 4.7 show the solution of this type. The solution with  $A > 2$  inflow speed is accelerated towards the center. Decreasing  $A$  ( $A \rightarrow 2$ ), it is shown that an outer part  $\xi \gtrsim 1$  reaches  $V \rightarrow 0$ . For  $A = 2^+$ , the solution reaches the singular line  $V = -\xi + 1$  at  $\xi = 1$  ( $V = 0$ )<sup>2</sup>. Since  $V = 0$  and  $\Omega = 2$  at  $\xi = 1$ , this solution with  $A = 2^+$  converges to the SIS at  $\xi = 1$ . This means that if there is an infinitesimally small amount of excess mass at the center of SIS, the accretion begins from the center while outside a radius the cloud is left static. The inner part of the solution  $\xi \lesssim 1$ ,  $V$  and  $\Omega$  are well expressed as  $V \propto \xi^{1/2}$  and  $\Omega \propto \xi^{3/2}$ .




Figure 4.7: Self-similar solution which shows the inside-out collapse. (*Left:*) Infall velocity ( $-V = -v/c_{\text{is}}$ ) is plotted against the similarity variable  $\xi \equiv r/c_{\text{is}}(t - t_0)$ . Three curves correspond to models  $A = 2.4, 2.1,$  and  $2.01$ . (*Right:*) Density ( $\Omega = 4\pi G\rho(t - t_0)^2$ ) is plotted against the similarity variable  $\xi \equiv r/c_{\text{is}}(t - t_0)$ .

<sup>2</sup> This means  $A = 2 + \epsilon$  and  $\epsilon > 0$  and  $\epsilon \rightarrow 0$

The power-law distributions of  $V \propto \xi^{1/2}$  and  $\Omega \propto \xi^{3/2}$  are explained as follows: Conservation of the total energy of an inflowing gas shell is expressed as

$$\frac{v^2}{2} - \frac{GM_r}{r} = -\frac{GM_{r_0}}{r_0}, \quad (4.72)$$

where  $r_0$  denotes the initial radius of a gas shell and  $M_{r_0} = M_r$  represents the mass inside the gas shell. Neglecting  $GM_{r_0}/r_0$  compared with the term  $GM_r/r$ , this gives

$$v \simeq \left( \frac{2GM_r}{r} \right)^{1/2} \propto r^{-1/2}, \quad (4.73)$$

where we assumed a major part of  $M_r$  comes from the mass of a protostar  $M_*$ , that is,  $M_r = M_* + \int_0^r \rho 4\pi r^2 dr \simeq M_*$ . Since the average density inside the radius  $r$   $\bar{\rho}(< r) = \int_0^r \rho 4\pi r^2 dr / \int_0^r 4\pi r^2 dr = 3\rho(r)$  for SIS distribution, the time necessary for a gas shell to reach the center is proportional to  $\propto \bar{\rho}(< r)^{-1/2} \propto \rho(r)^{-1/2} \propto r^1 \propto (t - t_0)^1$ , where we used a fact that the front of accretion expands with a constant speed  $c_{s\text{is}}$ . This means that the time necessary for the gas shell to travel from the radius of the accretion wave front to the center is proportional to  $t - t_0$ . Since the mass of the shell which begins accretion in a unit time is equal to  $\rho(r)4\pi r^2 c_{s\text{is}}$  and is constant irrespective of  $t - t_0$ . These two facts indicate that the mass accretion rate is constant in time. That is,

$$\dot{M} = 4\pi r^2 \rho v = \text{const.} \quad (4.74)$$

Since the mass swept by the sound wave per unit time is equal to  $\dot{M}_{sw} = 4\pi r^2 \rho_{\text{SIS}} c_s = 2c_s^3/G$ , the accretion rate is proportional to  $c_s^3/G$ . Shu (1977) obtained the accretion rate

$$\dot{M} = 0.975 \frac{c_s^3}{G}, \quad (4.75)$$

for SIS  $A = 2^+$ . Using this equation, equations (4.73) and (4.74) indicate that the spatial density distribution is expressed by a power-law as

$$\rho(r) \propto r^{-3/2}, \quad (4.76)$$

which is valid for the region except for the vicinity of the front of the accretion.

#### 4.5.1 Protostellar Evolution of Supercritical Clouds

What is a protostellar core formed in a supercritical cloud/cloud core? Is this different from the inside-out solution of Shu (1977)? A solution which corresponds to the protostellar core is obtained by Hunter (1977) and Whitworth and Summers (1985). This is a solution with  $t > t_0$  in equation (4.55). The asymptotic behaviors of the density and infall velocity reaching the center are different from that of the Larson-Penston self-similar solution for a prestellar collapse. That is,

$$\Omega \begin{cases} \rightarrow \text{finite} & (\text{LP}) \\ \rightarrow \text{infinite} & (\text{Inside} - \text{out}) \end{cases} \quad (4.77)$$

$$V \begin{cases} \rightarrow \text{finite} & (\text{LP}) \\ \rightarrow \text{infinite} & (\text{Inside} - \text{out}) \end{cases} \quad (4.78)$$

Using the boundary conditions suitable for the inside-out type solution, another self-similar solution is obtained. In Figure 4.6, such kind of solution is also plotted for  $t > 0$ .

Take notice that the solutions of  $t < t_0$  (prestellar) and  $t > t_0$  (protostellar) agree with each other at  $t = t_0$ . Even if the boundary conditions at the center for the **similarity variables**,  $V$  and  $\Omega$ , are completely different, the difference between the two is small in the physical variable  $v$  and  $\rho$ . Therefore, the evolution of a supercritical core is thought to be expressed by the Larson-Penston self-similar solution extended to the protostellar core phase by Hunter (1977) and Whitworth and Summers (1985).

Assume that we observe a protostellar core and obtain their density and infall velocity spatial distributions. Can we distinguish which solution is appropriate for the Shu's inside-out solution or the extended Larson-Penston solution? This seems hard, because the structure of density and velocity distributions are similar after the protostar is formed: the density and velocity show almost similar power-law as  $\rho \propto r^{-3/2}$  and  $v \propto r^{-1/2}$  irrespective of the inside-out solution or the extended Larson-Penston solution. The region where the infall velocity is accelerated toward the center (accretion-dominated region) is expanding after the protostar is formed. Therefore, to distinguish between the two solutions becomes harder and harder after the protostar is formed. The difference would be large and we would have a definite answer which solution is appropriate to describe the cloud collapse, if we can observe a very young protostellar core or a preprotostellar core which shows dynamical collapse. However, since the time-scale of such a phase is much shorter than the evolved protostellar phase or a younger preprotostellar core, the number of such kind of objects would be small ( $\tau_{\text{ff}} \propto \rho^{-1/2}$ ). Therefore, we are looking for such objects just before or after the protostar formation.

## 4.6 Accretion Rate

Using equation (2.26), the necessary time for a mass-shell at  $R$  to reach the center (free-fall time) is expressed as

$$T(R) \equiv \left( \frac{R^3}{2GM(R)} \right)^{1/2} \frac{\pi}{2} \quad (4.79)$$

(for detail of this section see Ogino, Tomisaka, & Nakamura 1999).

Consider two shells whose initial radii are  $R$  and  $R + \Delta R$ . The time difference for these two shells to reach the center  $\Delta T(R)$  can be written down using equation (4.79) as

$$\Delta T(R) = \frac{\pi R^{1/2}}{2^{3/2}(GM(R))^{1/2}} \left[ \frac{3}{2} - \frac{R}{2M(R)} \frac{dM(R)}{dR} \right] \Delta R. \quad (4.80)$$

Mass in the shell between  $R$  and  $R + \Delta R$ ,  $\Delta M \equiv M(R + \Delta R) - M(R) = (dM/dR)\Delta R$ , accretes on the central object in  $\Delta T(R)$ . Thus, mass accretion rate for a pressure-free cloud is expressed as  $\Delta M/\Delta T$ . This leads to the expression as

$$\frac{dM}{dT}(R) = \frac{2^{3/2} G^{1/2} M(R)^{3/2}}{\pi R^{3/2}} \frac{\frac{R}{M(R)} \frac{dM(R)}{dR}}{\frac{3}{2} - \frac{R}{2M(R)} \frac{dM(R)}{dR}}. \quad (4.81)$$

This gives time variation of the accretion rate. Consider two clouds with the same density distribution  $\partial \log \rho / \partial r$  but different absolute value. Since these two clouds have the same  $\partial \log M(R) / \partial \log R$ , the mass accretion rate depends only on  $M(R)/R$ , and is expressed as

$$\frac{dM}{dT}(R) \propto M(R)^{3/2}. \quad (4.82)$$




Fig. 8. Maximum of mass-accretion rate as a function of  $\alpha$  for the models with various  $\gamma$ . The unit of the mass-accretion rate is taken to be  $c_s^3/G$ , which is equal to  $1.6 \times 10^{-6} M_\odot \text{ yr}^{-1}$  for  $c_s = 0.19 \text{ km s}^{-1}$  ( $T = 10 \text{ K}$ ). The unit of time is taken to be  $1/\sqrt{G\rho_{\text{eq}}}$ , which is equal to  $6.3 \times 10^5 \text{ yr}$  for  $\rho_{\text{eq}} = 10^4 \text{ cm}^{-3}$ . The number attached with each line denotes the value of  $\gamma$ . For each line, we calculated many models with different  $\alpha$ , i.e.,  $\alpha = 1.1, 1.5, 2, 2.5, \dots, 10$ .

Figure 4.8: Mass accretion rate against the typical density of the cloud.

This indicates that the accretion rate is proportional to  $\rho^{3/2}$ , while the time scale is to  $\rho^{-1/2}$ . This is confirmed by hydrodynamical simulations of spherical symmetric isothermal clouds (Ogino et al.1999). When the initial density distribution is the SIS as  $\rho \propto r^{-2}$ , the mass included inside  $R$  is proportional to radius  $M(R) \propto R$ . In this case, equation (4.81) gives a constant accretion rate in time. In Figure 4.8 we plot the mass accretion rate against the cloud density.  $\alpha$  represents the cloud density relative to that of a hydrostatic Bonnor-Ebert sphere. This shows clearly that the mass accretion rate is proportional to  $\alpha^{3/2}$  for massive clouds  $\alpha > 4$ . This is natural since the assumption of pressure-less is valid only for a massive cloud in which the gravity force is predominant against the pressure force.

Similar discussion has been done by Henriksen, André, & Bontemps (1997) to explain a decline in the accretion rate from Class 0 to Class I IR objects. They assumed initial density distribution of

$$\rho \begin{cases} = \rho_0 & (r \leq r_N), \\ = \rho_0 \left(\frac{r}{r_N}\right)^{-2/D_1} & (r > r_N), \end{cases} \quad (4.83)$$

as shown in Figure 4.9. Since the free-fall-time of the gas contained in the inner core  $r \leq r_N$  is the same, such gas reaches the center once. It makes a very large accretion rate at  $t = (3\pi/32G\rho_0)^{1/2}$  as  $\dot{M} = (4\pi/3)\rho_0\delta(t - (3\pi/32G\rho_0)^{1/2})$ . If  $D_1 = 1$ ,  $\rho \propto r^{-2}$  for  $r \gtrsim r_N$ . Since  $M \propto R^1$  and  $t_{\text{ff}} \propto R$ , equation(4.81) predicts  $\dot{M} \propto R^0 \propto t^0$ . A constant accretion rate is expected for this power-law and the accretion rate is converged to a constant value after  $M(r_N) \ll M(r)$ . If  $D_1 = 2$ ,  $\rho \propto r^{-3}$  for  $r \gtrsim r_N$ . Since for this power  $M \propto R^2$  and  $t_{\text{ff}} \propto R^{1/2}$ , equation(4.81) predicts  $\dot{M} \propto R^{3/2} \propto t^3$ . They gave  $\dot{m} \propto t^{3D_1-3}$  for  $D_1 > 2/3$ .

### Problem

Show that  $\dot{M} \propto t^{3D_1-4/3D_1}$  if  $\rho \propto r^{-2/D_1}$  and  $D_1 < 2/3$  [Henriksen, André, & Bontemps (1997)].






Figure 4.9: A model proposed to explain time variation in accretion rate by Henriksen, André, & Bontemps (1997). The density distribution  $\rho(r)$  at  $t = 0$  (left) and expected accretion rate (right).

## 4.7 Outflow

In chapter 1, we mentioned ‘outflows’ ejected from the protostars and pre-main-sequence stars. L1551 IRS5 is a typical example which shows a number of outflows are ejected in the course of star formation. One is molecular outflow, whose lobes extends  $L \sim 0.5(D/160\text{pc})\text{pc}$  in two opposite directions from the IRS-5 (Snell et al 1980). This is traced by CO  $J = 1 \rightarrow 0$  emission line. They estimated an expansion velocity of the CO gas of  $V_{\text{CO}} \sim 15\text{km s}^{-1}$ , a mass of  $0.3 M_{\odot}$ . The dynamical age is equal to  $\tau_{\text{dyn}} = L/V_{\text{CO}} \sim 3 \times 10^4\text{yr}$ . Inside the CO outflow lobe, several Herbig-Haro objects are found, which are emission-line nebulosities. Proper motion studies have revealed the motion of such H-H objects. Cudworth & Herbig (1979) reported HH28 and 29 have  $0.15 - 0.20\text{arcsec yr}^{-1}$ , which corresponds to  $150 - 200\text{km s}^{-1}$ . In a neighbor of the IR source a stellar jet is found (Mundt & Fried 1983) by optical emission lines. Its size is equal to  $17\text{arcsec} = 4 \times 10^{16}\text{cm}(D/160\text{pc})$  and the jet indicates rather well-collimated shape (the opening angle  $\simeq 10^\circ$ ). The relationship between these two outflows (massive molecular outflow and less massive optical jet) are not clear yet.

### 4.7.1 Magneto-driven Model

To accelerate gas till supersonic speed, the magnetic force is considered to be an important player. Blandford & Payne (1982) have proposed the ‘‘magneto-centrifugal wind model’’ for the acceleration in two-dimensional axisymmetric configuration.

Figure 4.10 explains the magneto-centrifugal wind mechanism (Fig. 1 of Blandford & Payne 1982). Consider a disk under the gravity of a central star with mass  $M_*$ . Assume that gas in the disk at the radius  $r_0$  rotates with a Kepler speed of

$$v_{\phi}(r_0) = \left( \frac{GM_*}{r_0} \right)^{1/2}. \quad (4.84)$$

Consider a gas parcel whose original position was  $(r_0, z = 0)$ . The effective potential for the particle




Figure 4.10: Magneto-centrifugal wind model. Figure shows the effective potential for a gas parcel with conserving the angular rotation speed  $\omega_0$  of the disk (eq.4.85). The gas parcel is assumed to be located originally at  $(r_0, 0)$ . The  $x$ - and  $y$ -axes mean normalized distance from the central star as  $r/r_0$  and  $z/r_0$ . The darker color represents the deeper potential.

rotating with an angular rotation speed of  $\omega_0 = v_\phi(r_0)/r_0$  is written as

$$\phi(r, z) = -\frac{GM_*}{(r^2 + z^2)^{1/2}} - \frac{1}{2}r^2\omega_0^2 = -\frac{GM_*}{r_0} \left[ \frac{r_0}{(r^2 + z^2)^{1/2}} + \frac{1}{2} \left( \frac{r}{r_0} \right)^2 \right], \quad (4.85)$$

where  $\omega_0 = (GM)^{1/2}/r_0^{3/2}$ . (Be careful that ordinary effective potential is calculated for constant angular momentum. However, this is for constant angular rotation speed.) The gas will co-rotate with the same angular rotation speed as the disk, if the magnetic field is sufficiently strong. Equation (4.85) corresponds to the effective potential for the gas co-rotating with the disk at  $r_0$ . Figure 4.10 plots the isopotential contour lines for the effective potential [eq.(4.85)]. The contour line passing through the point  $(r_0, 0)$  represents  $\Phi = -3GM/2r_0$ . The angle between the isopotential line of  $\Phi = -3GM/2r_0$  and the  $r$ -axis is equal to  $\pm 60^\circ$  (see problem below). Further, it can be obtained that the effective potential increases upwardly while it decreases with reaching the central star or radially outwardly.

This leads to an important conclusion that

1. If the angle between the magnetic field line and the  $r$ -axis  $\theta_{\text{mag}}$  is larger than  $60^\circ$  ( $\theta_{\text{mag}} > 60^\circ$ ), gas parcel moving along the field line must climb the effective potential. It needs extra energy to depart from the disk.
2. If  $\theta_{\text{mag}} < 60^\circ$ , gas parcel moving along the field line slides down the effective potential. It can depart from the disk without extra energy.

Be careful that this is valid if the magnetic field is sufficiently strong. If the Kepler disk is threaded with the magnetic field lines with a shallow angle, and if the angular momentum is sufficiently transferred along the magnetic field line, in this case, the gas is accelerated by the extra centrifugal force and finally escapes from the potential well of the central star. This mechanism is called the “magneto-centrifugal wind” mechanism.




Figure 4.11: Pole-on (left) and side (right) views of the magnetic field lines, which drive the outflow.

### Problem

Show that the angle between the isopotential line which passes the point  $(r_0, 0)$  and the  $r$ -axis is equal to  $\pm 60^\circ$ . Calculate the Taylor expansion of the effective potential near the point  $(r_0, 0)$  and show that to satisfy  $\phi(r_0 + \Delta r, \Delta z) = \phi(r_0, 0)$ ,  $(\Delta z / \Delta r)^2$  must be equal to 3.

### Angular Momentum Transfer

In the axisymmetric case, the poloidal  $(z, r)$  and toroidal  $(\phi)$  components of the magnetic field  $\mathbf{B}$  and current  $\mathbf{j}$  are decoupled with each other. That is, the poloidal ( $B_p$ ) and toroidal ( $B_\phi$ ) magnetic fields are made by the toroidal ( $j_\phi$ ) and poloidal ( $j_p$ ) electric currents, respectively. As for the Lorentz force  $\mathbf{j} \times \mathbf{B}$ , the poloidal component comes from  $j_\phi B_p$  or  $j_p B_\phi$ , while the toroidal component does from  $\mathbf{j}_p \times \mathbf{B}_p$ . Even if there is no toroidal magnetic field (thus no poloidal electric current), there exists the poloidal component of the Lorentz force, which acts as a pressure to counter-balance the self-gravity (§4.2). On the other hand, the toroidal component of the Lorentz force appears only the case with the poloidal electric current and thus toroidal component of magnetic field  $B_\phi$ . This means that the angular momentum is transferred by the magnetic field only when  $B_\phi$  exists. Equation (B.14) explains how the angular momentum density  $\rho v_\phi r$  is transferred. The left-hand side of equation (B.14) represents the advection of the angular momentum density, while the right-hand side

$$\frac{r}{4\pi} \left[ \frac{1}{r} \frac{\partial}{\partial r} (r B_\phi) B_r + \frac{\partial B_\phi}{\partial z} B_z \right] = \left( \frac{\mathbf{B}_p}{4\pi} \right) \cdot \nabla (r B_\phi), \quad (4.86)$$

represents the torque exerted on the gas parcel.

The induction equation of the magnetic field [eq.(B.17)] shows that  $B_\phi$  is generated from poloidal magnetic field by the effect of rotational motion  $v_\phi$ . This indicates that the angular momentum is transferred as follows:

1. The toroidal component of magnetic field  $B_\phi$  is amplified by the dynamo process [equation (B.17)]. The magnetic field lines run like spiral viewing from the top like Figure 4.11.
2. Negative torque works near region attached to the fast-rotating disk, while positive one appears in the region connected to the radial magnetic field. The angular momentum is transferred from inside to outside along the same field line.

3. Angular momentum received by the gas near disk surface leads the excess centrifugal force which accelerate the gas. The disk inside which is affected by the negative torque loses its angular momentum and insufficient centrifugal force drives further accretion.

This angular momentum transfer on the same magnetic field has a character of *angular momentum redistribution*.

### Centrifugal Radius

In a diffuse cloud, the centrifugal force does not play an important role. Consider a gas parcel whose specific angular momentum is equal to  $j$ . When the gas element contracts till the distance  $r$  from the center, the centrifugal force per mass of

$$\frac{v_\phi^2}{r} = \frac{j^2}{r^3}, \quad (4.87)$$

works. Contraction stops when the gravitational acceleration  $GM/r^2$  is balanced by the centrifugal force. Its radius (centrifugal radius) is expressed as

$$r_c = \frac{j^2}{GM}. \quad (4.88)$$

Assuming that the disk is near the hydrostatic balance in  $z$ -direction, total column density is related to the volume density on the  $z = 0$  plane as

$$\sigma = \left( \frac{2c_s^2 \rho_c}{\pi G} \right)^{1/2}. \quad (4.89)$$

We can show that a nondimensional ratio of specific angular momentum to the mass is equal to the ratio of the free-fall time to the rotation period as follow:

$$\frac{c_s j}{GM} = \frac{c_s \omega_c r^2}{G \sigma \pi r^2} = \frac{\omega_c}{(2\pi G \rho_c)^{1/2}}. \quad (4.90)$$

Many numerical simulations confirm that the contracting disk in the runaway isothermal contraction phase has some ‘universality.’ The ratio of the free-fall time to the rotation period is approximately equal to

$$\frac{\omega_c}{(2\pi G \rho_c)^{1/2}} \simeq 0.3, \quad (4.91)$$

regardless of the initial conditions (Matsumoto, Nakamura, & Hanawa 1997). Thus,  $c_s j / GM \simeq 0.3$ . Finally, we can see the centrifugal radius as

$$r_c = \frac{GM}{c_s^2} \left( \frac{c_s j}{GM} \right)^2 \simeq 0.3 \frac{GM}{c_s^2}, \quad (4.92)$$

increases with time in proportional to the mass  $M$ , because gas element with large  $j$  contracts later. As the centrifugal radius increases with time, the launching point of the outflow also expands with time.




Figure 4.12: (Left) Relation between density and temperature at the center of the cloud. (Right) Cooling and heating rates, which are calculated for the cloud center are plotted against the density.

#### 4.7.2 Entrainment Model

Models that a protostar is driving a jet and the jet transfers its momentum to the ambient material and forms a molecular outflow are called as entrainment models. From hydrodynamical calculations, De Young (1986) pointed out there are two mechanisms of entrainment. The ejected jet forms a bow shock and matter hit by the bow shock is accelerated promptly (prompt entrainment). His numerical simulations predicted that gas within the radius from the jet axis of  $(1 - 3) \times$  jet radius is entrained in this mechanism. After the bow shock has passed away, slower mechanism of entrainment works, in which a turbulent layer is developed around the jet and the Reynolds stress in the layer transfers the momentum (Stahler 1994). In this mechanism, the ratio of entrained ambient gas to the injected mass by the jet is estimated as  $(1/3 - 3)$  (De Young 1986).

### 4.8 Evolution to Star

In the proceeding sections we have seen the evolution from gas cloud to stars. However we were restricted to the isothermal gasses. To understand the change in the temperature, we have to consider the radiation which keeps the gas isothermal. Spherical symmetric radiative hydrodynamical simulations have been studied to understand the evolutionary path from interstellar cloud to star. Figure 4.12 (left) illustrates the relation between  $\rho$  and  $T$  obtained by Masunaga & Inutsuka (2000). Several characteristic power-laws are seen in the figure. Figure 4.13 plots the density and temperature distribution.

#### Isothermal Phase

Below  $\rho \lesssim 10^{-13} \text{g cm}^{-3}$ , gas is essentially isothermal  $T \propto \rho^0$ . This corresponds to points number 1-3 of Figure 4.13. Since internal energy of gas is transferred to the thermal energy of dusts by collisions, the main coolant in this regime is the dust thermal radiation. The cooling rate per mass is

$$\Lambda = 4\kappa_{\text{p}}\sigma T^4, \quad (4.93)$$

where  $\kappa_{\text{p}}(10\text{K}) \sim 0.01 \text{cm}^2 \text{g}^{-1}$  and  $\sigma$  represent Planck mean absorption coefficient and the Stephan-Boltzman constant. Main heating process for  $\rho \lesssim 10^{-14} \text{g cm}^{-3}$  is cosmic-ray heating:  $\epsilon_{\text{CR}} \sim 2 \times$




Figure 4.13: Dynamical evolution of a spherical cloud studied with RHD simulation. Density (left) and temperature (right) distributions are plotted against the radius.

$10^{-4} \text{erg s}^{-1} \text{g}^{-1}$  (Goldsmith & Langer 1978). Balance between these two majors asks that temperature is constant as

$$T \sim 3\text{K} \left( \frac{\epsilon_{\text{CR}}}{2 \times 10^{-4}} \right)^{1/4} \left( \frac{\kappa_p}{0.01 \text{cm}^2 \text{g}^{-1}} \right)^{1/2}. \quad (4.94)$$

Heating rate due to dynamical compression,

$$\Gamma = -p \frac{d(1/\rho)}{dt} \simeq c_s^2 (4\pi G \rho), \quad (4.95)$$

increases according to the contraction and it balances with the above cooling at the density  $\rho_A \sim 10^{-14} \text{g cm}^{-3}$  (Masunaga, Miyama, & Inutsuka 1998).

The cloud in this phase experiences the dynamical contraction as described in section 4.5. Structure of  $\rho(r)$  and  $v_r(r)$  is well represented by the Larson-Penston self-similar solution.

### First Core

After  $\rho \gtrsim \rho_A$ , gas is no more isothermal and becomes adiabatic. Between  $10^{-13} \text{g cm}^{-3} \lesssim \rho \lesssim 10^{-9} \text{g cm}^{-3}$ , gas obeys the  $\gamma = 5/3$  polytropes. Above the density  $\rho \gtrsim \rho_A$ , the optical depth for the thermal radiation exceeds unity  $\tau \gtrsim 1$  and radiative cooling can not compensate the compressional heating. As long as the temperature is low as  $T \lesssim 100\text{K}$ , neither rotation nor vibration is excited for  $\text{H}_2$  molecule. Even  $\text{H}_2$  gas behaves like single-atom molecule. Thus  $\gamma \simeq 5/3$ .

Between  $10^{-9} \text{g cm}^{-3} \lesssim \rho \lesssim 10^{-7.5} \text{g cm}^{-3}$ , the exponent becomes  $\gamma = 7/5$ , which characterizes that the gas consists of two-atom molecule  $\text{H}_2$ .

In this phase, relatively large gas pressure supports against the gravity and the cloud becomes hydrostatic (points number 4-6 of Figure 4.13). This is called as ‘‘first core’’ made by the molecular hydrogen. The density structure of the first core is well represented by a polytrope sphere with the specific heat ratio of  $\gamma = 7/5$  or the polytropic index  $n = 2\frac{1}{2}$ . From equation (C.11) in Appendix C.1, such a polytrope has a mass-density relation as

$$M_{c1} \propto \rho_c^{1/10}, \quad (4.96)$$

where  $M_{c1}$  and  $\rho_c$  represent, respectively, the mass of the first core and the central density. At the beginning, the core mass is equal to  $M_{c1} \simeq 0.01 M_\odot$ . As long as the mass increases a factor 3, the central density increases 5 orders of magnitude.

### Second Collapse

For  $10^{-7.5} \text{g cm}^{-3} \lesssim \rho \lesssim 10^{-2.5} \text{g cm}^{-3}$ , the exponent becomes  $\gamma = 1.1$ . In this density range of  $\rho \gtrsim 10^{-7.5} \text{g cm}^{-3}$  the temperature exceeds  $10^3\text{K}$ . Due to this temperature, the hydrogen molecules begin to dissociate and the dissociation process absorbs the energy. By this, gas becomes much softer ( $\gamma \rightarrow 1$ ) than the first core. This introduces another dynamical contraction, the second collapse (points number 6-7 of Figure 4.13).

### Second Core

After the dissociation is completed, the exponent becomes  $\gamma = 5/3$ . This gas forms a second core composed of atomic hydrogen, which is called the second core (points number 7-8 of Figure 4.13).




Figure 4.14: Explanation of nested grid method.

Since the accretion rate [eq.(4.75)] is proportional to  $\mathcal{C}_s^3$  or  $T^{3/2}$ , the accretion rates onto the second core  $\dot{M}_{2\text{nd}}$  is larger than that of the first core  $\dot{M}_{1\text{st}}$  as

$$\frac{\dot{M}_{2\text{nd}}}{\dot{M}_{1\text{st}}} \simeq 5 \times 10^3 \frac{(T/10^{3.5}\text{K})^{3/2}}{(T/10\text{K})^{3/2}}. \quad (4.97)$$

Thus, the first core disappears quickly and after that the gas begins to accrete onto the second core, which will be a protostar. However, this is the case of non-rotating, spherical symmetric cloud collapse. As previously seen, the angular momentum plays a crucial role and forms a disk. The evolution must be different completely for such a case.

## 4.9 Example of Numerical Simulation

We have described the cloud run-away collapse and succeeding accretion process. In the latter phase, there is a possibility that the outflow is driven by the effect of magnetic Lorentz force. A restricted number of numerical simulation can simulate such evolution throughout. The size of the molecular cores  $\sim 0.1\text{pc}$  is more than  $3 \times 10^4$  times as large as the typical size of the first core  $\sim 1\text{AU}$ . Therefore numerical scheme to calculate such process must have a large dynamic-range. In the finite-difference scheme, the spatial dynamic range is restricted by the number of the cells. Although at least  $3 \times 10^5 \times 3 \times 10^5$  grid points (in 2D) seems necessary to resolve a factor of  $3 \times 10^4$ , to solve the MHD equation **with the Poisson equation** of the self-gravity have not been done. This is done by the Eulerian nested-grid simulation. This method uses a finite number of grid systems with different spatial resolutions are prepared. Coarse grid covers whole the cloud and is to see a global structure. Fine grid covers only the central part of the cloud and is to see the fine-scale structure appeared near the center. Figure 4.14 explains grid cells of the nested grid method. The size of the cells of the  $n$ -th level grid ( $L_n$ ) is taken equal to a half of those of the  $L_{n-1}$ .

A cylindrical cloud with coaxial magnetic field in hydrostatic balance is studied. Rotation vector  $\boldsymbol{\omega}$  and the magnetic field  $\mathbf{B}$  are parallel to the symmetric axis of the cylinder ( $z$ -axis). For the






Figure 4.15:  $\Omega_c = 10^{-13}\text{rad s}^{-1}$ ,  $B_c = 10\mu\text{G}$ ,  $\rho_c = 10^4\text{H}_2\text{ cm}^{-3}$ , and  $c_s = 190\text{m s}^{-1}(T/10\text{K})$ . Snapshots at the time of  $0.6066\tau_{\text{ff}} = 1.061\text{Myr}$  represented in different levels are shown: L1 (upper-left), L5 (upper-right), and L10 (bottom). The actual size of the frames of L5 (upper-right) and L10 (bottom) are, respectively, 1/16 and 1/512 smaller than that of L1 (upper-left). Magnetic field lines (dotted lines) and isodensity contours (solid lines) are presented as well as the velocity vectors by arrows.

energy equation, a double-polytropic equation of state is adopted, since the radiation hydrodynamical calculation (for example Masunaga & Inutsuka 2000) predicts  $\rho - T$  relations like shown in Figure 4.12, which consists of a number of power-laws for some density ranges.

$$p = \begin{cases} c_s^2 \rho & (\text{for } \rho < \rho_A), \\ c_s^2 \rho_A \left(\frac{\rho}{\rho_A}\right)^{7/5} & (\text{for } \rho > \rho_A), \end{cases} \quad (4.98)$$

where we take  $\rho_A = 10^{10} \text{H}_2 \text{ cm}^{-3}$ . Figure 4.15 is the model of  $\Omega_c = 10^{-13} \text{rad s}^{-1}$ ,  $B_c = 10 \mu\text{G}$ ,  $\rho_c = 10^4 \text{H}_2 \text{ cm}^{-3}$ ,  $c_s = 190 \text{m s}^{-1} (T/10\text{K})$ , and scale-length  $H = 0.341 \text{pc} (c_s/190 \text{m s}^{-1}) (\rho_c/10^4 \text{H}_2 \text{ cm}^{-3})^{-1/2}$ . This is a snapshot of  $0.6066 \tau_{\text{ff}} = 1.061 \text{Myr}$ . The cylindrical cloud breaks into pieces with a length in  $z$ -direction equal to the most unstable wavelength of the gravitational instability (Matsumoto, Nakamura, & Hanawa 1994) as

$$\lambda_{\text{max}} \simeq \frac{2\pi(1 + \alpha/2 + \beta)^{1/2}}{0.72 \left[ (1 + \alpha/2 + \beta)^{1/3} - 0.6 \right]^{1/2}} \frac{c_s}{(4\pi G \rho_c)^{1/2}}. \quad (4.99)$$

The magnetic fields prevent gas motions perpendicular to the field lines  $\mathbf{B}$ . Further, the centrifugal force works in the perpendicular direction to  $\boldsymbol{\omega}$ . A disk forms which running in the perpendicular direction to  $\mathbf{B}$  and  $\boldsymbol{\omega}$  and thus to the  $z$ -axis. Figure 4.15 illustrates a snapshot at the time of  $0.6066 \tau_{\text{ff}} = 1.061 \text{Myr}$ . Level 1 grid (upper-left) captures the global structure of the contracting disk perpendicular to the  $z$ -axis. Close up view is shown in L5, which has 16 times finer spatial resolution (upper right). Since speed,  $v_z$ , of infalling gas is larger than the sound speed, a number of shock fronts are formed near  $z \simeq 0.02H$  and  $z \lesssim 0.01H$ . Further inner structure is represented in L10 grid (bottom) which has 32 times finer resolution than L5. Another discontinuity in density is now forming near  $z \sim 0.0002H$ . Since gas inside of this discontinuity has the density larger than the critical density  $\rho > \rho_A$ , in this phase the first core (p.85) begins to be formed. In the isothermal collapse phase, a characteristic power-law for the Larson-Penston self-similar solution is seen in the radial distributions of density as  $\rho(r, z = 0) \propto r^{-2}$ , and  $B_z(r, z = 0) \propto r^{-1}$ .

After  $t > 0.6067 \tau_{\text{ff}}$  the first core is formed. Radial infall motion is accelerated toward the surface of the first core and the rotational motion  $v_\phi$  also increases. This amplification promotes the toroidal magnetic field,  $B_\phi$ . Since the torque is proportional to  $(\nabla \times B_\phi \mathbf{e}_\phi) \times \mathbf{B}_p$ , the amplitude of the torque increases after the first core formation.

Structure just after  $\sim 500$  yr has passed from Figure 4.15 is shown in Figure 4.15 In  $\tau \sim 500$  yr from the epoch of Figure 4.15, flow is completely changed and outflow is launched. Panel (b), which corresponds to a four-times close-up of panel (a), shows clearly that outflow is ejected from a region near  $r \simeq 1.5 \times 10^{-4}H$  and  $z \simeq 0.5 \times 10^{-4}H$ . And the outflow vectors and the disk has an angle of approximately 45 deg. The magneto-centrifugal wind model favors a small angle between the magnetic field and the disk,  $\theta_{\text{mag}} < 60 \text{deg}$ . This model predicts that the angle between the flow and the disk is also smaller than 60 deg. Numerical results is not inconsistent with this prediction of magneto-centrifugal wind model.

To see which force is driving such outflow, amplitudes of three forces are compared: the Lorentz force  $\mathbf{F}_m = (\nabla \times \mathbf{B}) \times \mathbf{B} / 4\pi$ , the centrifugal force  $\mathbf{F}_c = (\rho v_\phi^2 / r) \mathbf{e}_r$ , and the thermal pressure gradient  $\mathbf{F}_t = -\nabla p$ . The parallel components to the poloidal magnetic field of the three forces are as

$$\begin{aligned} \mathbf{F}_m \cdot \frac{\mathbf{B}_p}{|\mathbf{B}_p|} &= \frac{\nabla \times \mathbf{B} \times \mathbf{B}}{4\pi} \cdot \frac{\mathbf{B}_p}{|\mathbf{B}_p|}, \\ &= -\frac{1}{8\pi r^2} \frac{B_p}{|\mathbf{B}_p|} \cdot \nabla (r B_\phi)^2 \end{aligned} \quad (4.100)$$

(Ustyugova et al. 1999)

$$\mathbf{F}_c \cdot \frac{\mathbf{B}_p}{|\mathbf{B}_p|} = \rho v_\phi^2 r \cdot \frac{B_r}{|\mathbf{B}_p|}, \quad (4.101)$$

and

$$\mathbf{F}_p \cdot \frac{\mathbf{B}_p}{|\mathbf{B}_p|} = -\nabla p \cdot \frac{\mathbf{B}_p}{|\mathbf{B}_p|}. \quad (4.102)$$

Figure 4.16 illustrates the largest force at each grid point. Centrifugal force dominated region (region C), which is shown by asterisk (\*), overlaps the outflow region around point P1 ( $z = 10^{-4}H$ ,  $2.5 \times 10^{-4}H$ ). Just radially exterior to this region C, there is another region near point P2 ( $z = 2 \times 10^{-4}H$ ,  $3.5 \times 10^{-4}H$ ) filled with puls signs where the magnetic force is dominated (region M). Comparing with the flow vectors in the left, the strong outflow coincides with this region M and the above region C. Therefore, the outflow is driven by the centrifugal force and the Lorentz force (pressure gradient by toroidal magnetic fields).

Difference between models with different magnetic field strength but the same rotation speed is illustrated in Figure 4.17. Magnetic to thermal pressure ratio  $\alpha$  is taken 1 [panels (a), and (d)], 0.1 [panels (b), and (e)], and 0.01 [panels (c), and (f)]. Panels (a), (b), (c) in the left column are snapshots when the central density reaches  $\rho_A$  while panels (d), (e), (f) in the right column show the structure after  $\tau = 4.5 \times 10^{-3} \tau_{\text{ff}} = 8000\text{yr}$  passed. In the model with  $\alpha = 1$ , the outflow gas flows through a region whose shape resembles a capital letter U. This is similar to that observed in model ( $\alpha = 1$  and  $\Omega = 5$ ) shown in Figure 4.16. Decreasing the magnetic field strength  $\alpha = 1 \rightarrow 0.1$  and  $\rightarrow 0.01$ , the shape of disk changes from a flat disk to a round ellipsoid. The model with neither magnetic field nor rotation shows a spherical contracting core. Outflow is also affected by decrease in magnetic field strength. Globally folded magnetic field lines are seen in the outflow region in panel (e). Smaller-scale structure folding the magnetic field lines develops in the outflow region in panel (f). The structure seen in the poloidal magnetic field is well correlated to the distribution of toroidal magnetic field strength. That is, a packet of strong  $B_\phi$  pinches the gas and poloidal magnetic field lines radially inwardly. This means that the toroidal magnetic field is amplified by the rotation motion and becomes predominant over the poloidal one in the case with weak initial magnetic field. Thus, the hoop stress, which is expressed by the first term of the Lorentz force of equation (B.13) and comes from the tension of toroidal magnetic field, pinches the plasma contained inside of the loop of  $B_\phi$ . For this case, the magnetic force dominant region is widely spread compared with the centrifugal force dominant region, which is far from the model with strong magnetic field. In this case, the magnetic pressure gradient plays an important role. Gas of outflow is ejected perpendicularly from the disk and forms flow similar to the capital letter I. In conclusion, U-type and I-type outflows are generated for strong magnetic model and weak magnetic model, respectively.




Figure 4.16: Continuation of the evolution shown in Fig.4.15. Snapshots at the time of  $0.6069 \tau_{\text{ff}} = 1.066 \text{ Myr}$  just  $\sim 500 \text{ yr}$  after from Fig.4.15. Panel (a) plots the structure of L10 which is similar to Fig.4.15 (c). Spherical region inside  $\sim 1.2 \times 10^{-3} H$  has been swept by the outflow. Panel (b) is a snapshot of L12 grid which has 4-times finer spatial resolution than L10 (a). In panel (c), we plotted magnetic force dominated region with '+', centrifugal force dominated with '\*', and thermal pressure force dominated with blank ' '.




Figure 4.17: Difference in models with the same rotation speed  $\Omega = 2 \times 10^{-14} \text{ rad s}^{-1}$  but different magnetic field strength ((a) and (d):  $B_c = 10\mu\text{G}$ , (b) and (e):  $3\mu\text{G}$ , and (c) and (f):  $1\mu\text{G}$ ).



# References

- [1] Alves, J., Lada, C.J., & Lada, E.A., 2001, *Internal Structure of a Cold Dark Molecular Cloud Inferred from Extinction of background Starlight*, Nature, 409, 159
- [2] Alves, J., Lada, C.J., Lada, E.A., Kenyon, S.J., & Phelps, R. 1998, *Dust Extinction and Molecular Cloud Structure: L977*, ApJ, 506, 292
- [3] André, P. 1994, *Observations of Protostars and Protostellar Stages* in “The Cold Universe,” eds. T. Montmerle, C.J.Lada, I.F.Mirabel, & J.Tran Thanh Van, Editions Frontières, p.179
- [4] Binney, J., & Tremaine, S. 1988, *Galactic Dynamics*, Princeton Univ. Pr. Chapter 6.
- [5] Bonnor W. B., 1956, *Boyle’s Law and Gravitational Instability*, MNRAS, 116, 351
- [6] Blandford, R. D., & Payne, D. G., 1982, *Hydromagnetic flows from Accretion Discs and the Production of Radio Jets*, MNRAS, 199, 883
- [7] Chandrasekhar, S. 1939, *An Introduction to the Study of Stellar Structure*, (Univ. of Chicago Pr.; Chicago)
- [8] Cudworth, K. M., & Herbig, G. 1979, *In two large-proper-motion Herbig-Haro objects*, AJ, 84, 548
- [9] De Young, D. S. 1986, *Mass entrainment in astrophysical jets*, ApJ, 307, 62
- [10] Ebert R., 1955, *er die Verdichtung von H I-Gebieten. Mit 5 Textabbildungen*, Z. Astrophys., 37, 217
- [11] Elmegreen, B.G.; Seiden, P.E. & Elmegreen, D.M. 1989, *Spiral Arm Amplitude Variations and Pattern Speeds in the Grand Design Galaxies M51, M81, and M100*, ApJ, 343, 602
- [12] Fiege, J.D., & Pudritz, R.E. 2001, *Helical fields and filamentary molecular clouds - I*, MNRAS, 311, 85
- [13] Fridlund, C.V.M., & Liseau, R. 1998, *it Two Jets from the Protostellar System L1551 IRS5*, ApJL, 499, 75
- [14] Goldsmith, P. F., & Langer, W. D., 1978, *Molecular cooling and thermal balance of dense interstellar clouds*, ApJ, 222, 881
- [15] Gordon, M.A., & Burton, W.B. 1976, *Carbon monoxide in the Galaxy. I - The Radial Distribution of CO, H2, and Nucleons*, ApJ, 208, 346
- [16] Hanawa, T. 1999, , *Star Formation*, in “Active Universe” in Japanese (Shokabo, Tokyo) ed. by K. Shibata, J. Fukue, R. Matsumoto, & S. Mineshige
- [17] Hartmann, L. 1998, *Accretion Processes in Star Formation*, (Cambridge U.Pr.).
- [18] Heithausen, A., Bensch, F., Stutzki, J., Falgarone, E., & Panis, J.F., 1998, *The IRAM key project: small-scale structure of pre-star forming regions Combined mass spectra and scaling laws*, AAp, 331, L65
- [19] Henriksen, R., Andre, P., & Bontemps, S. 1997, *Time-dependent accretion and ejection implied by pre-stellar density profiles*, AAp, 323, 549
- [20] Hirano, N., Kameya, O., Nakayama, M., & Takakubo, K. 1988, *Bipolar outflow in B335*, ApJL, 327, L69

- [21] Honma, M., Sofue, Y., & Arimoto, N. 1995, *The Molecular Front in Galaxies. II. Galactic-scale Gas Phase Transition of HI and H<sub>2</sub>*, AAp, 304, 1
- [22] Itoh, Y. et al., 2000, *A Pair of Twisted Jets of Ionized Iron from L 1551 IRS 5* PASJ, 52, 81
- [23] Kennicutt, R.C. 1998, *The Global Schmidt Law in Star-forming Galaxies*, ApJ, 498, 541
- [24] Kennicutt, R.C., Tamblyn, P., & Congdon, C.W. 1994, *Past and Future Star Formation in Disk Galaxies*, ApJ, 435, 22
- [25] Kramer, C., Stutzki, J., Rohrig, R., & Corneliussen, U. 1998, *Clump Mass Spectra of Molecular Clouds*, AAp, 329, 249
- [26] Lada, C.J., 1999, *The Formation of Low Mass Stars*, in “The Origin of Stars and Planetary Systems” ed. C.J. Lada & N.D. Kylafis, (Kluwer) p.143
- [27] Loren, R. B., 1989, *The cobwebs of Ophiuchus. I - Strands of <sup>13</sup>CO - The mass distribution* , ApJ, 338, 902
- [28] Masunaga, H. & Inutsuka, S., 2000, *A Radiation Hydrodynamic Model for Protostellar Collapse. II. The Second Collapse and the Birth of a Protostar*, ApJ, 531, 350
- [29] Masunaga, H., Miyama, S. M., & Inutsuka, S., 1998, *A Radiation Hydrodynamic Model for Protostellar Collapse. I. The First Collapse*, ApJ, 495, 346
- [30] Matsumoto, T., Hanawa, T., & Nakamura, F., 1997, *Gravitational Contraction of Rotating Clouds: Formation of Self-similarly Collapsing Disks*, ApJ, 478, 569
- [31] Matsumoto, T., Nakamura, F., Hanawa, T. 1994 *Fragmentation of a magnetized filamentary molecular cloud rotating around its axis*, PASJ, 46, 243
- [32] Mizuno, A., Onishi, T., Hayashi, M., Ohashi, N., Sunada, K., Hasegawa, T., & Fukui, Y. 1994, *Molecular Cloud Condensation as a Tracer of Low Mass Star Formation Nature*, 368, 719
- [33] Mizuno, A. et al. 1995, *Overall Distribution of Dense Molecular Gas and Star Formation in the Taurus Cloud Complex*, ApJL, 445, L161
- [34] Mouschovias, T. C. 1976, *Nonhomologous contraction and equilibria of self-gravitating, magnetic interstellar clouds embedded in an intercloud medium: Star formation. II - Results*, ApJ, 207, 141
- [35] Mottel, F., & André, P. 2001, *The Circumstellar Environment of Low-mass Protostars: A Millimeter Continuum Mapping Survey*, AAp, 365, 440
- [36] Motte, F., André, P., Ward-Thompson, D., & Bontemps, S. 2001, *A SCUBA survey of the NGC 2068/2071 protoclusters*, AAp, 372, L41
- [37] Mouschovias, T. Ch., 1976a, *Nonhomologous contraction and equilibria of self-gravitating, magnetic interstellar clouds embedded in an intercloud medium: Star formation. I Formulation of the problem and method of solution*, ApJ, 206, 753
- [38] Mouschovias, T. Ch., 1976b, *Nonhomologous contraction and equilibria of self-gravitating, magnetic interstellar clouds embedded in an intercloud medium: Star formation. II - Results*, ApJ, 207, 141
- [39] Mouschovias, T. Ch., Paleologou, E. V., & Fiedler, R. A., 1985, *The magnetic flux problem and ambipolar diffusion during star formation - One-dimensional collapse. II - Results*, ApJ, 291, 772
- [40] Myers, P.C. 1978, *A compilation of Interstellar Gas Properties*, ApJ, 225, 380
- [41] Mundt, R., & Fried, J. W. 1983, *Jets from young stars*, ApJ, 274, 83
- [42] Nakano, T., 1979, *Quasistatic Contraction of Magnetic Protostars due to Magnetic Flux Leakage - Part One - Formulation and an Example*, PASJ, 31, 713
- [43] Nakano, T. Nishi, R., Umebayashi, T., 2002, *Mechanism of Magnetic Flux Loss in Molecular Clouds*, ApJ, 572, 199
- [44] Ogino, S., Tomisaka, K., & Nakamura, F. 1999, *Gravitational Collapse of Spherical Interstellar Clouds* PASJ, 51, 637.



- [45] Ohashi, N., Hayashi, M., Ho, P.T.P., Momose, & M., Hirano, N. 1996, *Possible Infall in the Gas Disk around L1551 IRS 5*, ApJ, 466, 957
- [46] Ohashi, N. Lee, S. W., Wilner, D. J. & Hayashi, M. 1999, *CS Imaging of the Starless Core L1544: an Envelope with Infall and Rotation*, ApJL, 518, L410
- [47] Onishi, T., Mizuno, A., Kawamura, A., Ogawa, H., Fukui, Y. 1996, *A  $C^{18}O$  Survey of Dense Cloud Cores in Taurus: Core Properties*, ApJ, 468, 815
- [48] Onishi, T., Mizuno, A., Kawamura, A., Tachihara, K., & Fukui, Y. 2002, *A Complete Search for Dense Cloud Cores in Taurus*, ApJ, 575, 950
- [49] Paleologou, E. V., & Mouschovias, T. Ch. 1983, *The magnetic flux problem and ambipolar diffusion during star formation - One-dimensional collapse. I - Formulation of the problem and method of solution*, ApJ, 275, 838
- [50] Reipurth, B., Yu, K.C., Rodríguez, L.F., Heathcote, S., & Bally, J. 1999, *Multiplicity of the HH 111 Jet Source: it Hubble Space Telescope NICMOS Images and VLA Maps*, AAp, 352, L83
- [51] Saito, M., Sunada, K., Kawabe, R., Kitamura, Y., & Hirano, N. 1999, *he Initial Conditions for Formation of Low-Mass Stars: Kinematics and Density Structure of the Protostellar Envelope in B335*, ApJ, 518, 334
- [52] Schmidt, M. 1959, *The Rate of Star Formation*, ApJ, 129, 243
- [53] Shu, F. H. 1978, *Self-similar collapse of isothermal spheres and star formation*, ApJ, 214, 488
- [54] Shu, F. H. 1983, *Ambipolar diffusion in self-gravitating isothermal layers*, ApJ, 273, 202
- [55] Shu, F.H., Milione, V., & Roberts, W.W.,Jr. 1973, *Nonlinear Gaseous Density Waves and Galactic Shocks*, ApJ, 183, 819
- [56] Snell, R. L., Loren, R. B., & Plambeck, R. L. 1980, *Observations of CO in L1551 - Evidence for stellar wind driven shocks*, ApJ, 239, 17
- [57] Spitzer, L. Jr. 1978, *Physical Processes in the Interstellar Medium*, Wiley, Cahper 13.
- [58] Stahler, S. W. 1983, *The Birthline for Low-mass Stars*, ApJ, 274, 822
- [59] Stahler, S. W. 1994, *The kinematics of molecular outflows*, ApJ, 422, 616
- [60] Stutzki, J. & Guesten, R., 1990, *High spatial resolution isotopic CO and CS observations of M17 SW - The clumpy structure of the molecular cloud core*, ApJ, 356, 513
- [61] Tafalla, M., Mardones, D., Myers, P. C., Caselli, P., Bachiller, R. & Benson, P. J. 1998, *L1544: A Starless Dense Core with Extended Inward Motions*, ApJ, 504, 900
- [62] Tamura, M., Hough, J.H., Hayashi, S.S., 1995, *Millimeter Polarimetry of Young Stellar Objects: Low-Mass Protostars and T Tauri Stars*, ApJ, 448, 346
- [63] Tomisaka, K. 2002, *Collapse of Rotating Magnetized Molecular Cloud Cores and Mass Outflows*, ApJ, 575, 306
- [64] Tomisaka, K., Ikeuchi, S., & Nakamura, T., 1988, *The equilibria and evolutions of magnetized, rotating, isothermal clouds. I - Basic equations and numerical methods*, ApJ, 326, 208
- [65] Tomisaka, K., Ikeuchi, S., & Nakamura, T., 1988, *Equilibria and evolutions of magnetized, rotating, isothermal clouds. II - The extreme case: Nonrotating cloud*, ApJ, 335, 293
- [66] Tomisaka, K., Ikeuchi, S., & Nakamura, T., *The equilibria and evolutions of magnetized, rotating, isothermal clouds. IV - Quasi-static evolution*, ApJ, 362, 202
- [67] Toomre, A. 1981, *What amplifies the spirals*, in "The structure and evolution of normal galaxies" Cambridge U.Pr., 1981, pp.111-136
- [68] Tothill, N. F. H., White, G. J., Matthews, H. E., McCutcheon, W. H., McCaughrean, M. J., & Kenworthy, M. A. 2002, *The Structure and Evolution of the Lagoon Nebula. I. Submillimeter Continuum and CO Line Mapping*, ApJ, 580, 285
- [69] Ustyugova, G. V., Koldoba, A. V., Romanova, M. M., Chechetkin, V. M., & Lovelace, R. V. E. 1999 *Magnetocentrifugally Driven Winds: Comparison of MHD Simulations with Theory*, ApJ,

516, 221

- [70] Ward-Thompson, D., Kirk, J.M., Crutcher, R.M., Greaves, J.S., Holland, W.S., André, P. 2000, *First Observations of the Magnetic Field Geometry in Prestellar Cores*, ApJL, 537, L135
- [71] Weintraub, D.A., Goodman, A.A., & Akeson, R.L. 1999, *Polarized Light from Star-Forming Regions*, in Protostars and Planets IV, ed. V.Mannings, A.P.Boss, & S.S. Russell, U. Arizona Pr. pp.247-271
- [72] Zhou, S. 1995, *Line formation in Collapsing Cloud Cores with Rotation and Applications to B335 and IRAS 16293-2422*, ApJ, 442, 685
- [73] Zhou, S., Evans, N.J.,II, Koempe, C., & Walmsley, C.M. 1993, *Evidence for Protostellar Collapse in B335*, ApJ, 404, 232

# Appendix A

## Basic Equation of Fluid Dynamics

### A.1 What is fluid?

Gas and liquid change their shape according to the shape of the container. This is a definition of **fluid**. Stress tensor

### A.2 Equation of Motion

Newton's second law of mechanics as

$$m \frac{d\mathbf{v}}{dt} = F, \quad (\text{A.1})$$

where  $m$ ,  $\mathbf{v}$ , and  $F$  represent the mass, the velocity of a particle and force working on the particle. In fluid dynamics, using the mass density  $\rho$  and the force working on the unit volume  $f$  equation (A.1) is rewritten as

$$\rho \frac{d\mathbf{v}}{dt} = \mathbf{f}. \quad (\text{A.2})$$

Which kind of force works in a fluid? Gas pressure force does work in any fluids. Beside this, if there is the gravity,  $\rho\mathbf{g}$  should be included in  $\mathbf{f}$ . If the electric currents is running in the fluid and the magnetic fields exist, the Lorentz force  $\mathbf{j} \times \mathbf{B}$  should be added.

To write down the expression of the gas pressure force, consider a fluid element between  $x$  and  $x + \Delta x$ . Pressure force exerting on the surface  $S$  at  $x$  is  $p(x)S$ , while that on the opposite side is  $-p(x + \Delta x)S$ . The net pressure force working on the volume of  $S\Delta x$  is equal to  $(p(x) - p(x + \Delta x))S$ , which is approximated as  $-\partial p / \partial x \Delta x S + O(\Delta x^2)$ . Thus, the pressure force working on the unit volume is written  $-\partial p / \partial x$ . Equation (A.2) can be rewritten as

$$\rho \frac{d\mathbf{v}}{dt} = -\nabla p + \rho\mathbf{g}. \quad (\text{A.3})$$

### A.3 Lagrangian and Euler Equation

The time derivative appearing in equation (A.1) expresses how the velocity of a **specific particle** changes. Therefore, that appearing in equation (A.3) is also expressing the same meaning, that is, the position of the gas element concerning in equation (A.3) moves and the positions at  $t$  and  $t + \Delta t$  are generally different. However, considering the velocity field in the space, the time derivative of the velocity should be calculated staying at a point  $x$ .

These two time derivative are different each other and should be distinguished. The former time derivative is called **Lagrangian time derivative** and is expressed using  $d/dt$ . On the other hand, the latter is called **Eulerian time derivative** and is expressed using  $\partial/\partial t$ . These two are related with each other. Consider a function  $F$  whose independent variables are time  $t$  and position  $\mathbf{x}$ , that is  $F(\mathbf{x}, t)$ . The difference  $\frac{dF}{dt}\Delta t$ , using the Lagrangian time derivative of  $F$ , represents the the difference of  $F(t + \Delta t)$  from  $F(t)$  focusing on a specific fluid element, whose positions are different owing to its motion. The element at the position of  $\mathbf{x}_0$  at the epoch  $t_0$  moves to  $\mathbf{x}_0 + \mathbf{v}_0\Delta t$  in time span of  $\Delta t$ . Thus the difference is expressed as

$$\begin{aligned} \frac{dF}{dt}\Delta t &= F(\mathbf{x}_0 + \mathbf{v}_0\Delta t, t_0 + \Delta t) - F(\mathbf{x}_0, t_0), \\ &\simeq \left(\frac{\partial F}{\partial \mathbf{x}}\right)_{t|\mathbf{x}_0, t_0} \cdot \mathbf{v}_0\Delta t + \left(\frac{\partial F}{\partial t}\right)_{x|\mathbf{x}_0, t_0} \Delta t, \end{aligned} \quad (\text{A.4})$$

where we used the Taylor expansion of  $F$ . The difference regarding to the Eulerian derivative is written down as

$$\frac{\partial F}{\partial t}\Delta t = F(\mathbf{x}_0, t_0 + \Delta t) - F(\mathbf{x}_0, t_0), \quad (\text{A.5})$$

and this is equal to the second term of the rhs of equation (A.4). Comparing equations (A.4) and (A.5), the Lagrangian derivative contains an extra term besides the Eulerian derivative. That is, the Lagrangian derivative is expressed by the Eulerian derivative as

$$\frac{d}{dt} = \frac{\partial}{\partial t} + \mathbf{v} \cdot \frac{\partial}{\partial \mathbf{x}} = \frac{\partial}{\partial t} + \mathbf{v} \cdot \text{grad}. \quad (\text{A.6})$$

Applying the above expression on equation of motion based on the Lagrangian derivative (A.3), we obtain the Eulerian equation motion:

$$\rho \left( \frac{\partial \mathbf{v}}{\partial t} + \mathbf{v} \cdot \nabla \mathbf{v} \right) = -\nabla p + \rho \mathbf{g}. \quad (\text{A.7})$$

## A.4 Continuity Equation

Another basic equation comes from the mass conservation. This is often called the continuity equation, which relates the change of the volume to its density. Consider a fluid element whose volume is equal to  $\Delta V$ . The mass contained in the volume is constant. Thus

$$\frac{d\rho\Delta V}{dt} = \frac{d\rho}{dt}\Delta V + \frac{d\Delta V}{dt}\rho = 0. \quad (\text{A.8})$$

The variation of the volume  $\frac{d\Delta V}{dt}$  is rewritten as

$$\frac{d\Delta V}{dt} = \int_{\partial\Delta V} \mathbf{v} \cdot d\mathbf{S} = \int_{\Delta V} \text{div}\mathbf{v}dV, \quad (\text{A.9})$$

where  $\partial\Delta V$  represents the surface of the fluid element  $\Delta V$ . From equations (A.8) and (A.9), we obtain the mass continuity equation for Lagrangian time derivative as

$$\frac{d\rho}{dt} + \rho \text{div}\mathbf{v} = 0. \quad (\text{A.10})$$

Using equation (A.6) this is rewritten to Eulerian form as

$$\frac{\partial \rho}{\partial t} + \text{div}(\rho\mathbf{v}) = 0. \quad (\text{A.11})$$

Basic equations using the Lagrangian derivative are equations (A.3) and (A.10), while those of the Euler derivative are equations (A.7) and (A.11).

## A.5 Energy Equation

The above basic equations (A.3) and (A.10) or equations (A.7) and (A.11) contain three dependent variables  $\rho$ ,  $p$ , and  $\mathbf{v}$ . The number of the variables, 3, is larger than the number of equations, 2. Therefore, an extra equation is needed to close the basic equations.

### A.5.1 Polytropic Relation

If the pressure of the fluid,  $p$ , is expressed only by the density,  $\rho$ ,

$$p = P(\rho), \quad (\text{A.12})$$

the number of dependent variables is reduced to two and the above equations are sufficient to describe the dynamics of the fluid. Occasionally, the pressure is assumed proportional to the power of  $\rho$  as

$$p = K\rho^\Gamma, \quad (\text{A.13})$$

where  $\Gamma$  is a constant. This assumption is called **polytropic relation**.

The fact should be reminded that the validity of the assumption comes from the physical condition of the system. In the case that the temperature of the gas is kept constant owing to the cooling and heating process, the gas pressure is proportional to the density

$$p = c_{is}^2 \rho, \quad (\text{A.14})$$

where  $c_{is} = (kT/\mu m_p)^{1/2}$  ( $k$ : Boltzmann constant,  $\mu$  average molecular weight, and  $m_p$  is the proton mass) represents the isothermal sound speed and is constant.

Another example is the isentropic fluid, in which the entropy is kept constant. In this case the pressure is proportional to  $\rho^\gamma$ , as

$$p = c_s^2 \rho^\gamma, \quad (\text{A.15})$$

where  $\gamma$  is the specific heat ratio  $= c_p/c_v$  and  $c_s = (\gamma kT/\mu m_p)^{1/2}$  represents the adiabatic sound speed and is constant. In these cases the polytropic relations of equations (A.14) and (A.15) plays a role as the third equation of basic equations of hydrodynamics.

### A.5.2 Energy Equation from the First Law of Thermodynamics

In more general cases, the last equation comes from the first law of the thermal physics,

$$\Delta Q = dU + pdV. \quad (\text{A.16})$$

where  $U$  and  $V$  represent the internal energy and the volume and  $\Delta Q$  is the heat added or subtracted from the system. Using the total energy per unit volume

$$\epsilon = \frac{1}{2}\rho v^2 + \frac{U}{V} \quad (\text{A.17})$$

the above equation gives the equation for total energy:

$$\frac{\partial \epsilon}{\partial t} + \text{div}(\epsilon + p)\mathbf{v} = \rho \mathbf{v} \cdot \mathbf{g}. \quad (\text{A.18})$$

Since the total energy per unit volume is expressed using the basic physical quantities as

$$\epsilon = \frac{1}{2}\rho v^2 + \frac{1}{\gamma - 1}p, \quad (\text{A.19})$$

equation (A.18) is the final basic equation for hydrodynamics. Equations (A.7), (A.11), and (A.18) are basic equations hydrodynamics using the Eulerian time derivative.

## A.6 Shock Wave

### A.6.1 Rankine-Hugoniot Relation

Passing through a shock front moving with a speed  $V_s$ , the physical variables  $\rho$ ,  $p$ , and  $u$  change abruptly. Since the basic equations of hydrodynamics is unchanged after chosing the system moving  $V_s$ , the continuity equation

$$\frac{\partial \rho}{\partial t} + \frac{\partial \rho u}{\partial x} = 0 \quad (\text{A.20})$$

gives an equation for a steady state as

$$\frac{\partial \rho u}{\partial x} = 0, \quad (\text{A.21})$$

from which we obtain the jump condition,

$$\rho_1 u_1 = \rho_2 u_2, \quad (\text{A.22})$$

after we integrate over a region overlapping the shock front. Here, the quantities with suffix 1 are for preshock and those with suffix 2 are for postshock.

Equation of motion for steady state

$$\rho u \frac{\partial u}{\partial x} = -\frac{\partial p}{\partial x}, \quad (\text{A.23})$$

gives

$$p_1 + \rho_1 u_1^2 = p_2 + \rho_2 u_2^2, \quad (\text{A.24})$$

where we used equation(A.22).

### Isotharmal shock

In the case of the gas is isothermal  $p = c_{is}^2 \rho$ , equation (A.24) becomes

$$\rho_1 (c_{is}^2 + u_1^2) = \rho_2 (c_{is}^2 + u_2^2). \quad (\text{A.25})$$

Eliminating  $\rho$  from equations (A.22) and (A.25), we obtain

$$(u_1 u_2 - c_s^2)(u_1 - u_2) = 0, \quad (\text{A.26})$$

which means

$$u_1 u_2 = c_s^2. \quad (\text{A.27})$$

From equation (A.22),

$$\frac{\rho_2}{\rho_1} = \frac{u_1}{u_2} = \frac{u_1^2}{c_s^2}. \quad (\text{A.28})$$

This indicates the postshock velocity  $u_1 \gg c_s$  the ratio of the postshock density to the preshock density becomes large.

# Appendix B

## Basic Equations of Magnetohydrodynamics

### B.1 Magnetohydrodynamics

Here, we derive the basic equations of magnetohydrodynamics. Differences from hydrodynamics are the Lorentz force in the equation of motion and the induction equation of magnetic field. The Lorentz force is written as

$$\mathbf{j} \times \mathbf{B} = \frac{1}{4\pi}(\nabla \times \mathbf{B}) \times \mathbf{B}, \quad (\text{B.1})$$

where we used the Ampere's law  $\nabla \times \mathbf{B} = (4\pi/c)\mathbf{j}$  and the displacement current is ignored. Thus, the equation of motion becomes as

$$\rho \left( \frac{\partial \mathbf{v}}{\partial t} + (\nabla \cdot \mathbf{v})\mathbf{v} \right) = -\nabla p - \rho \nabla \psi + \frac{1}{4\pi}(\nabla \times \mathbf{B}) \times \mathbf{B}, \quad (\text{B.2})$$

in the Eulerian form.

The Faraday's law  $\nabla \times \mathbf{E} = -(1/c)(\partial \mathbf{B} / \partial t)$  and the Ohm's law  $\mathbf{j} = \sigma(\mathbf{E} + \mathbf{v} \times \mathbf{B}/c)$  ( $\sigma$  is the electric conductivity) lead the induction equation of the magnetic field as

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}) - \frac{c^2}{4\pi} \nabla \times \frac{\nabla \times \mathbf{B}}{\sigma}. \quad (\text{B.3})$$

The factor  $c^2/4\pi\sigma$  is the coefficient of magnetic diffusibility. When we assume the electric conductivity  $\sigma \rightarrow \infty$ , diffusive MHD equation, equation(B.3), reduces to the ideal MHD equation as

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}). \quad (\text{B.4})$$

#### B.1.1 Flux Freezing

An important nature of the ideal MHD is derived from equation (B.4), that is, the matter and the magnetic field is coupled with each other. Consider a gas parcel threaded with the magnetic field  $\mathbf{B}$ . The magnetic flux anchored to the gas parcel  $\Phi_B \equiv \int \mathbf{B} \cdot d\mathbf{S}$  is changes

$$\frac{d\Phi_B}{dt} = \frac{d}{dt} \int \mathbf{B} \cdot d\mathbf{S} = \int \frac{\partial \mathbf{B}}{\partial t} \cdot d\mathbf{S} + \int \mathbf{B} \cdot \mathbf{v} \times d\mathbf{s}, \quad (\text{B.5})$$

where  $\mathbf{s} = \partial \mathbf{S}$ . The first term of the lhs of the equation comes from a time derivative of the magnetic flux density, while the last term represents the change of the integral region due to gas motion (see




Figure B.1: The change of flux anchored to a surface  $dS$ . One part of the change comes from the temporal change in magnetic flux density. The other comes from the change of the boundary of the surface due to gas motion. Time variation of the area is given by  $\mathbf{v}dt \times d\mathbf{s}$ .

Fig.B.1). ( $d\mathbf{S} = \mathbf{v} \times d\mathbf{s}$ ) Using the Stokes' theorem  $\int \nabla \times \mathbf{A} \cdot d\mathbf{S} = \int \mathbf{A} \cdot d\mathbf{s}$ , the last term is reduced to  $\int \mathbf{B} \times \mathbf{v} \cdot d\mathbf{s} = \int \nabla(\mathbf{B} \times \mathbf{v}) \cdot d\mathbf{S}$ . Finally, we obtain

$$\frac{d\Phi_B}{dt} = \int \left[ \frac{\partial \mathbf{B}}{\partial t} - \nabla(\mathbf{v} \times \mathbf{B}) \right] \cdot d\mathbf{S} = 0, \quad (\text{B.6})$$

where we used equation (B.4). This means that the magnetic flux anchored to a gas parcel does not change in the ideal MHD regime.

### B.1.2 Basic Equations of Idel MHD

Basic equations are as follows: The mass continuity as

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0, \quad (\text{B.7})$$

the equation of motion as

$$\rho \left( \frac{\partial \mathbf{v}}{\partial t} + (\nabla \cdot \mathbf{v}) \mathbf{v} \right) = -\nabla p - \rho \nabla \psi + \frac{1}{4\pi} (\nabla \times \mathbf{B}) \times \mathbf{B}, \quad (\text{B.8})$$

the equation of thermal energy as

$$\frac{\partial \epsilon}{\partial t} + \text{div}(\epsilon + p) \mathbf{v} = \rho \mathbf{v} \cdot \mathbf{g}, \quad (\text{B.9})$$

or some barotropic relation  $p = P(\rho)$  and the induction equation as

$$\frac{\partial \mathbf{B}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{B}). \quad (\text{B.10})$$

### B.1.3 Axisymmetric Case

The basic equations to be solved are the magnetohydrodynamical equations and the Poisson equation for the gravitational potential. In cylindrical coordinates  $(z, r, \phi)$  with  $\partial/\partial\phi = 0$ , the equations are expressed as follows:

$$\frac{\partial \rho}{\partial t} + \frac{\partial}{\partial z}(\rho v_z) + \frac{1}{r} \frac{\partial}{\partial r}(r \rho v_r) = 0, \quad (\text{B.11})$$

$$\begin{aligned} \frac{\partial \rho v_z}{\partial t} + \frac{\partial}{\partial z}(\rho v_z v_z) + \frac{1}{r} \frac{\partial}{\partial r}(r \rho v_z v_r) = \\ - c_s^2 \frac{\partial \rho}{\partial z} - \rho \frac{\partial \psi}{\partial z} + \frac{1}{4\pi} \left[ -\frac{\partial B_\phi}{\partial z} B_\phi - \left( \frac{\partial B_r}{\partial z} - \frac{\partial B_z}{\partial r} \right) B_r \right], \end{aligned} \quad (\text{B.12})$$



$$\begin{aligned} \frac{\partial \rho v_r}{\partial t} + \frac{\partial}{\partial z}(\rho v_r v_z) + \frac{1}{r} \frac{\partial}{\partial r}(r \rho v_r v_r) = \\ - c_s^2 \frac{\partial \rho}{\partial r} - \rho \frac{\partial \psi}{\partial r} + \frac{1}{4\pi} \left[ -\frac{1}{r} \frac{\partial}{\partial r}(r B_\phi) B_\phi + \left( \frac{\partial B_r}{\partial z} - \frac{\partial B_z}{\partial r} \right) B_z \right], \end{aligned} \quad (\text{B.13})$$

$$\frac{\partial \rho r v_\phi}{\partial t} + \frac{\partial}{\partial z}(\rho r v_\phi v_z) + \frac{1}{r} \frac{\partial}{\partial r}(r \rho r v_\phi v_r) = \frac{r}{4\pi} \left[ \frac{1}{r} \frac{\partial}{\partial r}(r B_\phi) B_r + \frac{\partial B_\phi}{\partial z} B_z \right] \quad (\text{B.14})$$

$$\frac{\partial B_z}{\partial t} = \frac{1}{r} \frac{\partial}{\partial r}[r(v_z B_r - v_r B_z)], \quad (\text{B.15})$$

$$\frac{\partial B_r}{\partial t} = -\frac{\partial}{\partial z}(v_z B_r - v_r B_z), \quad (\text{B.16})$$

$$\frac{\partial B_\phi}{\partial t} = \frac{\partial}{\partial z}(v_\phi B_z - v_z B_\phi) - \frac{\partial}{\partial r}(v_r B_\phi - v_\phi B_r), \quad (\text{B.17})$$

$$\frac{\partial^2 \psi}{\partial z^2} + \frac{1}{r} \frac{\partial}{\partial r} \left( r \frac{\partial \psi}{\partial r} \right) = 4\pi G \rho, \quad (\text{B.18})$$

where the variables have their ordinary meanings. Equation (B.11) is the continuity equation; equations (B.12), (B.13) and (B.14) are the equations of motion. The induction equations for the poloidal magnetic fields are equations (B.15) and (B.16) and for the toroidal magnetic field is equation (B.17). The last equation (B.18) is the Poisson equation.



# Appendix C

## Hydrostatic Equilibrium

### C.1 Polytrope

If we choose the polytropic equation of state<sup>1</sup>,

$$p = K\rho^{1+1/n}, \quad (\text{C.1})$$

the hydrostatic balance is expressed by

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left( \frac{r^2}{\rho} \frac{dp}{dr} \right) = -4\pi G\rho, \quad (\text{C.2})$$

where we used equations (4.1) and (4.2). A hydrostatic gaseous star composed with a polytropic gas is called polytrope. Normalizing the density, pressure and radius as

$$\rho \equiv \rho_c \theta, \quad (\text{C.3})$$

$$p \equiv K\rho_c^{1+1/n} \theta^{n+1}, \quad (\text{C.4})$$

$$r \equiv H\xi = \left( \frac{(n+1)K\rho_c^{(1-n)/n}}{4\pi G} \right)^{1/2} \xi \quad (\text{C.5})$$

we obtain a normalized equation as

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left( \xi^2 \frac{d\theta}{d\xi} \right) = -\theta^n, \quad (\text{C.6})$$

which is called Lane-Emden equation of index  $n$ . The boundary condition at the center of polytrope should be

$$\theta = 1, \quad (\text{C.7})$$

$$\frac{d\theta}{d\xi} = 0, \quad (\text{C.8})$$

at  $\xi = 0$ . Solution of this equation is plotted for several  $n$  in Figure C.1.

---

<sup>1</sup> Reference book for this appendix is chapter 4 of Chandrasekhar (1939).




Figure C.1: Solutions of eq.(C.6) are plotted for indices of  $n = 3$  or  $\Gamma = 4/3$  (the most extended one),  $n = 5/2$  or  $\Gamma = 7/5$  (the middle), and  $n = 3/2$  or  $\Gamma = 5/3$ (the most compact one). For  $n \geq 5$  gas extends infinitely and the solution has no zero-point.

Mass of the polytrope is written down as

$$\begin{aligned}
 M &= \int_0^r 4\pi\rho r^2 dr, \\
 &= 4\pi H^3 \int_0^\xi \theta^n \xi^2 d\xi, \\
 &= 4\pi H^3 \int_0^\xi \frac{d}{d\xi} \xi^2 \frac{d\theta}{d\xi} d\xi, \\
 &= -4\pi H^3 \xi^2 \frac{d\theta}{d\xi} \Big|_{\xi_1} \\
 &= -4\pi \left[ \frac{(n+1)K}{4\pi G} \right]^{3/2} \rho_c^{(3-n)/2n} \xi^2 \frac{d\theta}{d\xi} \Big|_{\xi_1}, \tag{C.9}
 \end{aligned}$$

where  $\xi_1$  represents the zero point of  $\theta$  or the surface radius normalized by  $H$ . For  $n = 3$  or  $\Gamma \equiv 1 + 1/n = 4/3$ , equation (C.9) reduces to

$$M = -4\pi \left( \frac{K}{\pi G} \right)^{3/2} \xi^2 \frac{d\theta}{d\xi} \Big|_{\xi_1}. \tag{C.10}$$

Thus, the mass does not depend on the central density for  $\Gamma = 4/3$  polytrope. For  $n = 5/2$  or  $\Gamma = 1 + 1/n = 7/5$ ,  $M$  is written down as

$$M = -4\pi \left( \frac{7K}{8\pi G} \right)^{3/2} \rho_c^{1/10} \xi^2 \frac{d\theta}{d\xi} \Big|_{\xi_1}, \tag{C.11}$$

where  $\xi_1 = 5.35528$  and  $-\xi_1^2 d\theta/d\xi|_{\xi_1} = 2.18720$ . Polytrope with  $\Gamma = 7/5$  gas, the mass-density relation becomes  $\rho_c \propto M^{10}$ . While, for  $n = 3/2$  of  $\Gamma = 1 + 1/n = 5/3$ ,  $M$  is written down as

$$M = -4\pi \left( \frac{5K}{8\pi G} \right)^{3/2} \rho_c^{1/2} \xi^2 \frac{d\theta}{d\xi} \Big|_{\xi_1}, \tag{C.12}$$

where  $\xi_1 = 3.65375$  and  $-\xi_1^2 d\theta/d\xi|_{\xi_1} = 2.71406$ .

## C.2 Magneto hydrostatic Configuration

In section 4.2, we obtained the maximum mass which is supported against the self-gravity using the virial analysis. In this section, we will survey result of more realistic calculation.

Formalism was obtained by Mouschovias (1976a,6), which was extended by Tomisaka, Ikeuchi, & Nakamura (1988) to include the effect of rotation. Magneto hydrostatic equilibrium is achieved on a balance between the Lorentz force, gravity, thermal pressure force, and the centrifugal force as

$$\frac{1}{4\pi}(\nabla \times \mathbf{B}) \times \mathbf{B} - \rho \nabla \psi - \nabla p - \rho \boldsymbol{\omega} \times (\boldsymbol{\omega} \times \mathbf{r}) = 0. \quad (\text{C.13})$$

In the axisymmetric case, the poloidal magnetic fields is obtained by the magnetic flux function,  $\Phi$ , or the  $\phi$ -component of the vector potential as

$$\begin{aligned} B_z &= \frac{1}{r} \frac{\partial \Phi}{\partial r}, \\ B_r &= -\frac{1}{r} \frac{\partial \Phi}{\partial z}, \\ B_\phi &= \frac{b_\phi}{r}. \end{aligned} \quad (\text{C.14})$$

Equation (C.13) leads to

$$-\frac{1}{4\pi r^2} \Delta_1 \Phi \frac{\partial \Phi}{\partial z} - \frac{b_\phi}{4\pi r^2} \frac{\partial b_\phi}{\partial z} = \rho \frac{\partial \psi}{\partial z} + \frac{\partial p}{\partial z}, \quad (\text{C.15})$$

$$-\frac{1}{4\pi r^2} \Delta_1 \Phi \frac{\partial \Phi}{\partial r} - \frac{b_\phi}{4\pi r^2} \frac{\partial b_\phi}{\partial r} = \rho \frac{\partial \psi}{\partial r} + \frac{\partial p}{\partial r} - \rho \omega^2 r, \quad (\text{C.16})$$

$$-\frac{\partial b_\phi}{\partial r} \frac{\partial \Phi}{\partial z} + \frac{\partial b_\phi}{\partial z} \frac{\partial \Phi}{\partial r} = 0, \quad (\text{C.17})$$

with

$$\Delta_1 \Phi \equiv \frac{\partial^2 \Phi}{\partial z^2} + r \frac{\partial}{\partial r} \left( \frac{1}{r} \frac{\partial \Phi}{\partial r} \right). \quad (\text{C.18})$$

Equation (C.17) indicates  $b_\phi$  is a function of  $\Phi$  as  $b_\phi(\Phi)$ , which is constant along one magnetic field line. Ferraro's isorotation law demands, that is, to satisfy the steady-state induction equation  $\omega$  is constant along a magnetic field. This means  $\omega$  is also constant along one magnetic field line,  $\omega(\Phi)$ . From this, the density distribution in one flux tube is written

$$\rho = \frac{q}{c_s^2} \exp \left[ - \left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right]. \quad (\text{C.19})$$

This means  $q$  is also constant along one magnetic field line,  $q(\Phi)$ . Since the forces are expressed by the derivative of function  $q$

$$(\nabla q) \exp \left[ - \left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right] = \nabla p + \rho \nabla \psi - \rho r \omega^2 - \rho r^2 \omega \nabla \omega, \quad (\text{C.20})$$

where equation (C.19) is used, equation (C.15) and (C.16) are rewritten as

$$\Delta_1 \Phi \frac{\partial \Phi}{\partial z} + \frac{\partial (b_\phi^2/2)}{\partial z} = -4\pi r^2 \left\{ \frac{\partial q}{\partial z} \exp \left[ - \left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right] + \rho r^2 \omega \frac{\partial \omega}{\partial z} \right\}, \quad (\text{C.21})$$

$$\Delta_1 \Phi \frac{\partial \Phi}{\partial r} + \frac{\partial (b_\phi^2/2)}{\partial r} = -4\pi r^2 \left\{ \frac{\partial q}{\partial r} \exp \left[ - \left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right] + \rho r^2 \omega \frac{\partial \omega}{\partial r} \right\}. \quad (\text{C.22})$$

Finally, using the fact that  $b$ ,  $q$ , and  $\omega$  are functions of  $\Phi$ , these two equations are reduced to

$$\Delta_1 \Phi = -\frac{d(b_\phi^2/2)}{d\Phi} - 4\pi r^2 \left\{ \frac{dq}{d\Phi} \exp \left[ -\left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right] + \rho r^2 \omega \frac{d\omega}{d\Phi} \right\}. \quad (\text{C.23})$$

Another equation to be coupled is the Poisson equation as

$$\Delta \psi = 4\pi G \frac{q}{c_s^2} \exp \left[ -\left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right]. \quad (\text{C.24})$$

The source terms of equations (C.23) and (C.24) are given by determining the mass  $\delta m(\Phi)$  and the angular momentum  $\delta L(\Phi)$  contained in a flux tube  $\Phi - \Phi + \delta\Phi$ . Mass and angular momentum distribution of

$$\delta m(\Phi) = 2 \int_0^{z_s(\Phi)} dz \int_{r(z,\Phi)}^{r(z,\Phi+\delta\Phi)} dr 2\pi r \rho(r, z), \quad (\text{C.25})$$

$$\delta L(\Phi) = 2 \int_0^{z_s(\Phi)} dz \int_{r(z,\Phi)}^{r(z,\Phi+\delta\Phi)} dr 2\pi r \rho(r, z) r^2 \omega(\Phi) \quad (\text{C.26})$$

is chosen arbitrary in nature, where  $z_s(\Phi)$  is the the height of the cloud surface where the magnetic potential is equal to  $\Phi$ . For example,  $m(\Phi)$  and  $L(\Phi)$  are chosen as a uniformly rotating uniform-density spherical cloud threaded by uniform magnetic field. Since  $\int_{r(\Phi)}^{r(\Phi+\delta\Phi)} 2\pi r dr = \delta\Phi \pi \partial r^2 / \partial \Phi$

$$q(\Phi) = (dm/d\Phi) / \int_0^{z_s(\Phi)} dz 2\pi \left( \partial r^2 / \partial \Phi \right) \frac{1}{c_s^2} \exp \left[ -\left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right], \quad (\text{C.27})$$

$$\omega(\Phi) = (dL/d\Phi) / \int_0^{z_s(\Phi)} dz 2\pi \left( \partial r^2 / \partial \Phi \right) \frac{1}{c_s^2} q(\Phi) r^2 \exp \left[ -\left( \psi - \frac{1}{2} r^2 \omega^2 \right) / c_s^2 \right], \quad (\text{C.28})$$

The source terms of PDEs [eqs (C.23) and (C.24)] are given from equations (C.27) and (C.28). While the functions  $q(\Phi)$  and  $\omega(\Phi)$  are determined from the solution of these PDEs after  $m(\Phi)$  and  $L(\Phi)$  are chosen. This can be solved by a self-consistent field method.

## Appendix D

# Basic Equations for Radiative Hydrodynamics

### D.1 Radiative Hydrodynamics

The basic equation for gas which is affected by the radiation is as follows:

$$\frac{d\rho}{dt} + \rho \nabla \cdot \mathbf{v} = 0, \quad (\text{D.1})$$

$$\rho \frac{d\mathbf{v}}{dt} = \rho \nabla \phi - \nabla \psi + \frac{\chi_{F_0}}{c} \mathbf{F}_0, \quad (\text{D.2})$$

$$\rho \frac{d\epsilon}{dt} + p \nabla \cdot \mathbf{v} = h_{\text{CR}} \rho + c \chi_{E_0} E_0 - 4\pi \chi_{P_0} B, \quad (\text{D.3})$$

$$\nabla^2 \psi = 4\pi G \rho, \quad (\text{D.4})$$

$$\epsilon = \frac{p}{(\gamma - 1)\rho}, \quad (\text{D.5})$$

where  $\chi$ ,  $h_{\text{CR}}$ , and  $B$  are the absorption coefficient, the cosmic-ray heating rate per unit mass, and the Planck function  $B = \sigma T^4/\pi$ . In equation (D.2), the term  $\frac{\chi_{F_0}}{c} \mathbf{F}_0$  represents the acceleration of gas due to the photon pressure. In equation (D.3),  $h_{\text{CR}} \rho$ ,  $c \chi_{E_0} E_0$  and  $4\pi \chi_{P_0} B$  represent, respectively the heating due to the CR particles, heating due to the absorption of radiation and cooling due to the emission. The frequency-integrated radiation energy density and radiation flux are defined as

$$E_0 = \int_0^\infty d\nu_0 \int d\Omega I(\nu, \mathbf{n}), \quad (\text{D.6})$$

$$\mathbf{F}_0 = \int_0^\infty d\nu_0 \int d\Omega I(\nu, \mathbf{n}) \mathbf{n}, \quad (\text{D.7})$$

where  $I(\nu, \mathbf{n})$  denotes the specific intensity of radiation at frequency  $\nu$  along the direction vector  $\mathbf{n}$ . These equations could be solved if the radiation transfer is solved. The frequency-averaged absorption coefficients are defined as follows:

$$\chi_{F_0} = \left( \int d\nu \chi(\nu) \mathbf{F}_0(\nu) \right) / \mathbf{F}_0, \quad (\text{D.8})$$

$$\chi_{E_0} = \left( \int d\nu \chi(\nu) E_0(\nu) \right) / E_0, \quad (\text{D.9})$$

$$\chi_{P_0} = \left( \int d\nu \chi(\nu) B(\nu) \right) / B, \quad (\text{D.10})$$

While the equation for the radiation transfer is basically as follows:

$$\frac{1}{c} \frac{\partial I(\nu, \mathbf{r}, \mathbf{n})}{\partial t} + \mathbf{n} \cdot \nabla I(\nu, \mathbf{r}, \mathbf{n}) = -\chi(\nu, \mathbf{r}) [I(\nu, \mathbf{r}, \mathbf{n}) - S(\nu, \mathbf{r})], \quad (\text{D.11})$$

where  $S(\nu, \mathbf{r})$  represent the source term.